with the minimum of sequelae.—We are, etc.,

IAN CRAFT
EZAT YOUSSEFJADIAN

Chelsea Hospital for Women,
London S.W.3

- 1 Wright, C. S. W., Campbell, S., and Beasley, J., *Lancet*, 1972, 1, 1278.
- 2 Craft, I., *Prostaglandins*, 1973, 3, 377.
- 3 Brenner, W. E., Dingfelder, J. R., Staurovsky, L. F., and Hendricks, C. H., *Prostaglandins*, 1973, 4, 819.
- 4 Mocsa, P., and Csapo, A. I., *Lancet*, 1973, 2, 683.
- 5 Skajaa, T., *Acta Obstetrica et Gynecologica Scandinavica*, 1961, 40, 68.
- 6 Bradley-Watson, P. J., Beard, R. J., and Craft, I., *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1973, 80, 284.
- 7 Craft, I., *Lancet*, 1973, 1, 1344.
- 8 Craft, I., *Prostaglandins*, 1973, 4, 755.
- 9 Craft, I., *Prostaglandins*, 1974, 5, 397.
- 10 Bowen-Simpkins, P., *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1973, 80, 824.

Meningococcal Disease

SIR,—In their interesting account of an outbreak of meningococcal disease in Devon (16 March, p. 50) Dr. D. M. Easton and his colleagues raise several points worthy of discussion.

They used a combination of penicillin, sulphonamides, and chloramphenicol in the initial treatment of most cases and suggest that if the low cerebrospinal fluid cell count in three of their cases "indicates only minimal inflammatory response in the meninges penetration of penicillin into the C.S.F. may be poor, so that the inclusion of chloramphenicol which passes well into the C.S.F. might be a beneficial addition to the treatment regimen." In meningococcal infection—especially of the fulminant type where relatively little involvement of the meninges is not uncommon—the major consideration is to deal with the systemic infection. Penicillin alone will do this adequately and, if inflammation of the meninges occurs will rapidly eradicate meningococci at that site without the need for the use of chloramphenicol.

The inclusion of sulphonamides in the treatment of meningococcal meningitis is of less certain help than formerly owing to the presence of resistant meningococci. Of the strains isolated in Scotland and forwarded to this laboratory for testing in 1973, 15% were fully resistant to sulphonamides. What is also of importance when considering prophylaxis is that in addition 51.5% of strains were partially resistant—that is, would not be eradicated from the nasopharynx by sulphonamide therapy. Hence if prophylaxis is to be used sulphonamides cannot be recommended.

This raises the question as to who should be protected and how. Firstly, I cannot agree with Dr. Easton and his colleagues that "family contacts should be screened and should all be given an adequate course of prophylactic treatment" if they imply that prophylaxis should be dependent upon screening. Screening will not pick up all carriers and in any case I would agree with Wenzel *et al.*¹ that it is the acquisition rate of meningococci that is important in terms of the dynamics of meningococcal infection, not the carrier rate. In other words, meningococcal disease is, like poliomyelitis, a "failure of carriage" (and it is interesting to note that in both these diseases asymptomatic infection leads to the production

of antibodies). Hence prophylaxis must be used early if at all and must cover the case contacts, not so much to eradicate carriage but to try to prevent acquisition leading to disease. There is no doubt that prophylaxis will abort epidemics, as has been shown in Africa² and in service camps in the U.S.A.³ but the problem is how this should be carried out in the face of sulphonamide resistance and the knowledge that penicillin will not eradicate meningococci from the nasopharynx of carriers.⁴ One approach might be to give penicillin in the hope of preventing disease (and maybe by reducing carriage rates to a low level to reduce the likelihood of acquisition by the non-carrier). However, there is evidence⁵ that penicillin not only will not eradicate carriage but will not prevent the onset of meningitis. The other approach is to give eradication treatment—for instance, using minocycline and rifampicin⁶ or rifampicin alone.⁷ Until reading the account of Foster *et al.*⁵ I might have conceded that in a family group penicillin, especially if given by injection, should be adequate, but I now feel that, as in an institutional outbreak where long-term carriage following inadequate prophylaxis could result in the seeding of the susceptible population once prophylaxis had been ended, eradication treatment should be considered. It would be interesting to hear the views of others on this important topic.

At one time the suggestion that family contacts should receive prophylaxis would not have been accepted because outbreaks in family groups were rarely reported, but in recent years outbreaks involving more than one member of a family have been reported⁸ and the experience in the Devon outbreak emphasizes that this can happen. Hence the urgency for reappraisal of the problem of prophylaxis is underlined.

Finally, Dr. Easton and his colleagues noted that group B meningococci predominated in their series. Were other groups isolated or were some strains either untyped or untypable?—I am, etc.,

R. J. FALLON

Department of Laboratory Medicine,
Ruchill Hospital,
Glasgow

- 1 Wenzel, R. P., Davies, J. A., Mitzel, J. R., and Beam, W. E., *Lancet*, 1973, 2, 205.
- 2 Vollum, R. L., and Griffiths, P. W. W., *Journal of Clinical Pathology*, 1962, 15, 50.
- 3 Kuhns, D. M., Nelson, C. T., Feldman, H. A., and Kuhn, L. R., *Journal of the American Medical Association*, 1943, 123, 335.
- 4 Dowd, J. M., Blink, D., Miller, C. H., Frank, P. F., and Pierce, W. B., *Journal of Infectious Diseases*, 1966, 116, 473.
- 5 Foster, M. T., jun., Sanders, E., and Ginter, M., *American Journal of Epidemiology*, 1971, 93, 346.
- 6 Devine, L. F., *et al.*, *American Journal of Epidemiology*, 1973, 97, 394.
- 7 Deal, W. B., and Sanders, E., *New England Journal of Medicine*, 1969, 281, 641.
- 8 Greenfield, S., and Feldman, H. A., *New England Journal of Medicine*, 1967, 277, 497.

SIR,—Recent experience in Wigan supports the suggestion of Dr. D. M. Easton and others (16 March, p. 507) that there is a recrudescence of meningococcal disease, often with complications, in Britain. In the six months from October 1973 to March 1974 we treated 10 such patients compared with four in the corresponding period a year earlier. Their ages ranged from 5 months to 7 years. All had confirmed meningitis and in nine the presence of men-

ingococci was demonstrated bacteriologically. A petechial rash was noted in nine cases.

Two children died, a girl of 3 years with severe septicaemia and shock and a boy of 6 who had septicaemia and gross cerebral oedema with coning and superficial cerebellar haemorrhage. Two other children had brief convulsive episodes and a third who had prolonged convulsions went on to develop bilateral subdural effusions and has considerable residual brain damage. One child had definite arthritis of the right elbow joint and another a transient arthralgia affecting the left knee.

None of these children had any history of prior contact with meningococcal disease and their homes were scattered throughout the area served by this hospital.—I am, etc.,

R. B. MCGUCKEN

Paediatric Department,
Royal Albert Edward Infirmary,
Wigan

Misleading Discs

SIR,—The antibiotic discs described by Mr. D. F. J. Brown and Dr. J. B. Selkon (23 March, p. 573) are not the only ones to mislead.

Bacteria of the *Haemophilus* genus require one or both of two growth-promoting substances for normal growth on culture. These are haematin, designated the "X-factor", and coenzyme 1, designated "V-factor". Coloured paper discs marked X, XV, and V, impregnated with standard quantities of these substances, are commonly used as a simple screening method for preliminary recognition of *haemophilus* species. Recently a culture of *H. influenzae* appeared as only a fine ring of growth around an XV disc, but outside a zone 14 mm in diameter without any growth. It was almost invisible without a hand lens. Another *haemophilus* grew as a ring around a zone without growth surrounding both the X and V discs. Experimental cultures of pneumococci and streptococci also showed zones of inhibition around the X and XV discs. The discs were in date and stored dry at 4°C.

A similar phenomenon was observed some years ago. The paper dye was not to blame since control blank discs of the three colours were inactive, and it was assumed that the X-factor (haematin) solution used must have been inhibitory. The makers were informed and corrected the matter, but the difficulty now seems to have recurred.—I am, etc.,

MAIR THOMAS

Pathology Department,
Elizabeth Garrett Anderson Hospital,
London N.W.1

Clinical Diagnosis of Reye's Syndrome

SIR,—I entirely agree with the views expressed in the last paragraph of Dr. Ellen S. Kang's letter (16 March, p. 518). Having an abiding interest in Reye's syndrome^{1,2} may I develop her idea a little farther? I suggest that what she terms "toxic encephalopathy with fatty visceral changes due to a specific toxin" should be called the Reye-Bourgeois syndrome because it was the original intention of Reye *et al.*³ to place the entity of "encephalopathy with fatty degeneration of