exportable, successful, practical, and acceptable. I hope that other centres will wish to try this type of approach for the shared management of this and other conditions requiring long-term follow-up—for example, iron deficiency anaemia, hypertension, and rheumatoid arthritis.

I am most grateful to Professor J. Crooks, Dr. W. D. Alexander, and Professor G. M. Wilson for helpful comments concerning the manu-

script. I should also like to acknowledge the members of the SAFUR organization—in particular Dr. Anthony Hedley and Professor R. Deans Weir—and Miss Margaret Brough for secretarial help and invaluable assistance in the collection and analysis of data.

Reference

- ¹ Hedley, A. J., Scott, A. M., Weir, R. D., and Crooks, J., *British Medical Journal*, 1970, 1, 556.

Medicine in China

Public Health in China

A. J. SMITH

British Medical Journal, 1974, 2, 492-494

Acupuncture and barefoot doctors may be the current talking points in any Western discussion of medicine in China, but in fact probably the most important achievements made in the years of Communist rule have been in the field of public health. Many infectious and parasitic diseases have been eliminated and others brought under control. Sixteen days spent in a country the size of China does not really give time for even a superficial impression; and necessarily much of the information we required came from conversations with our hosts rather than from our own observations. However, a great deal of information on methods and results has been published¹⁻³ in the last few years, and it is true to say that we saw nothing to contradict the claims made by the Chinese health authorities.

Health Priorities

In the 25 years of the People's Republic health policies have changed, sometimes quite abruptly, on matters such as the need for population growth or restraint; but a single theme has retained pride of place—that preventive medicine has priority over curative treatment. With this has been combined the unique Chinese capacity for mobilizing mass efforts by the people—and this was well illustrated by the first Patriotic Health Campaign launched in the early 1950s.

The primary goal of the health campaign was simple—the elimination of the four pests, flies, mosquitoes, rats, and sparrows (later sparrows were replaced as targets by bedbugs as the ecological value of the birds became apparent). Workers and their families were repeatedly encouraged to search for and destroy the pests: big campaigns were held in the spring each year before the national holiday on May Day. The astonishing result has been that the housefly is virtually extinct in China.

We visited Peking, Shanghai, and Kwangchow in late April and early May, with temperatures in the 70s and low 80s F. (25-30°C). We went round pig farms and dairy farms, we visited fish shops and butchers, and we used public latrines near the Great Wall of China in company with

thousands of workers having a day out. No flies were to be seen, until we reached the far south in Kwangchow, where there were some very occasional ones.

The barefoot doctors have been responsible for maintaining the impetus of the patriotic health campaigns at the commune level—and this means keeping the commune clean as well as free from flies and rats. Many communes have three set days each month when the entire population takes part in a scouring, cleaning, disinfecting, and rubbish clearing exercise. In the cities government office workers, who are expected to do some manual labour each week, may well spend that time helping to clear rubbish or clean the streets. Certainly we noticed very little rubbish or waste in the streets of the cities or in the yards of the houses of the agricultural workers: everything had a scrubbed or scoured appearance (though very little recent painting seemed to have been done).

Endemic Disease

In 1949 the bacterial and viral diseases in China included plague, cholera, smallpox, leprosy, poliomyelitis and tuberculosis and the parasitic disorders included malaria, kala azar, hookworm, filariasis, and schistosomiasis. Some of these were localized to only part of the country, but all represented major public health problems. They were tackled by a multiple attack: improvements in sanitation, elimination of intermediate hosts, vaccination and immunization campaigns, and identification and treatment of human carriers. The typical approach of the Chinese health authorities to endemic disease can be illustrated by their programme for the elimination of schistosomiasis, which formerly affected a large area of China along the valley of the Yangtze river—threatening the health of 100,000 people.

Schistosomiasis

The Chinese form of schistosomiasis is caused by *Schistosoma japonicum*, and the intermediate host is the small amphibious snail *Oncomelania hupensis*. The attack on the disease was described in detail for us at the Institute of Parasitic Diseases in Shanghai. Of the eight laboratories in the institute four are concerned with schistosomiasis, and one each of the others with kala azar, malaria, hookworm, and filariasis.

The key to the control programme is elimination of the snails. Here the Chinese have an advantage over many other

parts of the world affected by schistosomiasis: *O. hupensis* is an amphibian, living just above and below the waterline in the irrigation ditches and pounds, and not a truly aquatic snail.

Professor Yu Wei, the director of the preventive research work, explained that in each commune the whole water system had to be mapped and a systematic plan agreed for step-by-step eradication. The amphibious snails could be killed by being buried below the water line. The method used was to drain a ditch of water (fig. 1), dig trenches at the bottom, and then to cut the infected earth from the banks and bury it in the trenches. The snails died within three months of being buried. This radical method of snail-eradication was combined with the use of snail poisons, sprayed on the banks of less heavily infected areas, and added to the water in the paddy-fields in doses calculated to be the minimum necessary. Poisoning alone was not really satisfactory, since the available molluscicides were also toxic either to fish or to mammals.

FIG. 1—Procedure used in irrigation ditches in Yangtze valley to eradicate snails. These are most numerous in the soil just above the water level (1). First the irrigation ditch is drained (2); next trenches are dug at the bottom of the ditch (3). The snail-infected soil is cut away (4) and buried in the trenches (5). The ditch is then refilled with water (6), and the snails are drowned.

Also important was the work done to reduce the sites available for the growth of snails. Pools of stagnant water had to be removed, and the banks of waterways and irrigation ditches tidied (fig. 2) and cleaned so that annual snail-killing programmes could be effective.

Communal Effort

In every commune in the endemic areas, said Professor Yu Wei, a team of barefoot doctors had been trained to carry out the programme. They were shown how to teach the commune workers about the disease—an infected rabbit would be dissected so that everyone could see the late effects of schistosomal infection. Propaganda would be repeated at regular intervals.

The snail-eradication programme was combined with improvements to the sanitation standards of the villages. New latrines had been designed, using a triple chamber storage system which effectively destroyed the parasitic ova before the sewage was used for fertilizer.

FIG. 2—Clean and tidied bank of waterway in South China. Bank improvements like this help to maintain areas free of snails.

The third object of the programme was treatment of the inhabitants of the endemic areas: this required regular screening programmes to pick up those with the disease. Stool tests were satisfactory only when there was heavy infection; as the programme had developed skin tests had been devised (using either circarial antigen or an antigen made from the livers of infected rabbits) which could be administered by barefoot doctors and read within 15 minutes.

Results

In the early 1950s, before the programme began, the infection rate was 30 to 40% in the endemic areas. Among 20-year-olds the rate was nearly 60%. Professor Yu Wei told us that the figure had been cut to 25% by the late 1950s, 6% in 1960, and only 4% in 1965. The eradication campaign had taken 12 to 14 years to eliminate snails completely from an area; but in the clean areas there were now no cases of the disease among children and teenagers. The whole Yangtze valley had not yet been brought under control, however: there were still small patches where the snails had not yet all been destroyed.

Other Parasitic Disease

The Shanghai Institute told us that kala azar had virtually been wiped out. This had been achieved by repeated programmes of spraying against sandflies, treatment of patients, and a decision to kill off domestic dogs (a reservoir of the disease). Certainly we only saw one or two dogs while we were in China—the decision to get rid of them had been based on their role as reservoirs not only of kala azar but also of other diseases such as rabies.

We were told that there is still a form of kala azar among gerbils in the North West of China, but that it does not seem to be pathogenic for man. Nevertheless the institute still regularly sends out mobile teams to check that the disease is indeed extinct.

Malaria is still a problem in the south of China, especially in the semi-tropical jungle areas bordering southeast Asia. Most cases are of the falciparum type, we were told: the former important vector, *Anopheles minimus*, has been largely exterminated and the problem now is *A. balabacensis*. However, the numbers of cases were said to be falling each year as a result of spraying with D.D.T. and treatment/prophylaxis with chloroquine and primaquine.

Hookworm and filariasis are still public health problems, but again the combination of improvements in sanitation combined with mass treatment programmes are said to be bringing them under control.

Bacterial and Viral Diseases

Of the bacterial and viral diseases, plague, cholera, and smallpox are apparently extinct—a view shared by the W.H.O. Immunization and vaccination programmes, which fall within the scope of the barefoot doctors, seem to be followed with enthusiasm. Every health station, in both town and country areas, maintains records of the vaccination status of children in its area. At present the routine range includes DPT vaccine; measles; poliomyelitis (oral attenuated virus); BCG (given by injection or by mouth, the first dose within three days of birth); smallpox (though as in Britain the vaccination rate is falling); and vaccines against Japanese B encephalitis and meningococcal infection.

Screening Procedures

Though there is still a long way to go, China can claim to have the infectious and parasitic diseases under reasonable control—"we are in the phase of consolidation," was the attitude of the public health authorities. They are confident enough of the position to have started work on the next level of preventive medicine, screening tests for the population.

The decentralization of the medical services has marked advantages for screening campaigns—and certain drawbacks. The commune hospital we visited outside Shanghai, for example, had set out to x-ray the chest of every one of its 23,000 inhabitants once a year, and this was clearly a practicable proposition—but it was far from clear how many other communes would be following the same programme.

There does seem to be a nationwide campaign of screening for cervical cancer (which in China is more common than breast cancer, though we could find no national statistics on incidence). Gynaecology departments in hospitals took every possible opportunity to take cervical smears, and on the communes this was another of the responsibilities of the barefoot doctors. It seemed that the reading of the smears was done in the commune hospital laboratory.

Screening programmes for specific diseases have been started in some parts of China—the tests for oesophageal cancer in the northeast and for nasopharyngeal cancer in the south (see *B.M.J.*, 25 May, p. 429) are examples.

Barefoot Doctors the Key

While the health authorities can retain the enthusiasm of the barefoot doctors their preventive policies can be highly effective at the commune level. Good quality sanitation and immunization can achieve a great deal. The use of these keen but relatively unskilled health workers in screening campaigns is a more speculative venture, and the results will be of great interest.

References

- ¹ *Medicine and Public Health in the People's Republic of China*, ed J. R. Quinn. New York, U.S. Department of Health, Education and Welfare, National Institute of Health, 1973.
- ² Sidel, V. M., and Sidel, R., *Serve the People*, Observations on Medicine in the People's Republic of China. New York, Macy Foundation, 1973.
- ³ Horn, J. S., *Away with all Pests*. New York, Monthly Review Press, 1969.

Today's Treatment

Blood and Neoplastic Diseases

Treatment of Iron Deficiency and Iron Overload

J. H. DAGG

British Medical Journal, 1974, 2, 494-497

Iron Deficiency

Current practice in the management of iron-deficiency anaemia can give little cause for satisfaction, since follow-up studies indicate a relapse rate of around 30% in both hospital and general practice series. These disappointing results must reflect either an insufficiently energetic approach to the

identification and removal of the underlying cause, or to inefficiency in supplying replacement iron.

REMOVAL OF UNDERLYING CAUSE

Persistence of the primary cause of negative iron balance will guarantee a relapse after withdrawing iron therapy, and sources of manifest or occult bleeding should be identified, and, where possible, treated medically or surgically. Hormonal therapy for menorrhagia has proved useful and hysterectomy may be indicated near the menopause. Full gastrointestinal investigation is justifiable in all men and postmenopausal women (and in premenopausal women without menorrhagia)

Department of Medicine, University of Glasgow

J. H. DAGG, M.D., F.R.C.P.ED., F.R.C.P. Glasg., Consultant Physician