CORRESPONDENCE

Correspondents are asked to be brief

Isolation System for General Hospitals N. R. Grist, F.R.C.PATH.....	Encroachments on the Patients' Responsibility R. F. N. Duke, F.R.C.S.....	Complication of Proximal Gastric Vagotomy J. H. Wyllie, F.R.C.S.....
Tetracycline Poisoning in Renal Failure J. I. W. Davies, M.B.....	Acute Brucellosis Presenting as Epididymo-orchitis C. J. Mitchell, M.B., and T. J. Huins, M.B....	Drugs for Gastric Ulceration Surgeon-Captain T. L. Cleave, M.R.C.P....
Beta-blockade in the Presence of Renal Disease and Hypertension F. D. Thompson, M.R.C.P., and A. M. Joeke, F.R.C.P.	T.V. Programme on Heart Disease J. Yudkin, M.D.....	Pregnancy Associated with Copper-containing I.U.D. V. Sivanesaratnam, M.R.C.O.G., and others..
Autoimmune Haemolytic Anaemia in Ulcerative Colitis A. H. Goldstone, M.R.C.P., and B. Tolliday, F.I.M.L.T.....	Cephaloridine and Gentamicin in Prophylaxis of Surgical Wound Infection I. L. Rosenberg, F.R.C.S., and A. V. Pollock, F.R.C.S.	Clinical Public Health Doctors C. S. Smith, M.B., F.F.C.M.....
Urethral Prolapse in Children C. P. Douglas, F.R.C.O.G.....	Referees and Rejects R. J. V. Pulvertaft, F.R.C.P.....	Working Hours of Junior Staff J. G. Dumoulin, F.R.C.O.G., and others.....
Photosensitivity to Pyrimethamine? S. A. Craven, B.M.....	Epidemic Neuromyasthenia M. J. Dillon, M.R.C.P., and others.....	Consultant Crisis P. J. E. Wilson, F.R.C.S.....
Fenticlor, Actinic Reticuloid, and Antihistamines J. N. Burry, M.R.C.P.ED.....	Primary Responses to Transplantation Antigens G. D. Bonnard, M.D., and others.....	Finance and the Health Service J. C. Scott, F.R.C.S.; S. R. Brennan, M.R.C.P. 562
Looking after Schizophrenics W. W. Sargent, F.R.C.P.....	Sudden Death in Infancy J. A. J. Ferris, M.D., and S. R. Kendeel....	Practice Expenses R. B. L. Ridge, M.B.....
Confidentiality in Medicine J. S. Taylor, M.B.....	Skin Reactions to Practolol P. Wright, F.R.C.S.....	G.P.s' Pay R. J. Mountain, M.B.....
Screening—G.P. or Family Planning Doctor? N. A. Chisholm, L.R.C.P.ED.....	Vitamin A, Pregnancy, and Oral Contraceptives Isabel Gal, M.D.....	Mileage for Consultants G. S. Turner, M.B.....
		G.P.s and Family Planning A. Craig, M.B.....
		Authentication of International Certificates G. C. Mathers, M.B.....
		Minors and Contraception J. D. Beale, M.R.C.G.P.....

Isolation System for General Hospitals

SIR,—Infections of low communicability can be contained relatively easily by techniques designed to limit contact transfer. In planning for the future, however, it is important not to make facile and inappropriate application of historic experience. In earlier years poliomyelitis appeared to be poorly communicable—case - to - case transmission and more than one case per family were considered exceptional when most of the population was immune as a result of clinically trivial infection in early childhood. In present conditions, unless susceptible populations have been protected by vaccination, transmission routes may be more obvious and I have personally seen one family in which four members were afflicted by paralytic poliomyelitis (fatal in one child) during a village outbreak in the Netherlands.¹ The

analogy with hepatitis is clear. If social and hygienic conditions continue to improve we may not in future be able to get away with arrangements which appeared adequate in the past, particularly in the vulnerable populations concentrated in hospitals.

Dr. H. G. Easton (11 May, p. 331) may rest assured that the selective, economical, and efficient use of special facilities and limited resources in this field is the current concern of expert individuals, committees, and national and international conferences.²—I am, etc.,

NORMAN R. GRIST

University Department of Infectious Diseases,
Ruchill Hospital,
Glasgow

¹ Bijkerk, H., et al., *Nederlandsch tijdschrift voor geneeskunde*, 1972, 116, 549.
² Grist, N. R., *Lancet*, 1973, 2, 1340.

Tetracycline Poisoning in Renal Failure

SIR,—Dr. M. E. Phillips and his colleagues (20 April, p. 149) point out that some of the tetracyclines produce a fall in the glomerular filtration rate and can be poisonous to patients with renal impairment. Then the doctor who gives the drug may become the object of medicolegal proceedings. This unhappy situation can be avoided, they say, by prescribing doxycycline, which does not cause a rise in blood urea.

I accept entirely that this is a wise and timely paper. But there are three ways in which a diligent and conscientious general practitioner might fall into error:

(1) Renal impairment is likely to be symptomless and the patient often does not

understand the nature of his illness. The diagnosis can be missed, particularly if the clinical situation is dominated by an infection elsewhere.

(2) Proteinuria is not unknown in febrile illness. Such an illness is often revealed to the practitioner during anti-social hours when the chances of getting biochemical investigations done are not great.

(3) The general practitioner is urged by the Department of Health to avoid using expensive preparations. The present price of doxycycline (Vibramycin) is £26 per 100 tablets and that of oxytetracycline £1.17. It is thus unlikely that he will carry doxycycline in his emergency bag and make it

first choice for patients with respiratory infections in whom only the possibility of renal impairment exists.—I am, etc.,

J. I. W. DAVIES

Northampton

Beta-blockade in the Presence of Renal Disease and Hypertension

SIR,—It was with some surprise that we read the conclusion reached by Dr. D. J. Warren and his colleagues (27 April, p. 193) on the basis of observations on three patients with hypertension and renal failure that beta-blocking drugs "should not be given to patients with moderately severe renal failure." No mention is made of the number of patients treated in whom a diminution of renal function was not seen. In the three cases quoted beta-blockade was on no occasion the only therapy. Two of the patients received propranolol and one oxprenolol. If one were to draw the obvious conclusion from the opinion stated one would have to infer that beta-blockade was in itself a dangerous form of therapy.

During the past three years we have treated 69 patients for a minimum of six months with propranolol alone. Despite adequate control of the blood pressure, in no case has there been a fall in renal function such as to cause a change in the patient's clinical condition. In the accompanying table the mean findings before treatment with propranolol (initial) and at the last follow-up (final) are given for the group as a whole and also subdivided according to their initial creatinine clearance (I.C.C.).

From these results it is apparent that no

Group	No. of Patients	Mean Creatinine Clearance (ml/min)		Mean Blood Pressure (Lying) (mm Hg)	
		Initial	Final	Initial	Final
I.C.C. <30 ml/min	18	15.2	21.3	187/119	155/90
I.C.C. 31-60 ml/min	14	45.7	47.6	194/122	157/93
I.C.C. >60 ml/min	37	102.5	93.8	180/116	140/86
All patients	69	72.3	69.2	185/118	148/88

intrinsically harmful effect of beta-blockade on renal function can be inferred. It is the general experience of any unit treating hypertensive patients with renal functional impairment that transient reduction in renal function may occur during the early stages of effective blood pressure control. This usually returns to the same level or possibly increases above the initial value. The increased peripheral resistance that can occur with beta-blockade might be shared by the renal vasculature. Should reduction in renal function occur after the institution of beta-blockade this may be an indication for the addition of a peripheral dilator such as hydralazine.

We intend to describe fully in future publications our experience of the treatment of hypertension in the presence of renal disease using propranolol. This will substantiate our earlier findings¹ that there is no indication for a reduction of propranolol dosage with reduced renal function.—We are, etc.,

F. D. THOMPSON

Institute of Urology

A. M. JOEKES

St. Peter's Group of Hospitals,
London W.C.2

¹ Thompson, F. D., Joekes, A. M., and Foulkes, D. M., *British Medical Journal*, 1972, 2, 434.

Autoimmune Haemolytic Anaemia in Ulcerative Colitis

SIR,—We would like to report an interesting case of autoimmune haemolytic anaemia with an antibody showing rhesus specificity in a patient with relapsing ulcerative colitis. The anaemia apparently responded to steroid treatment of the colitis.

The patient, a 23-year-old nulliparous woman known to have colitis, was admitted on the present occasion with an exacerbation. Her haemoglobin concentration was 9.8 g/100 ml, mean corpuscular volume 119 pg, and reticulocyte count 7%. The blood film showed many spherocytes and the direct antiglobulin (Coombs) test was strongly positive. Serum folate (1.6 ng/ml) and red cell folate (131 ng/ml) levels were both low. She was treated with prednisone and folic acid.

The positive antiglobulin test showed IgG specificity and her serum contained antibody of mixed specificity, but an eluate of her red cells contained an antibody with anti- $\bar{e}$ specificity. On treatment the haemoglobin rose to normal, the reticulocyte count fell to 3%, and the antiglobulin test became negative. The serum, however, still contained antibody now appearing to have some anti- $\bar{e}$ specificity. Her probable rhesus genotype was cde/cde and this is consistent with the antibody being an auto-antibody.

There are several reports in the literature¹⁻⁵ describing the presence of positive direct antiglobulin tests in a total of eight patients with ulcerative colitis. Only a few

of these showed active haemolysis and only two of them showed any degree of rhesus specificity (in one patient an anti- $\bar{e}$ plus f from a red cell eluate and in another patient an anti- $\bar{e}$ in the serum). It is important to report such cases, which may help to elucidate the "aetiology and pathogenesis of this autoimmune haemolytic anaemia which does not have to be classified as idiopathic."⁶

We would like to thank Dr. Sheila Worledge for helpful discussion about this patient and Dr. T. M. Chalmers for permission to report the case.

—We are, etc.,

A. H. GOLDSTONE

Department of Haematology,
Addenbrooke's Hospital,
Cambridge

¹ Lorber, M., Schwartz, L. I., and Wasserman, L. R., *American Journal of Medicine*, 1955, 19, 887.

² Balint, J. A., Hammack, W. J., and Patton, T. B., *Gastroenterology*, 1962, 42, 740.

³ Fong, S., Fudenberg, H., and Perlmann, P., *Vox Sanguinis*, 1963, 8, 668.

⁴ Gastaldi, F., et al., *Rivista di Emoterapia ed Immunoematologia*, 1963, 10, 193.

⁵ Molaro, G. L., and Barillari, B., *Rivista di Emoterapia ed Immunoematologia*, 1964, 11, 43.

⁶ Dacie, J. V., *The Haemolytic Anaemias*, Part III, p. 826. London, Churchill, 1967.

Urethral Prolapse in Children

SIR,—Your leading article (4 May, p. 240) perhaps makes this condition sound more complicated than it really is. From an experience of some 25 patients under the age of 8 may I suggest two points? (1) The diagnosis is usually easily made by passing a urethral catheter—prolapse is the only tumour which symmetrically surrounds the catheter. (2) Treatment is best done as an outpatient, avoiding the psychogenic trauma of separation in a frightened young child. Under general anaesthesia (so rapidly eliminated today) two stitches (Dexon (polyglycolic acid) may be best) are inserted through normal urethra at right angles to one another. The prolapsed tissue distal to the sutures is then excised. The "cross" is then picked up with forceps and converted into four stitches, which can be tied with no fear of retraction of the mucosa.

I believe it is unwise to leave a catheter in the bladder as this only leads to infection. I have not found it necessary to catheterize subsequently for retention and I am not aware of urethral stricture developing subsequently (though my follow-up is not complete).—I am, etc.,

CHARLES P. DOUGLAS

Royal Free Hospital,
Liverpool Road Branch,
London N.1

Photosensitivity to Pyrimethamine

SIR,—I wish to report an apparent photosensitivity reaction to pyrimethamine (Daraprim). The Committee on Safety of Medicines has had only one similar notification, in a male of unknown age who was given a course of pyrimethamine. He de-

veloped a photosensitivity rash 10 days after taking his first tablet.

My patient was a 9-year-old boy of English parents who arrived by air in Lagos during March 1974. At 23.00 hr that day he was given one 25-mg Daraprim tablet. At 17.00 hr the following day he developed a pink, confluent, macular, non-pruritic rash over his cheeks, forearms, and legs which corresponded exactly with those areas exposed to the sun. There was no malaise. The rash disappeared overnight but recurred on the third day following exposure to the sun. It again disappeared overnight and reappeared on the fourth day following exposure to the sun. On the fifth day there was no rash. He had taken pyrimethamine on many occasions previously, though this tablet was the first of his present course. He had never before had a skin rash and had no known drug or other sensitivities. There was no family history of drug or other sensitivities. He was taking no other drugs.

Wellcome-Lepetit (Nigeria) Ltd. analysed the remaining tablets from the packet and tablets from their quality-control stock and were unable to detect any deviation from their standard product.—I am, etc.,

STEPHEN A. CRAVEN

Estate Clinic,
Ikeja, Nigeria

Fenticlor, Actinic Reticuloid, and Antihistamines

SIR,—In the article "Diseases of the Skin: Acne Vulgaris" (15 December 1973, p. 667), Dr. W. J. Cunliffe suggests fenticlor as a suitable topical antimicrobial agent in treatment. Fenticlor causes photocontact dermatitis¹ and two cases of chronic dermatitis from persistent photosensitivity² have followed its incorporation into a hair cream in the United Kingdom. Except for the absence of two chlorine molecules, fenticlor is identical with bithionol,³ known to cause distressing persistent photodermatitis which forces the sufferer to live a confined indoor life.⁴ Fenticlor and bithionol cannot be imported into Australia without Commonwealth authority, and the use of bithionol in consumer goods in the U.S.A. and Canada is restricted by government legislation.

The term "actinic reticuloid" describes a chronic dermatosis associated with severe photosensitivity and histological resemblance to lymphoma. Ives *et al.*⁵ in the original report of this condition suggest a relationship with the chronic photodermatitis caused by photoallergens such as bithionol and fenticlor. Wilkinson⁶ described a case of actinic reticuloid which first developed in 1969 but in which sensitivity to tetrachlorosalicylanilide, a halogenated salicylanilide related to fenticlor and bithionol, had been diagnosed in 1960. A direct causal relationship between actinic reticuloid and contact photoallergens has not been established, though such a relationship must be suspected. Sneddon⁷ described a case of actinic reticuloid which may have developed into a true lymphoma.

It has been proposed that persistent light reactions following the use of contact photoallergens are caused by the retention of those photoallergens in the dermal layer of skin for long periods.⁸ This concept might con-