ing the emphasis of the medical course away from the technically minded, curatively oriented Western style towards the preventive approach based on simple techniques in which the rural worker is regarded as being at the top of the pyramid of care rather than at the bottom.

The disparaging remarks which Dr. Senewiratne makes about medical auxiliaries do not accord well with experience in China¹ and Africa, where it is being found that a simply trained medical worker in close contact with the people and under the direction of doctors can improve basic health more effectively and much more cheaply than under a conventional Western-type system—as well as making the doctor's work, in organizing a team approach, more rewarding.

It is inevitable that medical training in a developing country patterned on Western methods will encourage migration, as will postgraduate courses in Europe and America. As Dr. Senewiratne points out, much insidious harm is done by such courses, which could be avoided if the teachers were instead seconded to the country concerned for a period. This would also have the valuable effect of widening the participant's viewpoint immeasurably.—I am, etc.,

TONY WATERSTON

Royal Hospital for Sick Children,
Yorkhill, Glasgow

¹ *Health Care in China*. Geneva, Christian Medical Commission, 1974.

Complications of Laparoscopy

SIR,—Though a straightforward laparoscopic sterilization is probably a shorter procedure than a tubal ligation after laparotomy, the advantage is marginal and if there is an increased incidence of complications laparoscopy would be unjustifiable. However, the shorter time taken for laparoscopic sterilization is not the chief justification for its use and Mr. E. E. Rawlings and Mr. B. Balgobin (29 March, p. 727), by keeping such patients in hospital for three to four days, nullify the main advantage.

Several consultants in this unit consider this procedure to be suitable for day cases, though the patients always have the option to choose to be kept in overnight or to change their mind after recovering from the anaesthetic. The table shows the number of patients admitted for laparoscopic sterilization as day cases and the number who were transferred for overnight recovery during the past year, since the day area in this hos-

pital opened. The dramatic change in August reflected the increasing confidence of the sister in charge of the unit, who was not previously experienced in looking after such cases.

Since August 1974 83 patients have been operated on in this way and, excluding the two laparotomies, have required 89 bed days. The same number of patients treated by Mr. Rawlings and Mr. Balgobin would have required 283 bed days. The saving is even greater since the day area is staffed by nurses working only between 8 a.m. and 5 p.m. No complications of any consequence have occurred in this small number of cases. The calibre of anaesthesia is obviously of great importance. Most of the patients have been given follow-up appointments and there is no doubt that this approach is popular with patients, who so often have young children.—I am, etc.,

J. W. CRAWFORD

University Department of Obstetrics and Gynaecology,
Ninewells Hospital,
Dundee

Injudicious First-aid

SIR,—We are deeply concerned by the implications of your leading article (5 April, p. 5). We wonder if the State-registered nurse who gave external cardiac massage to a shopper who had collapsed was criticized unjustly after the pathologist found internal laceration but "no underlying disease . . . which would have caused her collapse."

Every experienced pathologist is well aware that no definite cause of death can be found in a proportion of patients who die suddenly. If widespread coronary atheroma is present this may reasonably be presumed to be the cause of death; a very recent infarct will not be evident unless special stains are used, and many patients die of ventricular fibrillation in the absence of any fresh ischaemic event. Units providing resuscitation facilities outside hospital can testify to the frequency of primary ventricular fibrillation without apparent cause. The data from Seattle¹ are of special interest. Of 143 patients who survived out-of-hospital fibrillation, 48 had no previous history of cardiovascular disease and no fewer than 97 had no evidence of infarction on serial electrocardiograms. Moreover, we know of many conditions that can cause ventricular fibrillation, such as Wolff-Parkinson-White syndrome complicated by atrial fibrillation and the syndromes of prolonged Q-T interval, which would show no abnormality at necropsy. Thus it would be wrong to assume that the collapse of the patient mentioned in your leading article could not have been due to cardiac arrest simply because the pathologist found no evidence of underlying disease.

We believe that tissues which have become engorged and relatively stiff after circulatory arrest are more susceptible to damage by trauma. We wonder if the injuries described by the pathologist could have occurred because the patient was already dead when the massage was given, rather than the patient having died because of the injuries.

External cardiac massage is not free from risks even in experienced hands. We strongly support the notion that only skilled first-aid workers should be instructed in its use. However, the risks are small compared with the successes that can be achieved. We might

mention that 16 patients survived in our area in 1974 after resuscitation outside hospital by paramedical personnel. We very much hope that nurses, ambulancemen, and other first-aid workers are not exposing themselves to risk of censure by attempting resuscitation, for if they were to be discouraged in this way many lives might be lost unnecessarily.—We are, etc.,

D. A. CHAMBERLAIN
R. I. K. ELLIOTT
D. H. MELCHER

Royal Sussex County Hospital,
Brighton

¹ Baum, R. S., Alvarez, H., and Cobb, L. A., *Circulation*, 1974, 50, 1231.

SIR,—Death due to injudicious first-aid, as described in your leading article (5 April, p. 5), must be a rare event. I am not sure that injury due to first-aid is uncommon. I have certainly seen in the past two years two patients to whom this happened. The first was a man who was on holiday in East Anglia with his wife. He is a man subject to attacks of hypoglycaemia and on this occasion he left his hotel before breakfast to post a letter. He felt faint while waiting to cross the road. A passer-by was a nurse from an intensive care unit. She assumed cardiac arrest and immediately applied heart compression, as a result of which the patient was in hospital with a fractured sternum and several fractured ribs for several weeks. Recently a patient of mine, subject to faints, had an attack in church. On this occasion artificial respiration fractured several ribs. The warnings in your article should be heeded by all.—I am, etc.,

M. DRURY

Department of Medicine,
Queen Elizabeth Hospital,
Edgbaston, Birmingham

Hormonal Pregnancy Tests and Congenital Malformations

SIR,—Your leading article (30 November, p. 485) reviewed the evidence for and against a teratogenic effect of hormones administered to women during early pregnancy and concluded that the findings of associations between sex steroids and congenital anomalies "require confirmation or refutation from elsewhere."

The Committee on Safety of Medicines has received through its spontaneous reporting scheme only a small number of reports alleging a possible causal association between the use of drugs during pregnancy and the subsequent delivery of a malformed child. However, in order to detect possible associations the committee has collaborated with the medical division of the Office of Population Censuses and Surveys (O.P.C.S.) in an investigation of pregnancies which resulted in the birth of a malformed child in England and Wales during 1971 and 1972. This investigation has been based on notifications to the O.P.C.S. of babies born with a malformation. General practitioners who had cared for the mothers during pregnancy were identified with the help of family practitioner committees. Details of the maternal "drug histories" were obtained from the general practitioners' records during interviews with doctors employed by the Committee on Safety of Medicines. Each case history was paired with that of a normal

	Day Cases	Overnight Stay
1974		
March/April	16	10*
May	27	10
June	19	6
July	18	9†
Total	80	35
August	23	1†
September	11	0
October	5	1†
November	6	2
December	8	1
1975		
January	10	2
February	9	—
March	11	1
Total	83	8

*Laparotomy and sterilization carried out in two cases.
†Laparotomy and sterilization carried out in one case.