can be more perfectly enjoyed, but the import of sterile water from the U.S.A. for cystoscopic purposes is surely the height of extravagance. That our administrators can condone this and other expensive gimmicks such as disposable sigmoidoscopes shows little thought for economy. I hope Mr. Shaldon's letter will provoke a flood of similar criticism which in the fullness of time may put an end to this administrative folly.—I am, etc.,

I. E. W. GILMOUR

Leith Hospital,
Edinburgh

Dangers of Silent Gall Stones

SIR,—In your leading article (22 February, p. 415) you advocate elective cholecystectomy whenever gall stones are discovered. The evidence in the literature, as recently reviewed,¹ and the argument in your article are far from conclusive. The logic underlying a decision to advise cholecystectomy for a symptomless condition must depend on the incidence of complications, the possibility of developing carcinoma of the gall bladder, and the mortality of elective cholecystectomy. The most detailed study² (with which other studies³⁻⁵ agree) on the course of symptomless gall stones showed that 49% developed symptoms but only two-thirds of these had trouble sufficient to merit cholecystectomy. Further, the incidence of major complications was 18%, and twice as frequent in the group who had originally non-functioning gall bladders on cholecystography. The mortality of the group with silent stones was similar to that of the study as a whole, suggesting that elective cholecystectomy did not reduce mortality. It has been shown that patients with silent stones tend to develop symptoms within five years,⁶ not as suggested in your article in a linear fashion.

The danger of a conservative policy in the management of silent stones is that there will be an increased incidence of acute cholecystitis, particularly in the elderly age groups,⁷ but Sato and Matsushiro⁷ found that 74% of patients who were admitted with acute cholecystitis had had previous symptoms and only 6.5% of their patients who had had a cholecystectomy had been admitted in the acute phase. Thus the occurrence of acute cholecystitis as a complication of the conservative management of silent stones is unlikely to be a major contraindication to this policy.

Despite the well-known association between gall stones and carcinoma of the gall bladder there is no evidence that elective cholecystectomy will reduce mortality.¹ It has been estimated that the deaths from cholecystectomy must be reduced to less than 0.1% to achieve a reduction in overall mortality from carcinoma of the gall bladder, and though mortality figures as low as 0.3% are recorded,⁸ the general figure may be as high as 3.5%.⁹

Thus the case against elective cholecystectomy for silent stones could be argued on the basis that 50% of subjects will be symptom-free for life, cancer of the gall bladder is most unlikely to be a significant complication, and the mortality is similar whichever course is pursued. A more logical approach to this problem is to advise cholecystectomy for those patients who on detailed questioning are found to have

symptoms referable to the gall bladder and for those who have a non-functioning gall bladder at cholecystography. The remaining patients should be told to report to their doctor at the onset of symptoms, and elective cholecystectomy should then be performed at the earliest opportunity to avoid increased morbidity and mortality from acute cholecystitis.—We are, etc.,

R. C. G. RUSSELL
HUGH DUDLEYSurgical Unit,
St. Mary's Hospital Medical School,
London W.2

- 1 Peskin, G. W., *Surgical Clinics of North America*, 1973, 53, 1063.
- 2 Wenckert, A., and Robertson, B., *Gastroenterology*, 1966, 50, 376.
- 3 Comfort, M. W., Gray, H. K., and Wilson, J. M., *Annals of Surgery*, 1948, 128, 931.
- 4 Newman, H. F., and Northup, J. D., *International Abstracts of Surgery*, 1959, 109, 1.
- 5 Spiro, H. M., *Clinical Gastroenterology*, p. 739. Toronto, Macmillan, 1970.
- 6 Lund, J., *Annals of Surgery*, 1960, 151, 153.
- 7 Sato, T., and Matsushiro, T., *American Journal of Surgery*, 1974, 128, 368.
- 8 Glenn, F., and McSherry, C. K., *Annals of Surgery*, 1963, 157, 695.
- 9 Arnold, D. J., et al., *American Journal of Surgery*, 1970, 119, 714.

Merrison Report and Asian Doctors

SIR,—The Merrison Report¹ has made most Asian doctors angry. When the N.H.S. was in danger of collapse Asian doctors were not only welcome but were encouraged to come and work here. Now they are being humiliated in a general attack, though a big majority has faithfully and efficiently served the N.H.S. Britain has a right to impose conditions on new entrants, but it is reprehensible to kick the established ones in the teeth. Would it not be kinder and more honest to halt immigration of all overseas doctors than to allow them to come and then submit them to indignity and humiliation?—I am, etc.,

D. R. PREM
President,
Standing Conference of Asian Organizations
in the U.K.

Birmingham

¹ Report of the Committee of Inquiry into the Regulation of the Medical Profession, Cmnd. 6018. London, H.M.S.O.

Medical Problems of the North Sea

SIR,—Your leading article (5 April, p. 5) on medical implications of oil related industry was both timely and interesting. It appeared, however, by concentrating on the problems of the northern North Sea, to ignore areas south of Yorkshire as far as Britain is concerned and failed to comment on the fact that similar problems are faced by our colleagues in Scandinavia and Europe. While it is probably true to say that the sea is deeper and the climatic conditions worse in the north of the North Sea than in the southern area, I think it was a pity you failed to stress the importance of all medical groups who face the problems of the North Sea sharing their information and experience. I am sure in this way we are most likely to be of help to those who work in the North Sea.

My colleagues and I, who have worked in the southern North Sea for over eight years, are keen to learn from the experience of other groups and also to make available any useful information that we may have gathered in the course of this time. While

I am proud of my forebears from Aberdeen, I would hate to think that your leading article suggests that the problems of offshore medicine in the North Sea are confined to those on the coasts of Scotland.—I am, etc.,

I. K. ANDERSON

Royal Norfolk and Suffolk Yacht Club,
Lowestoft

Tailored Treatment for Varicose Veins

SIR,—I would like to agree with the spirit, if not the letter, of your leading article (15 March, p. 593). The principle you enunciate is one I have accepted for some years now—namely, that no single treatment suits all patients with varicose veins. This surely must be the lesson of the post-war stripping era and, as your article makes clear, evidence is now coming to light to condemn sclerotherapy as an all-or-none cure. There have always been reservations in my mind that Fegan's simplistic approach has been the logical or desirable treatment for varicose veins; I cannot accept, for instance, that all varicose veins are associated with incompetent perforators, nor can I accept that veins become varicose secondarily—that is, from below up—when patients will describe a distinct process by which veins appear proximally and spread distally. However, I would agree that Fegan has made a considerable contribution to the treatment of this condition by deflecting us from the equally illogical regimen of routine stripping of varicose veins; if Fegan had not existed, it would have been necessary to invent him. Stripping was a failure because so often it removed a normal long saphenous vein leaving the incompetent tributaries, and this is where I disagree with the detail of your article, where you advocate limited stripping from groin to knee, for I fear this will result in similar disappointment. Moreover, being a peripheral-artery surgeon, I am reluctant to remove the long saphenous vein, particularly in a male, in whom I know that this is the best substitute for an atheromatous superficial femoral artery at present available. The main trunk is affected far less often than was thought, and even if there are minor defects it can still be used for arterial grafting with appropriate repair.

Logical treatment of varicose veins is based on the following. (1) No injection treatment can be practised proximal to the knee. (2) If sclerosant therapy is indicated evidence should be sought for incompetence at the saphenofemoral junction, as it has been shown that if this is present the early recurrence rate is higher than if there is no incompetence at this point.¹ It is not easy to establish clinically whether there is saphenofemoral incompetence or not. A palpable thrill or a saphena varix, while constituting positive clinical proof, may be absent in many cases when functional incompetence at this junction is present. Therefore, as a policy, one must be prepared to explore the groin more often, while not necessarily ligating any veins unless they are obviously incompetent—that is, a proportion of negative explorations must be anticipated. (3) The best way to remove the obvious varicosities is a direct attack on them. It has always been a source of frustration that simple ligation, even if done at multiple