natal environment (due to difficulty in replacing passive with active immunity).

Sudden and unexplained deaths of infants are often preceded by easy deliveries or a short second stage of labour.³ So it is possible that cot deaths and unexplained stillbirths have a common aetiology, which should be explored by a combination of biochemical and histological tests. Also Drs. Tapp and Otridge have discovered an excess of plasma cells in the marrow of infants whose sudden and unexplained deaths were preceded by minor respiratory infections. So it is possible that infants who have a congenitally defective immune system have more difficulty in producing an effective leucocytosis (following infections) than in producing extra plasma cells.—I am, etc.,

A. M. STEWART

Regional Cancer Registry,
Queen Elizabeth Medical Centre,
Birmingham

1 Stewart, A. M., *British Journal of Cancer*, 1973, 27, 465.

2 Stewart, A. M., *British Medical Journal*, 1974, 2, 611.

3 Proestos, C. D., et al., *Archives of Disease in Childhood*, 1973, 48, 835.

Genitourinary Medicine

SIR,—I was sorry to see in your leading article (11 January, p. 51) that the Department of Health and Social Security and the Royal College of Physicians have been advised to recommend that the specialty of venereology should in future be called "genitourinary medicine." Though venereologists do deal with conditions of the lower urinary tract, as it is inseparably linked with the genitalia, they should not presume to look after its upper part with its very different functions which are more the province of urologists and nephrologists, who have rightly protested in your columns against this intrusion. The discipline, layout, and objects of our clinics do not lend themselves to taking on this branch of medicine as well as the venereal diseases. At one time the treatment of venereal diseases was in the hands of dermatologists, gynaecologists, urologists, and surgeons (and still is in many countries) to its detriment, as it had become a neglected and unimportant, though lucrative, sideline of these specialties. In Britain we have succeeded in making the management of these diseases a separate specialty of its own so that they could be dealt with by experts in that subject only. It seems to be a retrograde step to link them now with another quite different branch of medicine, which may well in due course take priority over the venereal or "sexually transmitted" diseases and lead to their neglect.

The lack of recruits with higher qualifications to this specialty is said to be due to its limited academic scope. Rather than try to expand this scope by taking in other fields of medicine, would it not be better to modify the academic qualifications required so as to attract those clinicians who are more interested in the art of trying to help with the personal difficulties of the patient rather than in academic medicine only? The actual cure of most of the venereal diseases has become very easy since the advent of antibiotics and chemotherapy, but the patients' fears and worries remain much the same. In helping with the latter the venereologists can be of more real

value to the patient than if he branches out into another area of medicine already fully covered by others.—I am, etc.,

JAMES JEFFERISS

The Praed Street Clinic,
St. Mary's Hospital, London W.2

Uterine Rupture in Labour

SIR,—Experience of rupture of the uterus in Wolverhampton confirms that of Mr. M. Brudenell and Mr. S. Chakravarti (19 April, p. 122). During the past 12 years 23 of our patients sustained uterine rupture, an incidence of 1 in 2048. Of these cases, only two occurred in late pregnancy, both with classical scars. Four lower-segment scars ruptured in labour and only one fetus survived. Traumatic rupture occurred 11 times, due to internal version in four cases and to difficult forceps delivery in the remainder. Spontaneous rupture occurred six times and in three of these cases there were obvious predisposing causes—that is, severe congenital abnormality, previous uterine perforation, and previous manual removals of adherent placentae.

In this small series, with a maternal mortality of one patient and a fetal mortality of only 70%, at least half of the cases occurred during the last four years. Amniotomy was performed on 15 patients, nine of whom received oxytocin in addition. The Cardiff infusion system, introduced in the last three years, was not involved. Epidural analgesia was given to four patients, of whom three were grand multiparae and the fourth was a gravida-5 in whom the analgesia abolished pain caused by the rupturing uterus. It is relevant, however, that only seven patients had severe pain, while serious haemorrhage or shock were obvious features in 13. We now regard grand multiparity as a contraindication to epidural analgesia, though in general we favour its use, especially in

breech birth, where it allows the obstetrician to assist delivery painlessly, without resort to general anaesthesia. The incidence of breech extraction has not been increased. Lower-segment scars are a contraindication to epidural analgesia in labour, but if this method of pain relief is considered essential, then continuous monitoring of the uterine contractions as well as the fetal heart rate is of greater importance than intermittent palpation of the lower-segment scar.

We have also found that the diagnosis is difficult. In one case laparotomy on the third day to perform sterilization revealed dehiscence in the upper segment in a patient who had had her seventh baby after a labour of two hours. The only symptom that she had had was a little pain associated with a little uterine tenderness and abdominal distension. The rent was firmly plugged by omentum and the cause was obviously uterine perforation, documented as occurring 12 months previously.

In our series hysterectomy for rupture of the uterus entailed an average of 2.7 l of blood transfused, while repair required only 1.8 l. One ureter divided during hysterectomy was successfully reimplanted and the overall morbidity of hysterectomy was greater than that of uterine repair.

I would disagree that rupture of the uterus always means loss of the mother's capacity to have further children. Ten of our patients had the ruptured uterus repaired. Six were not sterilized and four of these have had six living babies delivered by caesarean section. Our experience, however, leads me to agree with Mr. Brudenell and Mr. Chakravarti that amniotomy, oxytocin infusion, and epidural analgesia, so helpful to many patients, can become hazardous procedures in those with a weak or damaged uterus or who are grand multigravidae.—I am, etc.,

ALAN M. SMITH

New Cross Hospital,
Wolverhampton

B.M.A.: Need for Radical Change

SIR,—In some intangible way the B.M.A. is felt to be failing its members. The membership is reported to be gradually falling away, especially on the hospital side, and to be perilously near the significant 50% figure. This must be largely due to its often praise-worthy but ineffectual attempts to cover all fields: to be trade union, employment agency, expert spokesman on all medical matters, educator, charity, and even, at local level, organizer of social functions. That it fails in many of the more important of these activities is inevitable, given this brief.

The time has come for radical change, and what more appropriate time than now, when our long-serving Secretary, Dr. Derek Stevenson, is within a year of retirement? There have been disquieting rumours that the succession has already been decided within the upper echelons.

There can be little doubt that what members need most is a good trade union in these times of great social change in the country. Ergo, we need a Secretary who will see that these trade union activities take precedence. The name of the leader of a rival trade union has become almost a conversational cliché in this respect: "We need a Clive Jenkins!" is the cry. We do not, of

course, but do we actually need a doctor? The idea of two wings in the organization—a tough negotiating cadre and a bland advisory medical one, has been canvassed before. Why not have a tough negotiating Secretary anyway?

We must think very hard on this, and we must certainly advertise openly and widely for a new Secretary of the very highest calibre and not allow this to be settled in the corridors of power. The choice will be critical for the survival of the B.M.A. It cannot survive for much longer in its present form.—I am, etc.,

J. P. LEE-POTTER

Department of Pathology,
Poole General Hospital,
Poole, Dorset

Scottish Assembly and the Health Service

SIR,—The Glasgow pathologists' fear (3 May, p. 280) that the 40-hour week will endanger our Scottish traditions of clinical laboratory and academic training is correct. The desire for a return to a salaried service is widespread in Scotland. The Scottish consultants' views on the consultant contract are also quite distinctive.