CORRESPONDENCE

Ulceration of Small Intestine and Slow-release Potassium Tablets S. J. Heffernan, F.R.C.S.I., and J. J. Murphy, F.R.C.S.	746
Computerized Tomography of Brain I. Isherwood, F.R.C.R., and others.	746
Prevention of Overdoses B. M. Barraclough, F.R.A.C.P.	747
Beta-blockers and Fibrinous Peritonitis R. P. H. Thompson, D.M., and B. T. Jackson, F.R.C.S.; I. G. E. Cunningham, F.R.A.C.S.	747
Ocular Reactions to Beta-blockers J. R. W. Lyall, M.R.C.P.	747
Anaphylactoid Skin Reaction Following Intradermal Secretin H. W. Baenkler, M.D., and others.	747
Myocardial Infarction and Metabolism L. H. Opie, M.D.	748
Abortion (Amendment) Bill M. Sim, F.R.C.P.ED.; N. A. Chisholm, L.R.C.P.ED.	748
Psychiatry in Russia Susanne Shafar, M.R.C.PSYCH., and others.	748

Indomethacin, Prostaglandins, and Dysmenorrhoea V. R. Pickles, M.D.; G. G. Hill, M.B.; S. J. Carne, F.R.C.G.P.	749
Hormonal Pregnancy Tests and Congenital Malformations Isabel Gal, M.D.	749
Safer Cigarettes E. R. Trethewie, M.D.	749
G.M.C. and Indian Doctors R. T. Burkitt, F.R.C.S.	750
Immigrant Doctors Farzana Ali.	750
Alternatives to the Fluoridation of Water Winifred M. Sykes; D. Jackson, D.D.S.	750
Inhibition of Ristocetin-induced Platelet Aggregation by Haemaccel Jeanne Stibbe, M.D., and E. P. Kirby.	750
Tests for Lactose Malabsorption in Adults? G. L. Metz, M.B., and others.	751
Humidification of Inspired Air L. B. W. Jongkees, M.D.	751

Arthritis of Hepatitis: the Tourniquet Test R. A. Frayha, M.D., and S. Uthman, M.D.	751
SI Units J. C. B. Fenton, F.R.C.PATH.	751
Little and Big Bellyachers J. P. Childs, F.R.C.S.	752
Management of Depression G. Nicholson, M.B.	752
Screening Methods for Covert Bacteriuria in Schoolgirls Rosalind M. Maskell, B.M.	752
Pay Beds and Professional Freedom R. S. Murley, F.R.C.S.	752
Junior Hospital Staff Contract P. J. Wyld, M.B.; E. B. Nye, M.R.C.O.G.; H. P. Henderson, M.B.	752
Mrs. Castle's Conference Katherine M. Venables, M.B.	753
Economies in the Health Service D. M. J. Burns, F.R.C.S.	753
Practice Premises D. M. Wilks, M.B.	753
A.R.M. Agenda: Item 168 D. Lynch, M.R.C.G.P.	753

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are now being received that the omission of some is inevitable. Letters should be signed personally by all their authors.

Ulceration of Small Intestine and Slow-release Potassium Tablets

SIR,—We noted with interest the recent paper on oesophageal ulceration and stricture associated with Slow-K therapy by Drs. A. D. Howie and R. W. Strachan (26 April, p. 176), and a letter describing a case of mouth ulceration with slow-release potassium by Dr. B. R. McAvoxy (19 October 1974, p. 164). Because of the widely held view that Slow-K does not cause the small intestinal ulceration and stricture associated with other forms of potassium therapy, we should like to report two cases of small bowel ulceration and stricture in patients on Slow-K.

(1) A 73-year-old woman with a history of congestive cardiac failure had been treated for three months with digoxin, Lasix, and Slow-K. She was admitted with small bowel obstruction. At laparotomy an ulcer stricture of the terminal ileum was found and resected. The patient made a satisfactory recovery.

(2) A 65-year-old man with congestive cardiac failure after a myocardial infarct had been treated for four weeks with digoxin, Lasix, and Slow-K tablets. He was admitted with subacute intestinal obstruction, which subsequently became acute and required laparotomy. An ulcer stricture of the terminal ileum was found and resected, and the patient had an uneventful recovery.

The use of slow-release potassium tablets has been urged in the belief that it is the rapid release of potassium over a short segment of intestine, which is the factor precipitating mucosal ulceration and subsequent stricture formation. While this is undoubtedly true it must follow that the local concentration of potassium is the critical factor and that even the slow-release form, when held for any reason at one site, may reach a sufficient concentration to cause ulceration. As this is the third site of mucosal ulceration (mouth, oesophagus, and

now small intestine) to be reported in the *B.M.J.* in recent months it would seem that great care should be exercised in the prescription of slow-release potassium, and that patients on Slow-K should be watched for the symptoms of ulcer stricture.—We are, etc.,

SEAN J. HEFFERNAN

Mater Misericordiae Hospital,
Dublin

JAMES J. MURPHY

Royal City of Dublin Hospital,
Dublin

Computerized Tomography of Brain

SIR,—We note with interest the leading articles (10 May, pp. 295 and 300) "Non-invasive Investigations of the Brain" and "And Now the Whole Body." It has been our experience over the past two years with computerized tomography of the brain (E.M.I. scan) that this revolutionary technique has changed the pattern of radiological investigation and the clinical management of patients with neurological disorders. As a method of visualizing the brain and the pathological processes affecting it it is unparalleled by any other radiological method. Its non-invasive character, outpatient capability, and accurate demonstration of normal and abnormal tissues are indeed major diagnostic advantages.

Conventional neuroradiological procedures are now less often required, with consequent economy of inpatient facilities. Our experience in the department of neuro-radiology at the Manchester Royal Infirmary of the changing patterns of investigation for

space-occupying lesions is summarized in the table contrasting two equivalent time periods in 1972 and 1974. It is noteworthy that the total number of patients attending the department has increased by 50%.

	1974	1972
Computer tomography		
Inpatients	590	—
Outpatients	410	—
Angiography	259	318
C.S.F. pathway investigation	141	207
Isotope brain scan		
Inpatients	87	383
Outpatients	56	184
Totals { Inpatients ..	1077	863
Outpatients ..	466	184

In addition to these major advantages, however, it is important to stress the unique measuring capability of the equipment. Measurements recently carried out in our departments indicate that the method is much more sensitive to changes in tissue which accompany disease states than other conventional radiological techniques. Variations in density of one part in 1000 over a region of about 1 cm³ can be detected. Numerical display of data allows accurate measurement of x-ray absorption coefficients which are related to the mean electron density and atomic number of tissue. Changes of one part in 500 in the atomic number of tissue is just detectable for blocks of tissue 1 cm³. This change corresponds approximately to the difference in atomic number between intracellular and extracellular fluid. These changes are at present just within the limits of detection, though it may well prove possible in the future to detect changes in sodium-potassium flux.

A preliminary study has been undertaken exploiting the precision of the E.M.I. instrument in the quantification of bone mass and bone mineral concentration. Modification of the head device to obtain a cross-section of the wrist has enabled a reproducibility of 0.2% in bone mineral concentration of

cadaver arm to be achieved. Separate measurements of trabecular bone mass, cortical bone mass, and bone mineral concentration have also been achieved in patients. The results of this study have been presented in March 1975 at the International Symposium on Computer Tomography in Bermuda and will be further advanced at the European Society of Radiology in Edinburgh in June. It is important to stress that these results were obtained by modification of the E.M.I. brain scanner where the existence of a surrounding water reference makes accurate and absolute results possible.—We are, etc.,

IAN ISHERWOOD
R. A. RUTHERFORD
B. R. PULLEN

Manchester Royal Infirmary and
University of Manchester

Prevention of Overdoses

SIR,—The inclusion of emetics in hypnotics proposed by Dr. P. M. Vicary (31 May, p. 503) to prevent the fatal overdose has been suggested before. There are barbiturates available, collectively known as Barbemets, which contain a small amount of ipecacuanha. I attempted to assess the relative safety of these preparations by comparing the death rate per million prescriptions of Barbemets with conventional barbiturate preparations. Over the period 1965-70 there were 93m. prescriptions for barbiturates and 12 354 deaths from barbiturate poisoning, a death rate of 133 per million prescriptions. Over the same period 109 000 prescriptions of Barbemets were issued and one death was recorded from a Barbemet preparation, a death rate of nine per million prescriptions, which would suggest that Barbemets are indeed less of a risk, though the fact of only one death being recorded is insufficient for statistical confidence. In the course of the same inquiry nitrazepam was also investigated and gave a result of 11 deaths per million prescriptions, suggesting it to be safer than barbiturates.

However, Dr. Vicary asks how the commonly used hypnotics can be made safer for those who take overdoses, and the simple device of packaging in blisters or putting the active preparation in a mass of inert substance both warrant more official consideration, since the taking of large numbers of tablets is made much more difficult. For example, 50 tablets of nitrazepam 5 mg weigh 27.5 g, four times greater than 50 tablets of Amytal 100 mg, which weigh 7.7 g, a factor contributing to the greater safety of nitrazepam.

A more radical proposal for reducing barbiturate deaths is to stop the prescription of barbiturates by regulation. Even though the prescribing of barbiturates is diminishing as a result of voluntary restraint, deaths from barbiturates are reducing rather less fast.—I am, etc.,

B. M. BARRACLOUGH

M.R.C. Clinical Psychiatry Unit,
Graylingwell Hospital,
Chichester, Sussex

Beta-blockers and Fibrinous Peritonitis

SIR,—We read with interest the description by Mr. J. F. Gurry and others (31 May, p.

498) of a further case of sclerosing peritonitis associated with treatment with practolol. We would urge caution, however, in accepting their suggestion that the more recent use of propranolol may have contributed to the development of the lesion in their patient.

We have had referred to us two patients who developed sclerosing peritonitis, one two months and the other six months (also with pleural effusion) after stopping practolol; neither patient had taken any other beta-blocking drug. One of the patients reported by Brown *et al.*¹ had a similar history and several others have been reported.^{2,4} Since the abdominal effects of practolol can thus be delayed, it is difficult to implicate other beta-blocking drugs given at the same time or at a later date.—We are, etc.,

R. P. H. THOMPSON
B. T. JACKSON

St. Thomas's Hospital,
London S.E.1

- 1 Brown, P., *et al.*, *Lancet*, 1974, 2, 1477.
- 2 Bendtzen, K., and Søborg, M., *Lancet*, 1975, 1, 629.
- 3 Kristensen, K., Kristensen, J. S., and Thorborg, J. V., *Lancet*, 1975, 1, 741.
- 4 Halley, W., and Goodman, J. D. S., *British Medical Journal*, 1975, 2, 337.

** The authors sent a copy of this letter to Mr. Gurry and his colleagues, whose reply is printed below.—Ed., *B.M.J.*

SIR,—I have had the opportunity to read the letter from Dr. Thompson and Mr. Jackson, and in the absence of my two co-authors, who are overseas, I would like to comment on it.

It is most reassuring to learn that cases of sclerosing peritonitis have developed after cessation of practolol therapy without the subsequent use of propranolol. Though there have been previous reports of the delayed onset of peritonitis, the specific point of whether or not propranolol was used in the place of practolol has not been made until now.

In view of this report from Dr. Thompson and Mr. Jackson our cautionary note (31 May, p. 498) is no longer justified, nor is there any evidence to date to imply that propranolol therapy alone may cause this condition.—I am, etc.,

IAN G. CUNNINGHAM

St. George's Hospital,
London S.W.17

Ocular Reactions to Beta-blockers

SIR,—I should like to draw attention to the following case of corneal perforation occurring during treatment with beta-blocking drugs.

A 55-year-old man presented at this hospital in May 1974 complaining of dry eyes. In July 1974 his left cornea perforated. He had suffered from angina for five years and had been started on propranolol in June 1970; this was continued for eight weeks, when practolol 500 mg daily was substituted. He remained on this dose of practolol until April 1973 (32 months). In April 1973 he complained to his general practitioner about dry eyes and oxprenolol 40 mg three times daily was substituted for the practolol. He remained on oxprenolol until July 1974, at which time the cornea perforated.

This case is not easy to assess in terms of causation. I accept that this man was taking practolol for about 2½ years before starting

oxprenolol; nevertheless, he had been off practolol and taking oxprenolol for a full 15 months before the corneal perforation. Clearly oxprenolol is not above suspicion and I feel that this is sufficient reason for the case to be reported.—I am, etc.,

J. R. W. LYALL

Department of Medicine,
St. Thomas's Hospital Medical School,
London S.E.1

Anaphylactoid Skin Reaction after Intradermal Secretin

SIR,—Drs. S. R. Bloom and A. S. Ward (18 January, p. 126) reported on the impairment of secretin release in patients with duodenal ulcer, thus substantiating the original idea of Demling *et al.*,¹ who discussed deficiency of endogenous secretin as a potentially relevant factor in the pathogenesis of that disease. The fact that the secretin deficit in Drs. Bloom and Ward's short-history patients was as great as in their long-history patients may argue in favour of it being a primary defect.

While conducting a trial of exogenous secretin as a possible therapeutic agent in duodenal ulcer we obtained data possibly associated with immunogenic properties of that hormone. About 15 minutes after intradermal injection of 40 IU of synthetic secretin² dissolved in 0.05 ml of 0.9% saline subjects developed a local reaction resembling cutaneous anaphylaxis (urticaria plus surrounding erythema) indistinguishable from that following intradermal administration of 0.05 ml of histamine dihydrochloride (0.1 mg/ml of 0.9% saline). Urticaria and erythema vanished within 60-120 minutes. To study that phenomenon in more detail different groups of subjects who had not previously received secretin were tested in the same way. Eight out of eight duodenal-ulcer patients showed anaphylactoid skin reactions to intradermal secretin, two out of three gastric ulcer patients, four out of six cases of ulcerative colitis or Crohn's disease, and 13 out of 23 subjects suffering from various gastrointestinal diseases other than those listed above (for example, chronic pancreatitis, gastric and colonic polyps, colon cancer). Interestingly enough, six out of seven healthy volunteers did not respond, while one healthy individual developed a questionable reaction. Systemic symptoms were noted in none of the subjects after intradermal secretin injection, nor was sensitization against secretin detectable *in vitro* when passive haemagglutination or blast-cell-transformation techniques were employed.

Viewing the present preliminary data it may be suggested that there is—especially in duodenal-ulcer patients—a factor conditioning anaphylactoid skin reaction due to intradermal secretin. An immunological mechanism based on the presence of secretin antibodies cannot be excluded—though there were positive reactions in subjects who had never before received secretin and in spite of negative *in vitro* tests. On the other hand, non-specific release of histamine or other vasoactive substances by secretin has not been reported so far. Whether or not there are relations between the impaired secretin release in duodenal ulcer patients and a special immunological disposition of those patients against secretin is an un-