Acute gastric dilatation after trauma

Acute gastric dilatation is a condition which, if not diagnosed and therefore not treated, carries a high mortality.¹ While it usually complicates abdominal operations it has been reported after minor operations.² The widespread use of nasogastric suction after abdominal operations has led to a considerable fall in its incidence in recent years.²⁻⁴ Many young surgeons will therefore have no experience of this condition. We report here two unusual cases of acute gastric dilatation, both of which followed trauma to the upper abdomen.

Case 1

A 9-year-old girl was admitted after being in a road traffic accident. On physical examination she was extremely dyspnoeic, there was bruising over the right side of chest and epigastrium, and her abdomen was slightly distended and tender. A chest x-ray picture showed an extensive right-sided haemopneumothorax (see figure), and straight films of the abdomen showed pronounced gastric dilatation with a fluid level. A nasogastric tube was passed and a large amount of gas, together with clear fluid, was aspirated. This resulted in a dramatic improvement in the patient's condition, her abdominal pain was relieved immediately, her respiratory rate fell, and her abdomen became soft and free from tenderness.

The haemopneumothorax was treated by the insertion of a drain through the second intercostal space. During the next 36 hours the patient had recurrent episodes of gastric dilatation as evidenced by a tender, distended, and hyperresonant abdomen. This settled after a further 48 hours' nasogastric suction. Her progress thereafter was uneventful.

Case 2

An 8-year-old boy was admitted after he had been knocked down by a car. On admission he was extremely restless. Abdominal examination showed generalised tenderness and guarding. Plain x-ray examination of the abdomen showed gross gastric dilatation (similar to that shown in the figure). A nasogastric tube was passed, and this relieved his abdominal discomfort immediately. As there was still doubt about the possibility of an intra-abdominal injury, an exploratory laparotomy was carried out. At operation no abnormality was found. In the immediate postoperative period he continued to complain of abdominal discomfort, which was relieved by nasogastric suction. He went on to make an uneventful recovery.

Case 1. Chest x-ray picture showing right-sided haemopneumothorax.

Discussion

In these two cases the clinical features initially suggested the possibility of a major intra-abdominal injury likely to warrant urgent laparotomy. In the first case the extreme dyspnoea and acute distress caused by the gastric dilatation was very impressive. Equally impressive was the immediate relief provided by nasogastric suction, and this despite the concomitant presence of a haemopneumothorax with complete collapse of the right lung.

Although the precise cause of acute gastric dilatation is not known,

certain contributory factors—neurogenic, postanaesthetic, postoperative and nutritional—have been incriminated.²⁻⁵ In blunt abdominal trauma it is possible that a compression of force on the coeliac ganglion might cause reflex vagal inhibition.

The diagnosis should be borne in mind in cases of blunt abdominal injury in which the abdomen is distended and hyper-resonant. Plain x-ray examination of the abdomen confirms the diagnosis. Decompression should be carried out by continuous nasogastric suction for 24-36 hours. If there is no intra-abdominal injury the abdomen immediately becomes soft and the patient's distress is relieved.

We thank Mr W A T Robb for his help in preparing this manuscript.

¹ Starr, K W, *Annals of the Royal College of Surgeons of England*, 1953, **12**, 71.

² *British Medical Journal*, 1970, **4**, 128.

³ Jones, P F, *Emergency Abdominal Surgery*. p 44. Oxford, Blackwell Scientific Publications, 1973.

⁴ Bailey, H, and Love, M, *A Short Practice of Surgery*, 16th edn. London, Lewis, 1975.

⁵ Golby, M, *British Journal of Clinical Practice*, 1966, **20**, 584.

Royal Northern Infirmary and Raigmore Hospital, Inverness

A T KASENALLY, MB, FRCSED, registrar in general surgery

A G FELICE, MB, CHB, registrar in general surgery

J R C LOGIE, MB, FRCS, lecturer in surgery

Guide-wire technique for central vein cannulation

Since Aubaniac's¹ original description of infraclavicular subclavian puncture, several different approaches have evolved and a range of cannulation equipment is commercially available. None of the approaches or equipment are perfect for all conditions. Routine use of a floppy-ended guide wire provides a safe and simple technique for central vein cannulation using any of the commercially available cannula sets. It obviates the need for the cumbersome needle sheath.

Method

The materials used (in a sterile pack) were: gown, gloves, drapes, and swabs; and re-sterilisable guide wire, USCI, C R BARD catalogue number 9225-S, diameter 1 mm, length 70 cm.

The foot of the bed must be raised to engorge the neck veins. Any standard approach²⁻⁴ may be used to puncture the subclavian or jugular vein, with either a 14- or a 16-gauge needle. A small stab wound must be made in the skin first and adequate draping is important. Once the needle is in the vein, the bevel is directed towards the heart. The syringe is removed and venous blood should flow out. The floppy end of the guide wire is advanced down the needle and should meet no resistance. It is inserted about 20 cm and then the needle is removed completely.

A plastic or Teflon cannula is then passed over the guide wire, twisted slightly through the deep cervical fascia and then on into the central vein. The guide wire is then completely removed, and, after a tap has been put on the end, the cannula is sutured to the skin.

The technique may be easily modified for tunnelling of the cannula under the skin and for the insertion of wide-bore Silastic cannulae. The guide wire may also be used to change cannulae at the same site.

Discussion

Guide wires have been used in cardiological and radiological procedures for over 20 years; the original description was attributed to Seldinger.

I have now used the guide-wire technique in over 100 central vein cannulations. This technique has two major advantages. Firstly, it provides greater reliability in placing the cannula in the superior vena cava. This is presumably because the floppy end of the guide wire tends to be carried by the blood stream, whereas the more rigid cannula, on its own, sticks at corners and hence tends to be pushed in the wrong direction. The second advantage is the absence of the needle and sheath and the ease of subsequent nursing care. Ellis and Fielding,⁵ in their review of the commercially available cannulation sets, found that although those with the rigid needle were the easiest to insert, the presence of the needle on the cannula necessitated a protective but cumbersome sheath. This tended to

Teflon cannula being passed over floppy-ended guide wire into left subclavian vein by supraclavicular approach.

produce kinking of the cannula and limited the mobility of the patient's neck and shoulder. The guide-wire technique avoids these problems (see illustration). Subsequent nursing care is greatly simplified and provided the usual aseptic precautions are taken there is no increased risk of infection.

¹ Aubaniac, R, *Presse Médicale*, 1952, **60**, 1459.

² Parsa, M H, *et al*, *Bulletin of the New York Academy of Medicine*, 1972, **48**, 920.

³ Mogil, R A, *et al*, *Archives of Surgery*, 1967, **95**, 320.

⁴ Wilmore, D W, and Dudrick, S J, *Archives of Surgery*, 1969, **98**, 256.

⁵ Ellis, B W, and Fielding, L P, *Journal of the Royal College of Surgeons of Edinburgh*, 1974, **19**, 193.

Western General Hospital, Edinburgh EH4 2XU

JOHN A E GOY, FRACP, MRCP, medical registrar

Raised alpha-fetoprotein levels associated with minor congenital defect

The value of estimating the level of α -fetoprotein (α -FP) in amniotic fluid in diagnosing major defects of the neural tube is well proved.¹ Amniotic fluid levels of α -FP have also been reported to be raised in other serious conditions such as oesophageal atresia and renal tract abnormalities.² A raised α -FP level in association with a minor defect, pilonidal sinus, is reported.

Case report

A 23-year-old woman in her third pregnancy was referred for amniocentesis because her previous child had been born with meningomyelocele and hydrocephalus; he died at the age of 11 weeks. (She had had a spontaneous first-trimester abortion in her first pregnancy.)

A sonar scan confirmed the gestation of 18 weeks with normal outline of fetal head. It also indicated the presence of an ovarian cyst. A laparotomy was performed. The cyst was found to be benign, originating in the left ovary. It was enucleated and the ovary reconstructed. Amniocentesis was performed.

The α -FP level was noted to be 120 mg/l (upper limit of normal 30 mg/l). A maternal blood sample collected before the operation showed her α -FP level to be 40 μ g/l (below the 50th percentile for normal). In view of the high liquor level of α -FP, it was decided to repeat the amniocentesis. The level of α -FP two weeks later was 130 mg/l (upper limit of normal 20 mg/l), and the serum level had risen to 138 μ g/l (95th percentile of normal = 125 μ g/l). In view of these results, the patient was offered a termination of the pregnancy, which she accepted. The pregnancy was terminated by extra-amniotic infusion of a solution of prostaglandin E. The fetus had a postanal dimple, which detailed histological examination showed to be a neuroepithelial lined tract with marked vascularity opening on to the skin (seen in photograph). There were no other abnormalities in the fetus.

Pilonidal sinus of caudal end showing squamous epithelium with very congested vessels in the vicinity. ($\times 112$.)

Comment

All cases of raised α -FP levels reported have been associated with neural tube defects or some other gross abnormality of the fetus. The most likely reason for a false-positive diagnosis is contamination of the specimen with fetal blood. A few red blood cells were observed among the cellular material removed from the first liquor specimen for cytogenetic examination, but none were apparent in the second specimen. No haemoglobin F was detected in the liquor supernatants used for α -FP estimations. The high α -FP levels obtained in the absence of haemoglobin F were confirmed by Dr J H Brock. The other possibility is that a short spell of fetal hypoxia during enucleation of the ovarian cyst may have resulted in production of an excessive amount of α -FP. The amniocentesis performed two weeks later, however, still showed very high levels of α -FP.

Pilonidal sinus is a defect that is easily treated and compatible with a normal life. In retrospect termination of the pregnancy was most unfortunate. This case indicates the need for extreme care in interpreting amniotic α -FP values and the desirability of obtaining confirmatory evidence of a major neural tube defect before proceeding to termination. Additional investigations that may be helpful include the measurement of other markers such as α_2 -macroglobulin³ or procedures such as amniography.⁴

We thank Dr D J H Brock, university department of human genetics, Western General Hospital, Edinburgh, for examining both liquor specimens for α -FP and haemoglobin F, and Dr John McKenzie, reader in developmental biology, University of Aberdeen, for expert advice. Requests for reprints should be sent to Dr V Jandial, Aberdeen Maternity Hospital, Cornhill Road, Aberdeen.

¹ Brock, D J H, and Sutcliffe, R G, *Lancet*, 1972, **2**, 197.

² Seppala, M, *Obstetrics and Gynaecology*, 1973, **42**, 613.

³ Brock, D J H, *Clinical Genetics*, 1975, **8**, 297.

⁴ Connon, A F, *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1974, **8**, 753.

University of Aberdeen, Aberdeen AB9 2ZD

VIJAY JANDIAL, MB, MRCOG, senior registrar, department of obstetrics and gynaecology

HAZEL THOM, BSC, PHD, senior lecturer, department of child health

JANE GIBSON, MB, MRCPATH, lecturer, department of pathology