CORRESPONDENCE

Heat prostration in children with cystic fibrosis A J Williams, MRCP, and others.....297	High-dose corticosteroids in severe acute asthma J B Macdonald, MRCP.....300	Measles encephalitis during immunosuppressive treatment M M Reid, MB, and A W Craft, MRCP.....303
Cardiac muscle relaxation in hypothyroidism J E Sanderson, MRCP.....297	Rubella antibody tests in pregnancy W R G Thomas, FRCPATH.....300	United profession Sir George Godber, FRCP.....303
Accidents on holiday R Mangnall, FRCSGLAS.....298	Nitrazepam and diazepam S L O Jackson, MD.....300	Staffing in the hospital service S S Chatterjee, FRCP.....303
Vitamin A and lung cancer A Sakula, FRCP.....298	Raynaud's phenomenon as side effect of beta-blockers A J Marshall, MRCP, and others.....301	Pay restraint and the consultant W W Wilson, FRCS.....304
Urinary retention in women E Barbara Turner, MB.....298	Alternatives to barbiturate hypnotics D F Scott, MRCPSYCH.....301	Medical manpower and hospital staffing G I B da Costa, FRCSed.....304
Outpatient laparoscopic sterilisation T R M Bristow, FFARCS.....298	Child-resistant containers and child poisoning A W Craft, MRCP, and others.....301	An academic backwater? D J P Gray, FRCGP.....304
Incidence of dermatographism R P Warin, FRCP.....298	Priorities in the NHS M H Mills, MA.....301	Bank holidays and the NHS R H A Campbell, MRCP.....304
Effects of legal termination on subsequent pregnancy G Dixon FRCOG, and J A Richardson, FRCSed.....299	Craniopharyngiomas H B Kennedy, FRCSed, and R J S Smith, FRCSed.....302	Economy in prescribing P M Morris, MB.....304
Compensation for drug side effects J A L Gorrings, FRCPEd.....299	Sexual problems clinic S J G Spencer, FRCPSYCH; J H J Bancroft, FRCPSYCH, and Lesley Coles.....302	Hospital practitioner grade D Mary Ridout, FRCS.....304
Reducing outpatient attendances A P Warin, MRCP.....299	Hazards of modern competitive tennis E M R Frazer, MD.....302	Deputising services J Armstrong, MB.....305
Adverse reactions to practolol Sir Theo Crawford, FRCPATH.....299	Extrinsic allergic alveolitis R Penny, FRACP.....302	Apply to... J H Bulmer, FRCS.....305
Carpal tunnel syndrome and tennis elbow T C Beer, MRCP, and N Memon, MB.....299	Doctors, contraception, and sterilisation A R Hill, FRCSed.....303	Points from Letters Waste in the NHS (C D Korn); High-dose corticosteroids in severe asthma (H G J Herxheimer); System, or 2nd MB? (J George and F P Rugman); Non-accidental poisoning and child abuse (M S Dine); The long and the short of medicine (A A Lewis); Supporting brassiere in mastodynia (J McK Buchanan); Routine preoperative chest radiography (J Saperia); Never put off till tomorrow... (F M Owers).....305
Bromocriptine and breast cancer P A Knowles, MB.....300	Dosage of neomycin sulphate M G Thuse, FRCS, and D P Morgan, BPHARM.....303	
Eggs and hypercholesterolaemia A Elkeles, FFRRCSI.....300		

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters should be signed personally by all their authors.

Heat prostration in children with cystic fibrosis

SIR,—Standard paediatric textbooks note that children with cystic fibrosis are liable to heat prostration. This complication, due to the excessive transcutaneous loss of sodium, has hitherto not been recognised as a real problem in the United Kingdom. We wish to report three cases of hyponatraemic dehydration in children with cystic fibrosis admitted to this hospital in late June and early July this year. The environmental temperature locally has been unusually raised to 10°C above the June-July mean of 1974-5.

Case 1—Male aged 6 years. Presented with a 24-hour history of excessive sweating, anorexia, and weight loss. He had been reluctant to drink. He was severely dehydrated and hypotensive. Plasma urea 10.8 mmol/l (65 mg/100 ml), plasma sodium 129 mmol(mEq)/l, and plasma chloride 82 mmol(mEq)/l. He responded dramatically to intravenous rehydration.

Case 2—Male aged 2½ years. There had been profuse sweating and pyrexia for three days before admission. He had not eaten during this time but had been drinking water well. He was severely dehydrated. Plasma urea was 9.6 mmol/l (58 mg/100 ml) and plasma sodium 124 mmol(mEq)/l. There was a good response to intravenous rehydration.

Case 3—Female aged 2 years. There was a 48-hour history of refusal to drink and increasing drowsiness. She had vomited several times. She was severely dehydrated. Plasma urea 11.2 mmol/l (67 mg/100 ml) and plasma sodium 125 mmol(mEq)/l. In spite of vigorous rehydration she

remained oliguric for approximately 48 hours, but thereafter made a satisfactory recovery.

These three children with cystic fibrosis clearly illustrate the danger of the present unusual climatic conditions. The necessity of salt supplementation in these environmental circumstances should be emphasised to all concerned and, above all, to parents. The

Cardiac muscle relaxation in hypothyroidism

SIR,—Dr J J Manns and his colleagues state confidently in their recent paper (5 June, p 1366) that combined apex cardiography and phonocardiography can reliably measure the isovolumic relaxation time (IRT) of the left ventricle, based on the assumption that the O point coincides with mitral valve opening. For a long time this assumption has been considered to be dubious if not misleading. In 1965 Tavel *et al*¹ found that the O point occurred as early as 0 ms to as late as 52 ms after the left ventricular pressure and left atrial pressure crossover point and concluded that the O point "is not simply an expression of mitral valve opening." More recently, in a combined echocardiographic and apexcardiographic study, Prewitt *et al*² found that mitral valve opening, measured as the time of separation of the two cusps, preceded the O point in all but three of the 57 patients studied

danger of the administration of salt-free glucose solutions also should be borne in mind.

In the past the case has been argued for a safe electrolyte solution that would be commercially available.¹ In the absence of such a readily available form of electrolyte-glucose solution parents will need explicit instructions on how the salt supplement should be administered.

A J WILLIAMS
JOHN MCKIERNAN
FRANK HARRIS

Alder Hey Children's Hospital,
Liverpool

¹ *British Medical Journal*, 1971, 1, 125.

(mean interval 50±28 ms) and that the time of maximum separation of the mitral leaflets also preceded the O point (mean interval 14±26 ms). The O point corresponded more closely to the peak rate of outward wall movement. Dr Manns and his colleagues were, therefore, not measuring an isovolumic relaxation period, although their results do reflect an abnormal relaxation process.

In addition, the effect of ischaemic heart disease cannot be dismissed so lightly. Rubinstein *et al*³ in 1973 found that the IRT measured by combined echocardiography and phonocardiography was significantly prolonged in patients with ischaemic heart disease. It is unclear how much coincidental ischaemic heart disease was contributing to the results found by Dr Manns and his colleagues.

Finally, it is surprising that there was no

significant change in the heart rate after treatment, as bradycardia is a common finding in hypothyroidism. The authors do not show their correction factor or the effect it had on the result.

J E SANDERSON

Hammersmith Hospital,
London W12

¹ Tavel, M E, *et al*, *British Heart Journal*, 1965, 27, 829.

² Prewitt, T, *et al*, *British Heart Journal*, 1975, 37, 1256.

³ Rubinstein, J J, Pohost, G M, and Foster, J R, *Clinical Research*, 1973, 21, 446.

Accidents on holiday

SIR,—Mr M J Merlin's letter (3 July, p 42) highlights a problem which should be avoidable by good communications. It would be ethical to refer back to the general practitioner at home a patient who is treated for an injury on holiday, but prompt action by the GP can be frustrated if an outpatient appointment request disappears into the appointments machinery owing to somebody's failure to appreciate its urgency. It is not unknown for someone to come to a clinic for the first time with a fracture which has become redisplaced in a plaster cast and which has united too far to permit remanipulation.

Hospital operational plans vary, but one common feature is a trauma flow pattern starting at the accident department door. The surest procedure, therefore, is to write a full letter to the senior accident surgeon of the most convenient accident department for the patient, preferably naming the hospital. A copy is posted to the general practitioner at home, who is thus kept informed and can monitor progress. The letter itself is included with the x-rays and given to the patient to be taken in person to the accident department without fail the morning after arrival in his home district.

The only problem I have met from this is the understandable objection of the x-ray department to the loss of its films. The radiologists are naturally concerned at the possibility of missing a diagnosis through inability to report on the films and at the trouble caused if x-rays are unaccounted for when needed again. However, this can be overcome if the x-ray department is informed promptly whenever x-rays are sent out of the hospital.

ROGER MANGNALL

Bury St Edmunds,
Suffolk

Vitamin A and lung cancer

SIR,—In a recent leading article (3 July, p 2) the background to the possible cancer preventive value of vitamin A is discussed. This has consisted mainly of laboratory work both in vitro and in animal experiments. In man there has previously been only one study,¹ of an epidemiological nature, which associated bronchial carcinoma with a low dietary intake of vitamin A.

My colleagues and I have recently carried out a small pilot study² and reported 28 patients suffering from bronchial carcinoma who were found at the time of diagnosis to have significantly lower plasma levels of vitamin A than a control group of 10 healthy subjects and nine patients with non-malignant bronchial disease. Among these 28 bronchial carcinoma cases the plasma vitamin A levels

were significantly lower in those with oat-cell and squamous-cell histology than in those with the undifferentiated large-cell variety. The implications of these findings remain obscure at the present time.

However, in this context it is of interest to recall the findings of Dijkstra,³ who demonstrated a high incidence of bronchial carcinoma in those born in the winter months in Holland. He attempted to explain this by the following hypothesis. The fetus at birth has low levels of vitamin A. In the winter months the level of vitamin A in cow's milk is at its lowest. If the newborn infant is fed on cow's milk its vitamin A level remains low at the critical period when the lung is beginning to function and active bronchial development is proceeding. As a result of this vitamin A deficiency the bronchial mucosa may undergo squamous metaplasia and this later predisposes the individual to bronchial carcinoma when subjected to further bronchial insult (for example, by smoking).

If the bronchial damage by vitamin A deficiency has taken place at such an early stage it is unreasonable to expect success from the US National Cancer Institute's trial of added vitamin A for smoking adults in the hope that it will prevent the development of bronchial carcinoma.

Although successful prophylaxis with vitamin A remains to be proved, there is some evidence to suggest that vitamin A in high doses may have a part to play in the treatment of bronchial carcinoma, especially of squamous-cell type.⁴

ALEX SAKULA

Redhill General Hospital,
Redhill, Surrey

¹ Bjelke, E, *International Journal of Cancer*, 1975, 15, 561.

² Basu, T K, *et al*, *British Journal of Cancer*, 1976, 33, 119.

³ Dijkstra, B K S, *Journal of the National Cancer Institute*, 1963, 31, 511.

⁴ Micksche, M, *et al*, *Österreichische Zeitschrift für Onkologie*, 1974, 3, 70.

Urinary retention in women

SIR,—A cause of urinary retention in women not specifically mentioned in your leading article (26 June, p 1554) is primary genital infection with *Herpesvirus hominis*. This can cause extensive oedema and ulceration of the vulva and introitus. Many patients with herpes have some degree of dysuria, but in particular if periurethral ulceration is present the symptoms are very distressing.

In this clinic two patients with acute retention due to genital herpes have been seen during the past three years. Others, we believe, have avoided this complication with the help of simple advice—for example, to pass urine while standing or sitting in a cool bath. As this infection is now being seen more frequently the incidence of retention of urine in young women is likely to increase.

E BARBARA TURNER

Special Clinic,
Royal Infirmary,
Sheffield

Outpatient laparoscopic sterilisation

SIR,—At a time when the hospital service is forced to wear an ever-tightening financial strait-jacket one can only applaud the spirit and the contents of Mr J H Brash's article (5 June, p 1376). It is to be hoped therefore,

that not too much emphasis will be placed on the dire warnings contained in the letter from Major R H Jago (3 July, p 46).

In contrast to that of Major Jago my own experience does not bear out the view that controlled ventilation with endotracheal intubation is mandatory—or even desirable—in anaesthesia for laparoscopic sterilisation. I would venture to suggest that the anaesthetic technique used for the patients in Mr Brash's series is entirely suitable in the circumstances and, indeed, completely eliminates the possibility of such inconveniences as post-operative tracheitis or pharyngitis, damage to teeth or lips, etc.

But even more contentious is the statement in the last paragraph of Major Jago's letter that "the use of muscle relaxants is contraindicated in day-case anaesthesia." Few if any anaesthetists with experience in this field would concur in that view.

T R M BRISTOW

Royal Berkshire Hospital,
Reading

Incidence of dermatographism

SIR,—I was very interested to read the letter from Dr R F Q Johnson on this subject (19 June, p 1533). It has been difficult to get a simple machine giving different pressures. Various devices have been described in the literature but a useful and inexpensive one was described by Bettley¹ and modified by Kirby *et al*.² This machine can now be ordered from: Hook and Tucker Instruments Ltd Vulcan Way, New Addington, Croydon CRO 9UG. The incidence of dermatographism is discussed by Kirby *et al*² and the whole subject by Warin and Champion.³

There are a number of different types of dermatographism, but there are two common patterns. The first is simple dermatographism, occurring in some 5% of the population with the same incidence throughout all age groups, and this may well be just a physiological variant. The second pattern is symptomatic dermatographism, seen most commonly in young adults and associated with considerable itching; the outlook is good, and many cases settle down after a few months or years. There is an immunological basis for this second pattern and it can be transferred by the IgE immunoglobulin to the skin of a normal subject.

Less common types of dermatographism include those occasionally following an attack of acute urticaria, occurring at the site of urticaria pigmentosa, and where there has been a previous allergic reaction in the skin. There is a form of delayed dermatographism and one where the skin urticates on rubbing rather than scratching, producing a diffuse redness, stippled with tiny weals (cholinergic dermatographism). This last type can sometimes be demonstrated in cholinergic urticaria but may occur on its own and be the cause of a widespread pruritus. The incidence of dermatographism in chronic urticaria is probably no greater than in a series of control patients.

R P WARIN

Bristol

¹ Bettley, F R, *Journal of Investigative Dermatology*, 1962, 39, 1.

² Kirby, J D, *et al*, *British Journal of Dermatology*, 1971, 85, 331.

³ Warin, R P, and Champion, R H, *Urticaria*. London, Saunders, 1975.