The increased number of over-75s is the main problem facing the Health Service. The only resource that will be available will be flexibility. The twin problems of policies to help elderly people with mental illness and to define the role of old people's homes must be tackled. The family practitioners, community nurses, volunteers, and neighbours of elderly people with mental illness can only do so much. A realistic residential policy must be worked out and the keystone of this must be the old people's homes

J A MUIR GRAY
J S RODGERS

Oxfordshire Area Health Authority (Teaching),
Oxford

¹ Population Projections No 4 1973-2013, prepared by the Government Actuary. London, HMSO, 1974.

² Department of Health and Social Security, *Health and Personal Social Services for England 1975*. London, HMSO, 1976.

³ Department of Health and Social Security, *Community Hospitals: Their Role and Development in the National Health Service*. London, DHSS, 1974.

⁴ Wallingford Community Hospital Research Project. Oxford, Oxford Regional Hospital Board, 1973.

What do community physicians do?

SIR,—Dr A Roberts (17 July, p 178) was quite right in questioning whether I was not comparing the former medical officer of health (MOH) with the present community physician. My contention is that the problems are still there to be dealt with.

The MOH in the past was the only professional to oversee the Victorian environment and the crude medicosocial problems of the day. As these were tackled and were held at bay a number of technicians had to be trained and later took on almost independent professional responsibilities regarding the environment, water, sewage, housing, cleansing, and the like. The same occurred in the personal services of domiciliary midwifery, health visiting, and mental health. The one essential in the provision of these municipal community services was the legal requirement to make them work, a mandatory situation which does not apply to clinical medicine, where, if everyone will speak truthfully, most of my colleagues unquestioningly and automatically apply our Hippocratic tradition, but some do not.

It is this unique overall responsibility of the MOH to get things done in spite of everything which has disappeared. There is nothing in its place to cope immediately and decisively with emergencies or to provide that background of comprehensive professional knowledge which is essential to medicosocial planning. This is why community physicians as successors of MOHs feel that their hands are tied behind their backs so that they are merely the spectators of situations which need their involvement. We have been translated into a defective, hospital-orientated, lay administrative system in which each section works like battery hens laying eggs at their own rate without regard to others, without any central communication or collation of information, without any feedback from other sections, and without spirit. The present system has no central registry or mechanism for the interpretation or co-ordination of the snowstorm of circulars from the DHSS (Elephant droppings) which confine our every action.

Imagine the apoplexy of a surgeon if the rest of the team did their own thing in their own time or said they just didn't believe in

surgery anyway. This is the present position for community physicians, but we are still supposed to sort out anything and everything as well as being involved throughout the planning process.

W S PARKER

East Sussex Area Health Authority,
Brighton Health District,
Brighton, Sussex

Modified operation for ingrowing toenails

SIR,—Ingrowing toenails are one of the commonest conditions seen in surgical clinics. Operations, which tend to be left to junior staff, are frequently complicated by infection and later by nail regrowth due to an incompletely excised nail bed. We have, however, had very good results since using the following operation. It is usually performed under local anaesthesia, with a tourniquet.

An elliptical segment, including the edge of the nail and the inflamed nail fold, is excised (see figure). The incisions are cut down to the bone so that a V-shaped segment of inflamed tissue is removed. This ensures that all the nail bed and all infected granulation tissue are completely removed. The wound is then cleaned with an antiseptic solution.

No sutures are used to close the defect; instead, a length of Steristrip $\frac{1}{2}$ in (13 mm) is wrapped firmly around the toe. This produces very good approximation of the skin edges without leaving any open space beneath. A firm bandage is applied before the tourniquet is removed. This dressing is left undisturbed for eight days if possible.

This operation has been performed on numerous occasions and has been characterised by being surprisingly pain free and having a low incidence of infection.

B V PALMER
D LANG STEVENSON

Whipps Cross Hospital,
London E11

Rebound hypertension after acute methyldopa withdrawal

SIR,—We were interested in the report by Drs A C Burden and C T P Alexander (1 May, p 1056) of rebound hypertension after acute methyldopa withdrawal. Recently we observed a similar occurrence.

A 77-year-old hypertensive woman taking methyldopa 250 mg twice daily and cyclopentiazide was admitted with acute gouty arthropathy. Cyclopentiazide was discontinued without alteration in blood pressure control at 130/80 mm Hg, and allopurinol treatment was begun. After one week the blood urea remained elevated at 12.6 mmol/l (76 mg/100 ml) and it was considered that her blood pressure control might be excessive. Methyldopa was therefore withdrawn. Within a day her blood pressure had begun to rise, and 48 hours later, when it was 240/160 mm Hg, diazoxide 300 mg was given intravenously with immediate good effect. Satisfactory blood pressure control was later again obtained with methyldopa 250 mg twice daily.

In view of the fact that this patient's hypertension was easily controlled, both before and after cessation of therapy, by a small dose of methyldopa we consider the acute hypertensive crisis to have been a demonstration of rebound precipitated by acute methyldopa withdrawal. This would support Drs Burden and Alexander's suggestion that rebound hypertension occurs with hypertensive drugs other than clonidine.

J N SCOTT
D G McDEVITT

Belfast City Hospital and
Department of Therapeutics
and Pharmacology,
Queen's University,
Belfast

Short-term recovery of mental efficiency after anaesthesia

SIR,—The paper by Mr J E P Simpson and others (26 June, p 1560) is a beautiful piece of scientific research and writing, but the results are meaningless.

For day-case surgery the aim is to produce adequate intraoperative anaesthesia and post-operative recovery of consciousness which is as rapid and complete as possible. Premedication is part of the total anaesthetic management of a patient, and though the anaesthetic techniques compared by Mr Simpson and his colleagues are standard, premedication for day-case surgery is normally restricted to nitrazepam or some quiet reassurance. (If early recovery of consciousness is vital anaesthesia may be induced with propanidid rather than thiopentone.) Patients given a premedication of papaveretum 20 mg and hyoscine 0.4 mg would probably require 7-9 hours to return to 60% of mental efficiency without any general anaesthetic.

I hope Mr Simpson and his colleagues will repeat their excellent study with alternative and lighter premedication.

E LL LLOYD

Department of Anaesthetics,
Royal Infirmary,
Edinburgh

Propranolol in hypertension

SIR,—I believe that the article by Dr D B Galloway and others (17 July, p 140) might be seriously misleading in that they have not given sufficient emphasis to the fact that their study deals only with patients who are mildly hypertensive and who are treated with propranolol alone. The so-called "law of the initial value"^{1,2} suggests that the benefit of drugs may be proportionate to the severity of the condition treated, and my own experience

very strongly supports the view that in more severe degrees of hypertension the benefits of propranolol are dose-related up to amounts of at least 1 g daily, as well as exerting a markedly beneficial effect in some circumstances as soon as 30 minutes after ingestion.

As a hypertensive myself I have found it expedient if not imperative to limit the rate of intake of propranolol to 80 mg in any period of 40 minutes at least, and during a period of some weeks last year, after unwisely reducing the total daily dose to 400 mg, it became essential, particularly during the evenings, actually to consume the drug at an average rate of 2 mg/min. Whatever the long-term benefits of propranolol, I was left in no doubt that, given a certain level of active drugs already available to the body, definite effects occur within 30 minutes in slowing the pulse and reducing arrhythmia. Equally, it is quite clear that the necessary rate of intake varies with the time of day. The efficacy of relying on subjective sensations of need has been validated by objective observations of the consultant physician at regular review in the outpatient department.

As a psychiatrist I am inevitably reminded of the different clinical benefits of phenothiazine drugs in the short term and in the long term, but I hope that psychiatrists are better acquainted with the desirable upper limits of dosage in their own field than the highly respected physicians of Aberdeen appear to be. If read carefully the article from Aberdeen should not mislead, it is true, but I think one would have to be pedantic—or experienced—not to run the risk of being so misled.

J COLLINSON

Broadgate Hospital,
Beverley,
N Humberside

¹ Wilder, J, *Journal of Nervous and Mental Disease*, 1957, 125, 73.

² Bierman, E L, *Lancet*, 1976, 1, 1076.

Abortion and maternal deaths

SIR,—Your leading article (10 July, p 70) reminds us of the mortality of therapeutic abortion but does not mention the morbidity that more commonly occurs. The sequelae include pelvic infection resulting in sub-fertility and tubal pregnancy, now known to cause 9% of maternal deaths. Cervical laceration may be followed by middle-trimester abortion, premature labour, and an increased perinatal mortality.¹ Perforation of the fundus, unrecognised or undocumented, may cause rupture of the uterus in subsequent labour. Failure to give immunoglobulin to the rhesus-negative patient can be catastrophic. Therapeutic middle-trimester abortion of the young primigravida on purely social grounds should perhaps now be condemned.

Pregnancy in young girls is not especially hazardous provided they receive antenatal care. In Wolverhampton in the past five years 56 primigravidae were delivered while still under 16 years of age. Anaemia (haemoglobin less than 10.5 g/dl) occurred in 33% but was treated effectively since postnatal anaemia occurred in only 12.5%. Toxaemia was not unduly common (10.7%) although there was one eclamptic. The induction rate, 17%, was half that of the unit as a whole. A quarter of the patients had epidural analgesia and 23.2% were delivered by forceps. There were no caesarean sections. Unfortunately there were three stillbirths, but there were no neonatal

deaths. Undoubtedly the two 13-year-olds had complicated pregnancies. One had a difficult forceps delivery, while the other, who had no antenatal care, sustained abruption of the placenta and stillbirth. Whether the pregnancy is terminated or allowed to continue, it is surely vital to ensure that these young girls use effective contraception, because of those who become pregnant again, over 40% will do so within 12 months.¹

ALAN M SMITH

New Cross Hospital,
Wolverhampton

¹ Richardson, J A, and Dixon, G, *British Medical Journal*, 1976, 1, 1303.

Out-of-hours calls in general practice

SIR,—We were interested to read the account by Dr M G F Crowe and his colleagues (26 June, p 1582) of the way in which they provide out-of-hours care in their practice. They suggest that small partnerships which cover their own out-of-hours calls provide a more traditional, personal service to patients and that one of the results of this relationship is that a sizeable proportion of requests for help can be handled over the telephone, avoiding the need to visit the patient. They contrast this position with that of a deputising service, quoting from our study of the Sheffield deputising service, in 1970¹ the fact that 97% of calls resulted in a visit from a deputy.

We believe that this is not a comparison between like situations and that it gives a misleading impression. The Leicestershire doctors averaged 302 out-of-hours calls per year compared with 106 per year per subscriber received by the Sheffield service. Although the average list size of the Leicestershire practice doctors was approximately 25% greater than that of Sheffield doctors and their out-of-hours period was one hour longer on weekdays than that for which the Sheffield deputising service operated, these factors can account for only a small part of the three-fold difference in the number of calls. A more important factor is likely to be that the Sheffield subscribers passed only a proportion of their out-of-hours calls to the deputising service, dealing with the remainder either by giving advice over the telephone or by a personal visit to the patient. Although we do not know how often this occurred, it was clearly common practice for Sheffield subscribers to screen their calls since 47% of all those received by the deputising service in 1970 came via the general practitioner. In other words, the high proportion visited by the deputising service arose in part because the practitioners had already given advice to a number of patients and had made referral to the service unnecessary.

Dr Crowe and his colleagues also criticise our estimate that the deputising service handled between 36% and 69% of the night calls for its subscribers, suggesting, on the basis of the night-call rates in their practice, that the proportion should be much higher, over 90%. Here again it is very doubtful whether the circumstances of a semirural group practice are sufficiently similar to the pooled experience of 100 urban subscribing practices to justify such conclusions.

The problem remains that we do not know quantitatively how primary medical care is being dispensed out of hours in large areas of this country, nor do we know much about the effectiveness of the various methods or

their acceptability to the public. We recognise the limitations of assessing the size of the deputising service contribution by means of the indirect methods which we, and now Dr Crowe and his colleagues, have used. We envisage that a true picture can be obtained only through a study of a situation in which various methods of operating in the same place at the same time, a situation common to the majority of our large towns.

B T WILLIAMS
R A DIXON
J KNOWELDEN

Department of Community Medicine,
University of Sheffield Medical School,
Sheffield

¹ Williams, B T, Dixon, R A, and Knowelden, J, *British Medical Journal*, 1973, 1, 593.

Management of eclampsia

SIR,—In your leading article on this subject (19 June, p 1485) you refer to hypovolaemia in these patients and the danger of using diuretics. Although you mention death from intracerebral haemorrhage, no comment is made on the morbidity and mortality from renal failure.

Kennedy *et al*¹ included 20 women with pre-eclampsia or eclampsia in a large series of medical, surgical, and obstetric patients admitted to their unit with acute renal failure. Seven of the 20 died and those who survived had a higher incidence of residual renal impairment than surviving patients from the other groups.

Thomson *et al*² described the typical reversible renal lesions in pre-eclampsia of endothelial and mesangial cell proliferation and fibrinoid deposition. Such changes are aggravated, with decrease in glomerular function, when hypovolaemia reduces renal perfusion. It was suggested that prompt correction of hypovolaemia lessens the risk of irreversible damage. Bonnar³ reported that a similar problem in patients with abruptio placentae can be avoided by using central venous pressure (CVP) monitoring to maintain adequate circulating blood volumes.

Recently we have inserted CVP lines via the antecubital fossa in six patients with severe pre-eclampsia. This was performed in the first two cases when reduction in the urinary output was noted and was used in the subsequent four cases to monitor fluid replacement both before and after delivery. Venous pressure below -5 cm H₂O, measured from the angle of Louis, was noted in five of these patients. Normovolaemic states were restored with infusions of albumin and isotonic saline. Only then was frusemide or mannitol used to maintain urinary output. No deterioration in renal function occurred.

We believe that to avoid further renal damage CVP monitoring is an important aid in achieving a normovolaemic state in the management of severe pre-eclampsia or eclampsia.

ALLAN B MACLEAN
JOHN R DOIG
D R AICKIN

University Department of
Obstetrics and Gynaecology,
Christchurch Women's Hospital,
Christchurch, New Zealand

¹ Kennedy, A C, *et al*, *Quarterly Journal of Medicine*, 1973, 42, 73.

² Thomson, D, *et al*, *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1972, 79, 311.

³ Bonnar, J, *Clinics in Haematology*, 1973, 2, 213.