count, the clinical picture and the markedly increased liver enzyme values were misleading and the protein-bound iodine result was thought to be erroneous. Thus the patient was on parenteral nutrition for almost two weeks before a firm diagnosis was made. The treatment resulted in dramatic improvement within a few days.

MATTI SAARNI
ELLI KOIVUNEN

Hatanpää Hospital,
Tampere, Finland

SIR,—The paper by Dr F D Rosenthal and others (24 July, p 209) prompts us to record a recent patient with thyrotoxicosis who, while not vomiting, was decidedly liverish.

A 63-year-old farmer presented with three months' history of loss of weight (25 kg), shivering, sweating, and high backache. Mild scleral icterus and a pulse rate of 90/min were present together with a palpable liver edge and a palpable spleen. Initially the patient was investigated for neoplasm with a negative result. A serum thyroxine concentration of 412 mmol/l (32 mg/100 ml) reverted to within normal limits (46–148 mmol/l (5–11.5 mg/100 ml)) after treatment with radioactive iodine, as did abnormal liver function tests, the serum bilirubin level falling from 44.5 μ mol/l (2.6 mg/100 ml) to 13.7 μ mol/l (0.8 mg/100 ml), serum albumin rising from 28 to 37.5 g/l and total protein from 55 to 70 g/l, and the Thrombotest result rising from 45% to 100%.

Dr Rosenthal and his colleagues mention "slightly raised" liver enzyme levels in four cases out of seven and in their first case the serum bilirubin level "was raised" at 25 μ mol/l (1.5 mg/100 ml). It is worth remarking that our patient never suffered from anorexia of any kind during the course of his illness and all abnormalities disappeared after treatment with ¹³¹I. A subsequent liver scan was normal.

Liver abnormalities have, of course, been described in thyrotoxicosis but rarely of sufficient note to orientate the diagnostician clinically towards a gastrointestinal investigation.

THOMAS KIERNAN
M McELLIGOTT

Portiuncula Hospital,
Ballinasloe,
Co Galway

Management of diabetic pregnancy

SIR,—Your leading article (31 July, p 267) on the management of diabetic pregnancy is rather out of date. Firstly, you observe that the perinatal mortality is now "close to 10%", when in fact it is now much less than this. At this hospital it has decreased from 9% of 176 deliveries in the five years 1966–70¹ to 4% of 144 deliveries in the period 1971–4,² which is maintained at 3.5% when 1975 is included (total of 173 deliveries). This improvement is at least partially due to close collaboration between physician, obstetrician, and paediatrician, which is crucial in the management of diabetic pregnancy in order to achieve the careful antenatal and intrapartum monitoring which is needed.³ Furthermore, quoting a publication now nine years old, you observe that respiratory distress is the most common cause of death, whereas it is now exceptionally rare and we have had no such deaths for several years.² You make no mention of the increasing fetal loss from congenital malformations, which are now the

commonest single cause of perinatal death.^{2,4} Finally, you say that the insulin requirement falls "soon after delivery." In fact the fall is immediate and there is a great danger of very severe hypoglycaemia if the insulin dose, which may have increased two- or three-fold during pregnancy, is not reduced to pre-pregnancy levels on the day of delivery.

D A PYKE
J M BRUDENELL
P J WATKINS
H R GAMSU
N L ESSEX

Kings College Hospital,
London SE5

¹ Essex, N, *et al*, *British Medical Journal*, 1973, 4, 89.
² Essex, N, *British Journal of Hospital Medicine*, 1976, 15, 333.

³ Brudenell, M, in *Carbohydrate Metabolism in Pregnancy and the Newborn*, H W Sutherland and J M Stowers. Edinburgh, Churchill Livingstone, 1975.

⁴ Soler, N G, *et al*, *Quarterly Journal of Medicine*, 1976, 45, 303.

* * * The recent figures for perinatal mortality obtained at King's College Hospital are unusually good, but the figure of "nearly or slightly below 10%" may be regarded as more typical in the country generally. The leading article did not say that hyaline membrane disease or "respiratory distress" is the commonest cause of death but rather that it has been found to be the commonest cause of death of babies born prematurely to diabetic mothers, the "has" being used because a relatively old source of information was quoted. The decreased incidence of this cause of death is doubtless due to several factors, of which a significant one is the study of amniotic fluid mentioned in the previous sentence. Considerations of space prevented direct reference to the importance of congenital malformations, but this is one of the main factors which make it, as we said, "increasingly difficult to reduce the perinatal mortality in diabetics any further." The final point, whether the insulin requirement falls "soon after delivery" or "immediately," is hard to judge as the mother usually has an intravenous drip of glucose running at this time, but certainly the insulin dose should be decreased from the time of delivery.—Ed, *BMJ*.

Urinary retention in women

SIR,—I am glad that the letters from Mr S L R Stanton and Mr T Moore were published together (14 August, p 420). I see about one woman each year who has "female prostatic obstruction" exactly as described by Mr Moore; as he says, they respond well to the removal of the obstruction by endoscopic resection. So I am surprised that Mr Stanton found no such case in his series of 27 women with retention or a slow stream.

But why does Mr Stanton consider pressure-flow studies with sphincter electromyography so important? It is safe to assume that his 27 patients were thought to have "acute retention," "chronic retention," or "a noticeably slow stream" as the result of a good history and a simple examination. To determine a drug-induced aetiology requires only careful inquiry and the trial substitution of the drug thought responsible. Similarly, a neurological aetiology is surely determined by the history and the relevant clinical examination. Maybe there is now a urodynamic finding that is actually diagnostic of hysteria, but I doubt it.

I have seen two women with retention which had been elsewhere considered to be due to hysteria. One, with repeated painful episodes of acute retention, had a cyst of the urethra and trigone; the other, with chronic retention, had a "female prostate"—both responded well to endoscopic resection.

In 1976, when impecunious area health authorities like mine cannot provide a consultant with a desk, let alone an office, it behoves the teaching hospitals to consider carefully, when advocating such very expensive gadgetry, whether equally good results could not be obtained with a pin and a bit of cotton-wool.

ROGER HOLE

North Ormesby Hospital,
Middlesbrough,
Cleveland

SIR,—While I am in agreement with Mr S L R Stanton (14 August, p 420) that hysteria may be an important aetiological factor in many cases of acute retention of urine in women, I must disagree that chronic retention "may be considered to be the result of a single or repeated episode of acute urinary retention."

Once obvious obstructive urological (for example, stone, tumour), neurological, and gynaecological lesions and a history of drug ingestion have been excluded we believe that chronic urinary retention is not a single entity but consists of at least two different aetiological groups. The first is characterised by a residual urine volume of less than 500 ml and displays the self-perpetuating spiral of chronic inflammatory changes at the bladder neck,^{1,2} residual urine, and infection; the second is characterised by the "modesty type" of bladder with a residual urine volume in the region of 1 l or more. While a hysterical personality trait may be found in some of the women in this second group there is, I would propose, no evidence that the first group may progress to the second and no evidence that, except perhaps in an immediate postoperative period, episodes of acute retention of urine in women may pass unnoticed and progress to chronic retention.

GERALD J JARVIS

St Mary's Hospital,
Leeds

¹ Fox, M, Jarvis, G J, and Henry, L, *British Journal of Urology*, 1976, 47, 797.

² Evans, A T, *Journal of Urology*, 1971, 105, 245.

Abortion and maternal deaths

SIR,—Dr Ann Cartwright in her letter (24 July, p 232) is quite right. There exists much evidence of undue delay in reaching decisions to abort. Certainly, there is enough to justify reforming action now rather than at some unspecified time in the future.

For example, a recent survey¹ in the Cardiff area shows that a significant proportion of family practitioners consider the service at the main hospital (University Hospital of Wales) to be unsympathetic or unsatisfactory or inadequate in substantial part because of delays by consultants in making appointments. Incidentally, the survey shows that about half the women in Cardiff seeking help with termination have to be referred outside the area because of prevailing attitudes at the University Hospital. Only 10 of the 197 family practitioners in the South Glamorgan Area (15% of respondents) were prepared to

report that the service (help with terminations) at UHW was sympathetic or satisfactory or adequate.

Another survey² shows that at the University Hospital of Wales only one in five patients referred to the professorial department of gynaecology for consideration of termination was seen within one week of referral; 30% had not been seen within two weeks; 57% of vaginal terminations were performed at or beyond the twelfth week. "Early interview and assessment would enable most terminations to be performed not later than the twelfth week of pregnancy."

The reforming action needed now is simple: a DHSS circular to area health authorities requiring them to make appropriate agency arrangements with one or more of the established non-profit-making voluntary agencies. Family practitioners would then be able to refer locally and the conscientious objections of NHS consultants and others need then not be challenged.

CASPAR BROOK

David Owen Centre for Population Growth Studies, Cardiff

¹ Family Planning Association, *Where family practitioners in South Glamorgan refer Requests for Help with Termination*, April May 1976 (6 Windsor Place, Cardiff).

² Chalmers, I, and Anderson, A, *Lancet*, 1972, 1, 1324.

Meditation or methyldopa?

SIR,—Your leading article (12 June, p 1421) discusses sympathetic nervous activity in hypertension, suggests that mild cases may benefit from transcendental meditation (TM) or yoga, and points out that the scope for such treatments is undefined. These are but fragments of two major problems facing us today. The first is the extent to which hypertension and its related cardiovascular disorders are the outcome of a morbid relationship between the individual and his environment—that is, of a relationship that creates excessively severe and prolonged arousal leading to fatigue and exhaustion.^{1,2} The second is the extent to which relief from these disorders should be obtained by modifying the morbid relationships, as opposed to "treating" the raised blood pressure pharmacologically as if it was a "disease" in its own right.

Of 67 patients admitted for routine investigation and treatment of hypertension (range 160/100-230/130 mm Hg) 33 (50%) became normotensive when the arousal and exhaustion were treated. In 26 (39%) the blood pressure fell to normal levels on some occasions but failed to stay there consistently. In one patient admission to hospital was not well tolerated and his blood pressure rose. Only in seven (9%) did the blood pressure fail to respond to the treatment of the arousal. Of these seven, two had renal failure, one had renal carcinoma, and four had to be regarded as fixed-essential hypertension. Of the 33 patients who recovered normal blood pressure one-third had arrived at the point of exhaustion and ill health² because their behaviour was maladaptive, and two-thirds had been brought down by circumstances causing extreme frustration and anger or despair and hopelessness.³⁻⁸

The hospital methods for reducing arousal included the sleep regimen as employed in myocardial infarction,⁹ lighter levels of sedation, a warm and caring atmosphere, and the creation of a support group for the period

of the patient's need. An attempt was made to teach every patient to recognise the level of arousal and fatigue that made his blood pressure high and to be responsible for his own care. In the longer term the patients were taught to use diazepam as a defence against morbid arousal and encouraged to learn how to reduce arousal at will, a method advocated by Mackenzie.¹⁰ Some were taught to relax and others adopted TM or yoga. All were taught to regard sleep deprivation as dangerous.

Economically we believe that a great saving could be made if the initial routine investigation became the general practitioner's observation of the blood pressure response to the reduction of arousal and the removal of exhaustion at home. It is reasonable to agree with Pickering's¹¹ contention that established hypertension can be the outcome of long periods of transiently raised blood pressure and that the transient rises have much to do with the patient's "behaviour of mind" or arousal. A greater medical awareness of the heightened sympathoadrenal activity of life on fatigue and exhaustion, a wider recognition of maladaptive behaviour, and a much closer inquiry into the patient's attitudes and circumstances might lead to "essential" hypertension becoming inessential.

P G F NIXON
DAVID H DIGHTON

Charing Cross Hospital, London W6

¹ Nixon, P G F, in press.

² Kagan, A R, and Levi, L, *Social Science and Medicine*, 1974, 8, 225.

³ Lorenz, K, *On Aggression*. London, Methuen, 1970.

⁴ Kennedy, A, *Lancet*, 1957, 2, 261.

⁵ Chambers, U N, and Reiser, M F, *Psychosomatic Medicine*, 1953, 15, 38.

⁶ Birley, J L T, *Proceedings of the Royal Society of Medicine*, 1973, 66, 96.

⁷ *British Medical Journal*, 1971, 2, 215.

⁸ *British Medical Journal*, 1975, 1, 458.

⁹ Nixon, P G F, *Practitioner*, 1973, 211, 5.

¹⁰ Mackenzie, L, *Diseases of the Heart and Aorta*. London, Oxford Medical Publications, 1908.

¹¹ Pickering, G, *Lancet*, 1976, 2, 1403.

Consent and intrauterine contraception

SIR,—In the medicolegal columns (31 July, p 310) you quote the advice of the Family Planning Association on the obtaining of husband's consent for the insertion of an intrauterine contraceptive device. In my opinion the advice is unhelpful and confusing if not actually contradictory. It seems that the alternative to the husband's agreement is for the doctor to be satisfied that the insertion is "reasonably necessary for the proper treatment of the wife." This being so, unless the financial inducement is greater than one had imagined, what is the need to weigh up the wife's assurance about her husband's agreement? Surely there is no other reason for inserting such a device than that it is "reasonably necessary."

In what way does the keeping of routine notes exonerate the doctor from obtaining the husband's consent, if indeed consent is necessary? Is not the purpose of the notes to record when the wife withholds permission to communicate with her husband, or that the husband withholds his consent? Does such a record really upset the husband's legal rights? This surely is the crux of the matter. Has the husband the right to give his wife a child regardless of her own wishes? The insertion of an intrauterine device at best, or worst, depending upon one's viewpoint, does not permanently deprive him of this right, if

indeed he has it. If he has such a right he has the normal processes of the law to recover it if he considers he has been improperly deprived of it. However, the doctor who deprived him of his right albeit temporarily may need protection.

It is not sufficient for the FPA to issue advice or make recommendations as to practice in its clinics. The doctors practising in such clinics need a clear statement of the law and if the law is ill defined or unsatisfactory have the right to expect that our legislators will amend it to bring it into line with other recent social legislation affecting the status of women. Meanwhile, if a man has the legal right to have children by his wife and to decide when to have them, regardless of her wishes, it is dangerous for a doctor to interfere with that right. Now that most FPA clinics have been or are being transferred to area health authorities the DHSS surely has an obligation to clarify the legal position and if need be to initiate action to change it.

In the meantime many doctors working in family planning clinics are not seeking the agreement of the husband or the wife's assurance of it. They believe such agreement to be unnecessary and derogatory to the status of women. Neither for that matter do they seek the permission of the wife before supplying her husband with condoms. The BMA would be doing these doctors a service if it sought to clarify this matter now rather than leave it to the courts to decide.

NORMAN TALBOT

Margaret Pyke Centre, London W1

Uraemic bullae

SIR,—During the summer of 1975 two middle-aged male patients with chronic renal failure who had not yet started haemodialysis treatment developed bullae on the dorsum of the hands, particularly on the thumbs, and on other exposed areas. One patient had obstructive uropathy and the other had chronic glomerulonephritis. Neither of them had any past history of skin trouble or porphyria. The first patient, aged 47, was taking Slow-K sodium bicarbonate, vitamins B and C, aluminium hydroxide, and large doses of frusemide. The second patient, aged 54, was taking vitamin D, aluminium hydroxide, iron, folic acid, and frusemide 2 g daily. Their blood urea levels were 51.5 mmol/l (310 mg/100 ml) and 30.3 mmol/l (182 mg/100 ml) and plasma creatinine levels 1.9 mmol/l (21 mg/100 ml) and 1.24 mmol/l (14 mg/100 ml), respectively. Similar lesions have been described by Gilchrist *et al*¹ and Korting² in patients on haemodialysis; they postulated that the lesions in their patients were a complication of dialysis. Recently Professor A C Kennedy and Dr A Lyell (19 June, p 1509) have described remarkably similar lesions in patients on high doses of frusemide. All but one of their patients had severely reduced renal function.

The clinical similarity of the lesions in these four groups of patients, all of whom were uraemic, is striking. The aetiology is now known. Frusemide could be responsible in our patients, as in the Glasgow patients. However, many patients undergoing conservative treatment for chronic renal failure take this drug. In view of the occurrence of similar lesions in patients on haemodialysis who were not taking frusemide we postulate that these