Clinics in General Practice

A case of asthma

K A O MOURIN, J COLLINS

British Medical Journal, 1976, 2, 1003-1004

The trainee's problem

A 13-year-old boy was brought by his mother to the surgery recently with generalised rhonchi over both lung fields, both inspiratory and expiratory, and most severe in his left lower lobe. He was given ampicillin and the infection gradually settled.

This attack had followed the pattern usual for him—namely, an increase in frequency and severity of his nocturnal cough, the onset of an audible wheeze, and production of small amounts of purulent sputum. Between episodes he is normally free of wheeze, although nocturnal cough is present more nights than it is absent.

Since the family moved to this area four years ago the boy has been a frequent attender at the surgery, usually for respiratory tract infections but twice with vague and undiagnosed abdominal pain and once after an episode of "fainting and collapse" at school. The frequency of his respiratory infections, however, is not diminishing; these occur mainly during the winter and average four or five episodes annually, although already this year he has had more attacks than at any time, and this in turn is affecting his schooling.

A brief account of the family history may be interesting. This boy is the second child in a family of four. He has an older brother aged 14, a younger brother aged 11, and a sister aged 5. His father, aged 37, is presently suffering from sciatica and often cannot work. His mother, aged 33, suffers from attacks of urethritis and has cervical spondylosis. There is a history of allergy on the maternal side, with the boy's maternal uncle and grandfather both affected by asthma. The three boys all have eczema and suffer from hay fever and various allergies (these include allergy to plants and animals but not to food or drugs), although this is most severe in the patient's case. His elder brother has been seen on numerous occasions both by ourselves and the hospital because of frequency of micturition and loin pain, which after many negative urine cultures and radiological and bladder studies has now settled with no specific treatment.

His mother is naturally worried about the frequency of the boy's chest infections, the family history of asthma, and time off school. The principal problem is whether a course of disodium cromoglycate (Intal) or bronchodilators, or both, is now indicated. Should the patient be referred to the asthma clinic for lung function studies and skin testing to establish a baseline measurement before treatment?

General practitioner's comments

The problem here concerns the whole family, not just the boy. Looking at the problem in clinical, psychological, and social terms helps to identify those areas where expert help and advice are needed as well as those perhaps best dealt with away from hospital.

Clinically the boy presents with classic atopic disorders and, although all too little is known about the cause and therefore prevention of these, effective empirical therapeutic measures are available. Children may wheeze because of allergy, infection, exercise, cold air, and other lung irritants, and even spontaneously. I should want first to know if his nocturnal symptoms indicate a postnasal drip from a chronically infected sinus, bronchiectasis, open windows, or an allergic response to household pets, dust, or mites.

The psychological and social issues raised by this story require careful thought, as these may be crucial in deciding whether referral is necessary or inadvisable at this stage in the illness. The boy's history of vague abdominal pains is common in children under stress. His elder brother had long-investigated urinary symptoms, father has recurrent "sciatica," mother has "urethritis" and "cervical spondylitis"—all rather tenuous labels. The family pattern of disease is repeating itself. What are the time relationships of the parents' symptoms and the boy's? Are mother's periods related to his bellyache or wheeze?

The general practitioner needs to look inside the home himself. Cigarette smoke, dogs, cats, budgerigars, dusty bedrooms, broken windows, shared rooms—all tell a story. Is the house downwind from the local tip or smoking factory chimney? Where does the father work, doing what, and how often? Where did he come from four years ago, and why? Are there social or environmental factors that can be changed?

What are the family's expectations about illness? Are they "under-the-hospital" people whose past experience has led them to expect hospital investigation and management? These patients' disorders are often so ill-defined, inexplicable, intractable, and unresponsive to conventional treatment that it is not surprising that in sheer frustration, or in fear of missing something extraordinary but identifiable by expert investigation, they are referred to hospital clinics. Intensive investigations so often fail to identify a genuine cause or allay symptoms but merely perpetuate their sick role and reinforce their expectations.

My advice with this problem, which is perhaps a more subtle one than it initially seems, is again divided into a physical part and a psychosocial part.

The asthma clinic can help the doctor to assess the underlying condition by chest and perhaps sinus x-ray examination and lung function studies. Skin testing or other tests for allergies may help, though often they add little to a good history. The doctor should, however, be clear in his own mind—and state explicitly in his letter—just why he is referring this particular patient, what aspects of the problem he needs help with, and whether he wishes to continue the general management of the patient himself.

Dereham, Norfolk

K A O MOURIN, MB, MRCP, general practitioner

St Bartholomew's Hospital Medical College, London EC1

J COLLINS, MD, MRCP, senior lecturer in medicine (now consultant physician, Brompton Hospital, London)

On the other hand, I should appreciate more information about the social and psychological background, about each family member's ability to cope with information about himself, his "illness," and the effect of this on the rest of the family. Close co-operation with health visitor and social worker would help.

If the doctor has a capacity for and interest in counselling, supportive care, and rehabilitation he could probably manage this problem on his own. Patient education should aim towards positive actions that the boy, his mother, and the whole family can take in preventing and coping with his episodes of chest troubles. Influenza prophylaxis (unless contraindicated by egg or feather sensitivity) and the development of positive attitudes towards health and the family's ability to cope with disturbances in physical and psychosocial wellbeing should all be planned, probably in consultation with the health visitor.

I personally would be prepared to try treatment on my own for, say, six months with, firstly, a bronchodilator and, secondly, if necessary, disodium cromoglycate. If either clinical control is not achieved or patient satisfaction apparent I would certainly think seriously of referral for specialist opinion, advice, and support.

Consultant's comments

Much of the detail of this boy's problem is clear from the history, and there can be little doubt that we are dealing with asthma. Nocturnal cough is often a prominent symptom of this condition in both children and adults. The frequent calls for treatment indicate that the time is ripe for a collaborative approach by the general practitioner and a hospital physician. Decisions about treatment should be based on a review of his disability and the possible part taken by constitutional and environmental factors. In this way the relative merits of different treatments are more likely to become apparent.

At the referral visit to the hospital clinic further information sought should include more-detailed accounts of the allergies to animals and plants; any temporal association, both immediate and delayed, between exposure to these things and the development of wheezing or coughing should be established. An exercise history would be useful—Do school activities and play-ground games lead to symptoms? Similarly, details of the family's housing and of the boy's bedroom will help in planning his treatment.

Clinical examination should pay particular attention to physical development and the presence of chest deformity and growth retardation, in addition to eliciting signs of airways obstruction. The investigations should include assessment of blood and sputum eosinophil levels, x-ray film of the chest, and skin-prick tests for common allergens with the addition of any others that seem important from the history. The lung function tests should be comprehensive to confirm the presence of reversible airflow obstruction, disturbance of other aspects of pulmonary function, and to assess the baseline on which to judge any effects produced by treatment. Lung size should be assessed by measurement of total lung capacity and its subdivisions. These measurements will reflect lung growth and may also be affected by airways obstruction. Measurements of expiratory airflow should include forced vital capacity (FVC), forced expired volume in one second (FEV₁), the ratio FEV₁:FVC, and peak expiratory flow rate. Although not mandatory, it would also be useful to assess respiratory gas exchange by measurements of carbon monoxide transfer. At this visit a simple exercise test would be useful, for it may precipitate airflow obstruction even if this is not present at the preliminary testing. Subsequent exercise tests would also help in showing whether

disodium cromoglycate effectively prevents exercise-induced airflow obstruction.

The aims of treatment should be to render the boy wholly free of symptoms as much of the time as possible and to treat exacerbations promptly. Stress should be laid on the many increases in therapeutic choices for asthma that have recently become available and an attitude of cautious optimism should be encouraged. The best way to assess response to treatment is with diary records of symptoms of wheeze and cough, use of bronchodilator aerosols, and at least twice-daily measurements of the peak expiratory flow rate. The boy and his parents should be given suitable instruction and provided with a peak-flow meter or gauge for temporary use at home. The social and financial circumstances of the family are not clear from the details given, but in most cases it should be possible to effect simple measures for environmental control aimed at reducing exposure to household dust and other potential allergens. Striking improvement may sometimes follow simplifying bedroom furniture, removing carpets, regular vacuum cleaning of floors and bedding, and substituting synthetic materials for feathers and wool.

On general principles it is usual to begin treatment with bronchodilators, often using a combination of oral and inhaled preparations. In this boy's case, however, his symptoms are perennial, and on the evidence of his cough he probably has some degree of airways obstruction most days. For this reason the suggestion of the general practitioner is sound and starting with a combination of disodium cromoglycate and a bronchodilator aerosol is likely to give more satisfactory control of his asthma. He should be taught the use of a Spinhaler and a pressurised aerosol. Treatment should begin with disodium cromoglycate, one capsule four times daily, preceded by one puff of salbutamol, terbutaline, or rimiterol aerosol. He should also be encouraged to use the Spinhaler 30-45 minutes before exercise or games if exercise tests have shown that this protects him from exercise-induced airflow obstruction. Initially, assessments should be repeated at one to two week intervals, and he should be taught to regard any need for additional doses of bronchodilator as indicating poor control of symptoms. At this stage assessment may be shared by the general practitioner and hospital physician.

Within four to six weeks it should be apparent whether this regimen is working and if it is it can be gradually tapered to the minimum doses required for good control. If control is not satisfactory, some additional advantage may be achieved by adding an oral bronchodilator of the theophylline group, which have a different and possibly synergistic action from that of the B sympathomimetic group already mentioned. Further poor response should lead to a change to a trial of one of the steroid aerosols in conventional doses, substituting this for the disodium cromoglycate. With this sequential approach most patients with asthma resembling that shown by this boy may be expected to come under control. Subsequently, the responsibility for follow-up is probably best left to the general practitioner, although ready access for further combined management should be maintained.

Postscript to the problem

The boy was seen at the asthma clinic and, after lung function studies, was started on disodium cromoglycate (Intal) four times a day, with the recommendation that a bronchodilator be added if necessary. Since then he has been well. His mother, however, has now brought the elder boy to the surgery because he has a wheeze too and she wonders if . . .