preceded the development of the classical clinical features of multiple sclerosis by almost four years.

Case report

A 36-year-old man began to have paroxysmal attacks of dysarthria, ataxia, and weakness of his legs in February 1972. Each attack would last up to 10 seconds and he would experience at least a dozen such episodes each day. They would occur more frequently after a hot bath or any sudden exertion. In between his attacks he was free of symptoms and signs but during them he had bilateral horizontal nystagmus, dysarthria, and ataxia of his legs. In view of his unusual symptoms he was investigated in detail. Among the normal results of tests were cerebrospinal fluid, air encephalography, and four-vessel arteriography.

Treatment with carbamazepine resulted in the complete cessation of his attacks. He was followed up at monthly intervals and eventually, in January 1976, developed numbness of both lower limbs and ataxia of gait. Examination then disclosed bilateral horizontal nystagmus, cerebellar and pyramidal signs in all four limbs, with extensor plantar responses and absent abdominal reflexes. Examination of his cerebrospinal fluid then showed a protein of 0.45 g/l of which 23% was IgG. The Lange colloidal gold curve was 1232100000.

Comment

The diagnosis of multiple sclerosis, although presumed, could be established only when this patient developed the classical features of this condition four years after his presentation with a paroxysmal brain stem disturbance. The dramatic response of these paroxysmal symptoms to carbamazepine has been noted by others.^{3,4}

¹ Miley, C E, and Forster, M F, *Neurology*, 1974, **24**, 458.

² Andermann, F, *et al*, *Neurology*, 1959, **9**, 211.

³ Espir, M L E, Watkins, S M, and Smith, H V, *Journal of Neurology, Neurosurgery, and Psychiatry*, 1966, **29**, 323.

⁴ Harrison, M, and McGill, J I, *Journal of Neurology, Neurosurgery, and Psychiatry*, 1969, **32**, 230.

Department of Neurology, Manchester Royal Infirmary, Manchester M13 9WL

W H PERKS, MB, MRCP, registrar in neurology

R G LASCELLES, MD, FRCP, consultant neurologist

Unusual cerebral inclusions in Reye's syndrome

We wish to report on unusual inclusion bodies, revealed by electron microscopy, in the brain of a patient with Reye's syndrome.¹

Case report

A 13-year-old boy was admitted with a two-day history of confusion and drowsiness after an influenza-like illness lasting three weeks. He was unresponsive to painful stimuli and had irregular respiration with periods of apnoea. The arms were areflexic, but reflexes in the legs were brisk with extensor plantar responses. The serum contained persistently raised levels of aspartate aminotransferase (67-121 IU/l) and bilirubin (37-61 µmol/l). The serum calcium concentration fell from normal to 1.4 mmol/l causing tetany. The cerebrospinal fluid was normal. Serial EEGs showed high amplitude repetitive complexes at 2 Hz initially but became isoelectric later. Carotid angiography showed no intracranial blood flow. Viral studies for herpes simplex, measles, and influenza A and B were negative. Treatment was supportive but assisted ventilation was finally discontinued and the patient died on the fourth day after admission.

At necropsy the liver was found to be enlarged to 2610 g. The brain weighed 1655 g and showed signs of severe oedema: the gyri were flattened, the sulci obliterated, and the cerebellar tonsils herniated. Midline structures (septum pellucidum and subependymal areas of the third ventricle), the thalamus, and the lentiform nuclei showed liquefactive necrosis—a feature of the respirator brain death syndrome.²

Light microscopy showed abundant microvesicular fatty change affecting all liver cells, but no inflammation or necrosis. Similar changes were also present in other parenchymatous organs, notably in the renal tubular epithelium, pancreas, and myocardium. The brain showed non-specific cellular changes, which had resulted from oedema and anoxia. Thus the criteria of Reye's syndrome—namely, encephalopathy and microvesicular fatty change of viscera—were fulfilled.

Electron-microscope examination of the cerebral cortex revealed a spectrum of extensive neuronal damage ranging from mild degeneration to necrosis. The neuronal mitochondria were badly affected: their matrices were swollen and their cristae disrupted. Myelin sheaths had lost their structural integrity: there were widened, irregular spaces between individual lamellae. Astrocytic cytoplasm appeared to be swollen.

Discussion

All these changes, described already,³ may be due to hypoxia. They also may be artefactual, resulting from delayed fixation. It was, however, interesting to find large, complex inclusion bodies in some neurones and vascular pericytes. These inclusions were irregular and occupied most of the cytoplasm in many cases. They consisted of lipid droplet-like areas and masses of electron-dense granules. In the neurones they resembled an agglomerate of lipofuscin granules, while in the pericytes they were larger and more complex in structure (see figure). The pericytic inclusions were similar to residual bodies found

A large inclusion in the cytoplasm of a vascular pericyte. L=lumen ($\times 7465$).

in the hepatocytes of patients with Reye's syndrome.⁴ Osmiophilic dense bodies, different from those described here, were found in pericytes and were thought to reflect a mild disturbance of lipid metabolism.⁵ Thus probably these inclusions in pericytes, together with the lipofuscin-like bodies in neurones, are of greater importance in the diagnosis of the cerebral lesions of Reye's syndrome than the other changes, all of which can be produced by anoxia.

¹ Reye, R D K, Morgan, G, and Baral, J, *Lancet*, 1963, **2**, 749.

² Towbin, A, *Human Pathology*, 1973, **4**, 583.

³ Partin, J C, *et al*, *Journal of Neuropathology and Experimental Neurology*, 1975, **34**, 425.

⁴ Svoboda, D J, and Reddy, J K, *Laboratory Investigation*, 1975, **32**, 571.

⁵ Chang, L W, *et al*, *Archives of Pathology*, 1973, **96**, 127.

Bland-Sutton Institute of Pathology, and Department of Neurological Studies, The Middlesex Hospital Medical School, London W1P 8AA

P L LANTOS, MD, MRCPATH, senior lecturer and consultant neuropathologist

P P ANTHONY, MB, MRCPATH, senior lecturer and consultant histopathologist

C R A CLARKE, MB, MRCP, registrar in neurology