were performed on spontaneously breathing patients via the nasal route under local anaesthesia. The indication was usually for retained sputum with lobar collapse. Bronchoscopy improved the air entry in all cases, quite dramatically so in some. One such case is illustrated in fig. 3.

This 17-year-old youth sustained contusion of his right lung and a pneumothorax due to a motor cycle accident. The pneumothorax was successfully treated by an intercostal drain, but over the next three days, despite physiotherapy, first his left lower lobe and then his left upper lobe collapsed (fig 3a). This collapse was resistant to hourly physiotherapy. Fibreoptic bronchoscopy showed some increased secretions in the right tree and complete occlusion of the left main bronchus by blood clot and mucus. This was aspirated and was seen to extend to the segmental orifices of both lobes. Eventually all visible blood was aspirated. The left chest movement and air entry were

dramatically improved, as was the chest x-ray picture (fig 3b).

There have been no complications from the procedure and it is well tolerated, even by very sick patients. Therapeutic bronchoscopy with a flexible fibroscope is therefore a valuable procedure in selected cases and should be available in intensive therapy units.

References

- ¹ Lindholm, C E, *Critical Care Medicine*, 1974, **2**, 250.
- ² O'Donnell, J E, *Anaesthesia and Intensive Care*, 1975, **3**, 139.
- ³ Sackner, M, *American Review of Respiratory Disease*, 1975, **111**, 62.
- ⁴ Lett, Z, and Ong, G B, *Anaesthesia*, 1974, **29**, 623.
- ⁵ Lewis, R B, *Anaesthesia*, 1974, **29**, 625.
- ⁶ Albertini, R E, et al, *Journal of the American Medical Association*, 1974, **230**, 1666.

Bone and Joint Diseases

Hip fractures up to date

P A RING

British Medical Journal, 1976, **2**, 1429-1431

Fractures of the femoral neck are the commonest serious leg injuries seen in the elderly. They have a high associated morbidity and mortality, not because of the injury but mainly because of the physical and mental consequences in the elderly of staying in hospital with a period of relative immobility. Both morbidity and mortality can be diminished by early and energetic treatment, and most of these patients may, with a combination of good orthopaedic and geriatric care, return to their previous environment with little or no additional disability.

Although a history of a fall in these patients is common, some will admit that the leg just gave way, and the injury is clearly associated with a senile osteoporosis. Thus many of these fractures are caused by disease.

Diagnosis

The history of a fall, or the story that the leg suddenly gave way, is a characteristic one, and the diagnosis in most elderly patients should not be in doubt. The leg lies in an externally rotated position and appears shortened, and any attempt at movement is painful. In some patients, however, the fracture may be impacted, and indeed walking is sometimes possible although uncomfortable after such an injury. The risk of neglect in these patients is obvious; such a fracture, which is a relatively favourable one, may disengage and become displaced.

The immediate management is the provision of suitable analgesics. Splinting the limb by tying it to the opposite member is a useful way of diminishing the discomfort during a journey to hospital.

RADIOGRAPHS

Good anteroposterior and lateral films are necessary not only to confirm the diagnosis but also to establish the level of the lesion since this influences the orthopaedic management to a considerable extent. Three main groups of fracture can be identified.

Subcapital

The fracture line is essentially at the junction of head and neck. It may lie in a relatively horizontal plane or may incline more to the vertical; the latter position is more unfavourable since the fracture is inherently unstable. Subcapital fractures may be classified again in more than one group.

Impacted—There is no disengagement at the fracture site; indeed, the outer side of the fracture is impacted into the femoral neck giving an essentially stable position. These are the fractures on which walking is often possible after the injury has occurred.

Undisplaced—These are very similar to the impacted fractures. There is no appreciable displacement on the femoral head relative to the neck, but both parts of the femur are disengaged at the fracture line.

Displaced—Here the amount of displacement may vary widely from a simple varus angulation of the neck to a complete separation of neck and shaft.

Basal neck fractures

Basal neck fractures occur at the junction of the neck and the trochanteric region; separation of the fragments may vary from slight to a very major angulation.

Trochanteric

The fracture line passes through the trochanteric region separating the lesser trochanter that lies as a free fragment and occasionally the greater trochanter as well. Displacement may again vary in severity

from a fracture that is visible only as a hairline crack to one with major displacement of the femoral shaft on the femoral neck and head.

The importance of the distinction in the radiographic appearances lies in the method of surgical treatment and does not influence the general management.

Treatment

Because the risks of prolonged immobilisation in the elderly are well known treatment is surgical save under the most exceptional conditions. Even in a patient who is normally confined to bed or a wheelchair it may be preferable to fix the fracture securely in order to relieve pain, make nursing easier and therefore more effective, and lessen what would otherwise be an additional disability. There is no place for long-term traction nor plaster immobilisation in the hope of securing bony union, but it may be in the best interests of the demented or the paralysed patient not to treat the fracture at all, particularly if it seems to be a source of relatively little discomfort.

Surgery must be carried out at the earliest possible moment which for most patients will be within a few hours of the accident. Although on general medical assessment some of these patients appear poor operative risks, in few is the risk diminished by a day or two spent on general medical care, and it is usually better to treat a fracture as an emergency and then deal with the medical problems that many of these elderly patients pose.

Operative principles—The blood supply of the femoral neck is derived from a vascular anastomosis that passes from the trochanteric region up the femoral neck close to the bone and into the femoral head. Thus fractures at the base of the neck and above carry with them the risk of disturbing the blood supply of the bone above, and such a vascular disturbance may in time produce ischaemic necrosis and may in part be responsible for non-union at the site of the fracture. The nearer to the head of the femur the fracture lies the greater this risk, and for this reason removal of the femoral head and part of the neck and replacement by a prosthesis may in many patients be superior to the results of internal fixation. The balance of the argument is often a fine one, and there is no doubt that a soundly united femoral neck fracture in a good position with a viable head carries with it a much greater chance of a permanent long-term solution to the femoral neck fracture than removal of the implant and a simple prosthetic replacement of the femoral head, a procedure that not only carries with it a higher mortality but may over the years be associated with acetabular wear.

It is generally accepted that the impacted subcapital fracture is best treated by simple internal fixation, and may occasionally be managed conservatively throughout. It is also generally accepted that the widely displaced subcapital fracture in which contact between the femoral neck and the head has been lost carries with it such a high risk of avascular necrosis that primary prosthetic replacement is the procedure of choice. Between these two extremes, however, there may in some patients be a place for internal fixation and in others a place for prosthetic replacement. To some extent the age of the patient may influence such a decision. It may be better to strive for union in the relatively young patient, even at the risk of a period of protected weightbearing, while in the elderly a rapid solution by prosthetic replacement may often be preferable.

Internal fixation—To achieve satisfactory internal fixation the fracture must first of all be reduced, a manoeuvre that must be gently and simply executed to avoid the risk of further vascular damage. Simple traction, gentle abduction, and gentle internal rotation should bring the fragments back to the anatomical position. If this cannot be achieved internal fixation is certain to fail. The traditional method of fixation is the Smith-Petersen trifin nail that ideally passes through the lateral femoral cortex, rests on the calcar, and penetrates the centre of the femoral head, a position of three-point fixation. Various other methods have been produced to improve the fixation of the two fragments—

multiple pins, Garden's lag screws, Smyth's triangle with two lag screws, and various systems of sliding nail. The more complex the system the more difficult it is to obtain a good technical result, and it seems unlikely, despite the statistical evidence, that more complex methods of fixation carry with them an appreciable advantage over the simpler and time-honoured Smith-Petersen procedure. Indeed, in many patients the fate of the femoral fracture is sealed at the time of the accident and perhaps at the time of the operative manipulation.

If the vascular supply is good and the fracture is easily reduced then the conditions for internal fixation to succeed are favourable, and the method of fixation may well be immaterial provided it is well applied. No method of internal fixation, however secure, can overcome the problem of the avascular femoral head.

Prosthetic replacement—Removal of the femoral head and the insertion of a prosthesis overcomes the problems of non-union and avascular necrosis. This can be done either with the Moore's prosthesis or by using a Thompson prosthesis inserted into the femoral shaft with the aid of bone cement. Undoubtedly both these procedures carry a higher morbidity and mortality than those of simple internal fixation, and during the postoperative period there are risks of the prosthesis dislocating and also of infection. Dislocation at best means a period of traction in recumbency. Infection is always a disaster and usually necessitates removal of the implant. In practice both these complications commonly end fatally, clearly both to a large extent avoidable by good surgical technique.

Basal neck fractures—Fractures of the base of the femoral neck again carry with them the risk of delayed union and a risk, although a smaller one, of avascular necrosis of the femoral neck and head. Nevertheless, this risk, in distinction to the high fracture, is not sufficiently great to justify the use of prosthetic replacement even when displacement is wide, but fixation with a simple trifin nail is inadequate, and this must be supplemented, as in the trochanteric fracture, by a side plate.

Trochanteric fractures—Secure internal fixation of these fractures can be achieved by the insertion of a trifin nail with a side plate, the latter screwed to the femoral shaft. There are advantages in a one-piece appliance but a two-part appliance serves equally well. Clearly, whatever device is used its strength must be sufficient to permit immediate weightbearing, and unless the fracture is grossly comminuted and the fixation poor, this is usually possible. More recently Enders has described a method of internal fixation using pins up the femoral shaft that avoids the need of a wide open exposure and thereby diminishes the risk of postoperative infection. This method has not yet gained wide acceptance.

Postoperative management

The object of internal fixation is to allow early mobilisation, and two or three days after the operation these patients can stand and walk bearing full weight upon the injured limb. The problems thereafter are twofold, firstly, the avoidance of complications immediately associated with the injury and operation, and, secondly, the general medical care and rehabilitation.

The commonest postoperative complication of a hip fracture is pulmonary embolism, and Seivitt some years ago advocated prophylactic anticoagulants to overcome this particular difficulty. Their use has not gained wide acceptance, partly because they are not without risk, and partly because over the past few years other methods of avoiding both venous thrombosis and pulmonary embolus have gained ground. The routine risk of low-dose heparin and the use of dextran for two or three days postoperatively both have their advocates, but for this particular group of patients in the early postoperative period an elastic supporting stocking to encourage venous return, early mobilisation, and keeping the legs horizontal when sitting are probably as important as any of the medical measures of prophylaxis.

Many of these patients present general medical problems often

unrecognised until the fracture occurs, and in addition many present social problems that are brought to light only on their admission for hospital care. It is in these cases that the close association between the orthopaedic, geriatric, and social services departments of a hospital are of the greatest importance. All these patients need a full medical and social assessment at a relatively early stage, and many need a planned period of rehabilitation not only for the hip fracture but also for general medical care before they are fit to return home. An active and energetic geriatric rehabilitation unit is essential if these patients are to return to full activity, and with such a service most in fact will do so. In the postoperative period the orthopaedic aspects of this type of injury play only a small part.

Prevention

Clearly in many patients a hip fracture is in fact a pathological fracture through osteoporotic bone, and its prevention, if this

were possible, would be a major improvement in the care of the elderly. A few patients with hip fractures will in fact be found to be suffering from osteomalacia and a correction of this may prevent a similar injury on the other side at a later date. There is some evidence that giving oestrogens to postmenopausal women reduces the risks of senile osteoporosis and might therefore be expected to diminish the incidence of hip fractures, but in general it seems unlikely that in an aging community much can be done to diminish the incidence.

Over 30 000 people each year in England and Wales, almost all elderly, and most of them women, are admitted to hospital after a fracture of the hip. Some will die as a result of the injury or the treatment, or more commonly from intercurrent disease. Some will inevitably end their days in a long-stay bed or in sheltered accommodation, but with good surgical treatment and a high standard of geriatric and nursing care most can return to their place in the community with no additional disability caused by the injury.

Aspects of Student Health

Psychological problems of students

C J LUCAS

British Medical Journal, 1976, 2, 1431-1433

A consideration of the psychological problems of students must take place against a complex and changing background. Rapid social and technological changes are taking place in society. Values are changing, or under challenge; uncertainty pervades increasing areas of life. Whether changes are seen as adverse or beneficial, they are likely to contribute to a climate of conflict and stress that individuals will react to in their own personal ways.

Within higher education there have been major and continuing changes. The 1960s saw rapidly rising student numbers and the development of new universities. Within institutions knowledge has exploded, courses have diversified, and options have increased. More has been questioned than ever before. Ironically, economic recession has brought a reduction of subsequent vocational opportunities. How such complex factors may influence late adolescent development and identity formation must largely be a subject for speculation, but they are likely to have contributed to the various modes of unrest and disturbance that have been enacted in universities in recent years, and which find their counterpart in the consulting room.

Those working with students in clinical services have themselves, over the years, experienced change and confusion, as conceptual models of "health" and "illness," "normality" and "disturbance" have lost the illusory clarity of earlier years. The development of new therapeutic approaches has added to the range of possible provision for such problems, but theoretical

confusion and difficulties in evaluation have contributed to uncertainty.

Developments this century

To understand the present position it will be useful to recapitulate briefly some of the important developments of the past few decades. A more fully documented account of some of the earlier aspects is provided by Mair.¹

The Boer War led to a recognition of a deplorably low standard of physical health among recruits. The concern evoked eventually resulted, after various commissions and inquiries, in the establishment of the school health service. A subsequent royal commission in 1902 commented on the lack in universities of facilities comparable with those in schools. The emphasis at this time was exclusively on detecting physical illness and promoting health by physical recreation. So far as the universities were concerned, all other student problems were regarded as educational or spiritual and the concern of tutors and religious advisers. For many years, therefore, medical activities were confined to detecting physical illness (particularly tuberculosis) and to promoting health by physical exercise, to which end many universities appointed directors of physical education. Thinking continued predominantly on these lines in the years between the two world wars.

The Goodenough report (1944)² and reports from the Social and Preventive Medicine Committee of the Royal College of Physicians (1943 and 1946)³ advocated the further development of this preventive approach, with its implicit notion that physical health was something separate from intellectual and personal maturation. Thus by the mid-1940s the structure envisaged was that services should be attached to departments of preventive and social medicine. It was thought that NHS facilities would