mutation rate in man was between 15 and 100 rads, so the risk of genetic damage is likely to be insignificant. Were a woman of reproductive age to be treated, I would recommend application of the "10-day rule" as in diagnostic radiography to lower the risk of irradiation of a young embryo.

Court Brown and Doll⁴ described an increased incidence of leukaemia after irradiation for ankylosing spondylitis. This was, however, rare, with only 60 definite cases among 14 554 treated. Radiotherapy for ankylosing spondylitis usually entails treatment of much haematogenous bone marrow in the spine and pelvis, unlike the areas treated in this series in which leukaemia is therefore probably not a serious risk. Steiner⁵ reported 12 cases of bone sarcoma developing after radiotherapy and reviewed the published cases. The given doses of radiation were much higher than that used in this series, and the overall risk appears to be very small.

I thank Professor H L F Currey for helpful advice, and Miss S Shah for preparing the typescript.

¹ Currey, H L F, in *An Introduction to Clinical Rheumatology*, ed M Mason, 2nd edn, p 244. London, Pitman Medical, 1975.
² Dewing, S B, in *Radiotherapy of Benign Disease*, p 55. Springfield, C C Thomas, 1965.
³ Alexander, P, in *Atomic Radiation and Life*, p 180. Harmondsworth, Penguin, 1965.
⁴ Court Brown, W M, and Doll, R, *British Medical Journal*, 1965, **2**, 1327.
⁵ Steiner, G C, *Cancer*, 1964, **18**, 603.

(Accepted 15 March 1978)

London Hospital, London E1 1BB
B S MANTELL, MRCP, FRCP, consultant in radiotherapy and oncology

Is penicillamine therapy in rheumatoid arthritis influenced by previous treatment with gold?

The clinical response and side effects of both gold and penicillamine are similar.¹ Gold binds to connective tissues and may remain there long after treatment has stopped.² Penicillamine can release gold from the tissues and has been used to treat gold toxicity.³ Some of the side effects attributed to penicillamine in patients who have received gold may in fact be due to the gold itself. We have investigated whether previous gold treatment influenced the clinical response and incidence of side effects in our patients with rheumatoid arthritis (RA) who were being treated with penicillamine.

Patients, methods, and results

One hundred and fourteen patients with RA had been treated with penicillamine; 75 of these had received gold, which was stopped either because of side effects or a poor therapeutic response. In most cases penicillamine had been started within one year of stopping the gold. During treatment with penicillamine a total of 68 side effects had occurred in 50 patients. Marrow depression had occurred on nine occasions and 19 patients developed rashes. Ten rashes had occurred during the first three months of treatment and the others after the ninth month. Improvement was judged to have occurred in 56 patients.

The influence of previous gold treatment on clinical response and side effects is shown in the table. Although the incidence of all side effects was the same in both groups, marrow depression and rashes were more common in the gold-treated patients. Two patients with marrow depression had had

Influence of gold on penicillamine treatment in 114 patients with rheumatoid arthritis

	Previous gold (% of 75 patients)	No gold (% of 39 patients)
Marrow depression ..	9	5
Rash ..	20	10
All complications ..	45	41
Improvement ..	52	54

neutropenia when on gold, and four had had a rash. Seven patients with a rash thought to be due to penicillamine had also developed a rash with gold. One had developed alopecia, one mouth ulcers, and one neutropenia. Eight of the ten patients with an early rash due to penicillamine had been on gold treatment.

Comment

Although our observations suggest that there is no difference between the clinical response or the overall incidence of side effects with penicillamine whether patients had previously been treated with gold or not, the most important fact is that rashes and marrow depression were twice as common in the gold-treated group. Of the 15 gold-treated patients who developed a rash with penicillamine, seven had had a rash while on gold. A multicentre trial group⁴ reported that treatment with gold resulted in no appreciable difference in the prevalence of side effects in RA patients on penicillamine, though noting that there were more rashes in the gold-treated group. Tsang *et al*⁵ found that 14 of 17 patients with advanced RA who had not responded to gold improved with penicillamine. They found no correlation between the toxic manifestations of the two drugs, although two patients had mouth ulcers, one proteinuria, and one a rash with both penicillamine and gold.

Our results show that rashes, and possibly marrow depression, may be more common if patients have received gold. Theoretically, also, some of the early rashes noted with penicillamine might in fact have been caused by gold being released from the tissues by penicillamine. It would be sensible to take extra care in the supervision of patients starting on penicillamine if they have had gold treatment.

¹ Huskisson, E C, *et al*, *Annals of the Rheumatic Diseases*, 1974, **33**, 532.
² Block, W D, *et al*, *Journal of Pharmacology and Experimental Therapeutics*, 1941, **73**, 200.
³ Davis, C M, *American Journal of Medicine*, 1969, **46**, 472.
⁴ Multicentre Trial Group, *Postgraduate Medical Journal*, 1974, **50**, suppl No 2, p 77.
⁵ Tsang, I K, *et al*, *Arthritis and Rheumatism*, 1977, **20**, 666.

(Accepted 15 March 1978)

St Stephen's Hospital, London SW10
M WEBLEY, MB, MRCP, senior registrar (present address: Westminster Hospital, London SW1)
E N COOMES, MD, FRCP, consultant physician

Synovial cyst causing an inguinal mass

Diagnosis of a lump in the groin may be difficult and the possibility of malignancy occurring often leads to biopsy. We present two cases in which synovial cysts caused such diagnostic difficulty.

Case 1

A 69-year-old woman had had intermittent pain in the right groin for two years. She had suffered from severe rheumatoid disease for 13 years and her medication had included prednisone. During admission for elective knee surgery, examination of the right groin showed a firm fixed swelling, 3 x 3 cm, above the inguinal ligament and lateral to the femoral artery. It was not tender or reducible. The results of full blood count and biochemical tests were unhelpful. X-ray films of the right hip showed appreciable femoral head deformity with protrusio acetabuli. At surgical exploration a cyst extending posteriorly into soft tissues and containing 150 ml dark fluid was aspirated and excised. The aspirate showed scanty white blood cells on microscopy, and culture, including that for tubercle bacilli, was sterile. Histology of the cyst wall showed only necrotic material.

Case 2

A 61-year-old woman gave a four-month history of discomfort in the right groin associated with a lump that was slowly enlarging. Her history included clinical osteoarthritis of the left hip. Examination showed a firm mass, 4 x 7 cm, lateral to the femoral artery in the femoral triangle. It was

not tender or pulsatile and there was no cough impulse. The results of haematological and biochemical screening were normal. X-ray films showed moderate degenerative changes in both hips. At operation a cystic swelling deep to the muscles in the femoral triangle was excised. The clear fluid contained scanty lymphocytes and was sterile. Histology of the cyst wall showed dense fibrous tissue.

Comment

We believe that our two patients had synovial cysts as a result of hip disease. Synovial cysts are protrusions of synovial membrane caused by synovitis with overproduction of synovial fluid. They occur more commonly with inflammatory arthritis, although one of our patients had osteoarthritis. The joint capsule in the hip has an anteromedial defect and in 15% of all hips there is direct contact with the iliopsoas bursa.¹ When synovitis is active, pressure and necrosis may cause more frequent contact.² The cysts may be asymptomatic but may cause pain, leg oedema by iliac vein compression,³ or bladder compression.⁴ Ultrasound and injection of radio-opaque contrast may help in diagnosis. Two non-surgical methods of treatment successfully used are radiotherapy⁵ and oral corticosteroids.⁴ Both patients were initially thought to have a malignancy. We therefore suggest an alternative diagnosis in patients with hip disease and a lump in the groin.

We thank Mr Kemp at the Royal National Orthopaedic Hospital and Mr Ralphs, Mr Slack, Dr M Corbett, and Dr Boyle at the Middlesex Hospital for permission to report patients under their care.

Requests for reprints should be sent to Dr R A Williams.

¹ Chandler, S R, *Anatomical Record*, 1933, **58**, 235.

² Coventry, M B, *et al*, *Journal of Bone and Joint Surgery*, 1959, **41A**, 721.

³ Scalco, G B, *Angiologia*, 1976, **28**, 7.

⁴ Watson, J D, and Ochsner, S F, *American Journal of Roentgenology*, 1967, **99**, 695.

⁵ Hucherson, B C, and Freeman, C E, *American Journal of Orthopedics*, 1962, **4**, 220.

(Accepted 2 March 1978)

Royal National Orthopaedic Hospital, 234 Great Portland Street, London W1N 6AD

R A WILLIAMS, BSC, MRCP, senior registrar (present address: Department of Rheumatology, the Middlesex Hospital, Arthur Stanley House, London W1N 9PG)

Department of Rheumatology, the Middlesex Hospital, Arthur Stanley House, London W1P 9PG

L J MARKS, MRCP, registrar

Tertiary hyperparathyroidism with adenoma formation after renal transplantation

St Goar introduced the term tertiary hyperparathyroidism specifically to describe the development of autonomous hyperparathyroidism due to a parathyroid adenoma that had arisen from a background of secondary hyperplasia in a patient with chronic renal failure.¹ Nevertheless, the number of reported cases is small.

Case report

A 54-year-old woman with chronic renal failure due to chronic glomerulonephritis had been maintained on home haemodialysis for five years before cadaveric renal transplantation in February 1973. The graft functioned satisfactorily and at six months she had a creatinine clearance of 80 ml/min and was maintained on prednisolone 10 mg/day and azathioprine 75-100 mg/day. Before transplantation the serum immunoreactive parathyroid hormone (PTH) concentration was raised at 1.8 µg/l, but no other abnormalities suggested underlying bone disease. After transplantation minimal hypercalcaemia and hypophosphataemia occurred but serum immunoreactive PTH and calcium concentrations fell within a few months to near normal values. Hypophosphataemia, however, persisted together with a hyperchloraemic metabolic acidosis. The development of Looser's

zones and the histological findings indicated hypophosphataemic osteomalacia, and so phosphate supplements were given.

Fourteen months after transplantation proteinuria of up to 15 g/day was found and a transplant biopsy showed focal proliferative glomerulonephritis. Thirty-six months after transplantation the creatinine clearance had fallen to 55 ml/min and phosphate supplements were stopped. Forty-five months after transplantation plasma calcium concentration rose to 2.85 mmol/l (11.4 mg/100 ml) but two months later had risen to 3.3 mmol/l (13.2 mg/100 ml) when symptoms of hypercalcaemia occurred. Normal plasma concentrations of phosphate, alkaline phosphatase, and hydroxyproline were found, but serum immunoreactive PTH was raised at 2.4 µg/l. The serum 25-hydroxycholecalciferol concentration (25-OHD₃) had been normal (9.2-10.5 µg/l) until the onset of gross proteinuria but subsequently remained very low (<2.5 µg/l). An iliac crest bone biopsy showed nothing abnormal.

At operation four parathyroid glands were identified. Two were normal and two enlarged (1.5 × 1.5 × 1.5 and 1.0 × 0.8 × 0.8 cm). The two enlarged glands and one of the normal-sized glands were removed. Histological examination showed that the largest gland had typical features of a parathyroid adenoma with a rim of compressed normal parathyroid tissue (see figure) and appreciable nuclear pleomorphism. The other two glands showed chief cell hyperplasia with reduction in the normal fat content. The postoperative course was uneventful and both plasma calcium and serum immunoreactive PTH concentrations returned to normal.

Showing part of the adenomatous gland (A) with an apparent rim of normal parathyroid tissue (N). (H and E. × 71.)

Comment

Our patient had severe symptomatic hypercalcaemia 45 months after transplantation, when her renal function was declining. Phosphate supplements may have masked the development of hypercalcaemia or contributed indirectly to the development of tertiary hyperparathyroidism. This is unlikely, however, since hypercalcaemia developed nine months after withdrawing oral phosphate supplements, and hypocalcaemia was not evident during phosphate treatment. Another factor may have been the heavy loss of 25-OHD₃ in the urine in association with severe proteinuria.²

The histological features of the parathyroid glands supported the diagnosis of the true tertiary hyperparathyroidism, which is rare in patients with chronic renal failure or after transplantation. Few cases of parathyroid adenomata developing in a background of hyperplasia have been described.^{1,3} Some have considered hypercalcaemic hyperparathyroidism in patients with chronic renal failure after transplantation to be a manifestation of tertiary hyperparathyroidism.^{4,5} However, histological study of the parathyroid glands in these patients has shown generalised hyperplasia. In most of the patients the state of functional autonomy in the parathyroid glands is presumably due to the considerable increase in the number of secreting