are best disposed of via a WC pan in the isolation suite. Where this is not possible the bedpans and urinals should be bagged within the unit and passed through the exit hatch and taken to the bedpan or bottle washer. Bedpans and bottles should be sterilised by machines close to the washing unit. The only exceptions to the immediate use of the public drainage system for disposing of excreta from infected patients are the enteric fevers, smallpox, Lassa fever, Ebola fever, and Marburg disease. In such cases prolonged contact with a suitable concentration of phenolic disinfectant is essential before disposal.

Dead bodies of patients who have suffered from diagnosed or suspected smallpox, Lassa fever, Ebola fever, Marburg disease, or serum hepatitis should be dealt with by staff wearing gloves and impermeable protective clothing, and should be enclosed in a large plastic bag before being taken from the room in which the patient died.

Waste from susceptible patients does not usually need any special arrangements.

References

- ¹ Lowbury, E J L, *et al*, *Control of Hospital Infection*. London, Chapman and Hall, 1975.
- ² Lowbury, E J L, *British Journal of Dermatology*, 1969, **81**, suppl No 1, p 55.

- ³ Midland Group, *Infection-control Nurses' Association Conference*, Stirling, Maidstone, Kent, Kimberly-Clark, 1976.
- ⁴ Curley, A, *et al*, *Lancet*, 1971, **2**, 296.
- ⁵ Plueckhahn, V D, and Collins, R B, *Medical Journal of Australia*, 1976, **1978**, 819.
- ⁶ Forfar, J O, Gould, J C, and McCabe, A F, *Lancet*, 1968, **2**, 177.
- ⁷ Lowbury, E J L, Lilly, H A, and Ayliffe, G A J, *British Medical Journal*, 1974, **2**, 369.
- ⁸ West Midlands Group, *Contact (Infection-control Nurses' Association)*, 1977, new series No 1, p 14.
- ⁹ Kohn, J, *Lancet*, 1973, **1**, 415.
- ¹⁰ Liljedahl, S O, *et al*, *Journal of Medical Microbiology*, 1972, **5**, 473.
- ¹¹ Hare, R, and Thomas, C G A, *British Medical Journal*, 1956, **2**, 840.
- ¹² Lidwell, O M, *et al*, *Journal of Applied Bacteriology*, 1974, **37**, 649.
- ¹³ Summers, M M, Lynch, P F, and Block, T, *Clinical Pathology*, 1968, **18**, 13.
- ¹⁴ Dineen, P, and Drusin, L, *Lancet*, 1973, **2**, 1157.
- ¹⁵ Bethune, D W, *et al*, *Lancet*, 1965, **2**, 480.
- ¹⁶ Speers, R, *et al*, *Lancet*, 1965, **1**, 478.
- ¹⁷ Maurer, I M, Efstration, A, and Watson, H M, *British Hospital Journal and Social Service Review*, Institutional Cleaning Supplement, May 1962, p 19.
- ¹⁸ Gerken, A, Cavanagh, S, and Winner, H I, *Lancet*, 1972, **1**, 1214.
- ¹⁹ Beard, M A, McIntyre, A, and Rountree, P M, *Medical Journal of Australia*, 1969, **2**, 758.
- ²⁰ Speers, R, and Shooter, R A, *Lancet*, 1966, **2**, 469.
- ²¹ Kelsey, J C, and Maurer, I M, *Monthly Bulletin of Ministry of Health and Public Health Laboratory Service*, 1967, **26**, 110.
- ²² Kelsey, J C, *Lancet*, 1972, **2**, 1301.
- ²³ Public Health Laboratory Service, *Pharmaceutical Journal*, 1971, **207**, 96.
- ²⁴ Lacey, B A, *et al*, *Journal of Applied Bacteriology*, 1965, **28**, 331.

Personal Therapeutics

Childhood epilepsy

D P ADDY

British Medical Journal, 1978, **2**, 811-812

Although the following scheme for the drug treatment of childhood epilepsy expresses my own preferences, it has been arrived at over the years after much discussion with many colleagues who share an interest in the welfare of these children and after consideration of the written views of many others.

Recent years have seen the rise to popularity of some relatively new drugs such as carbamazepine and sodium valproate (the latter having been greeted by some as the greatest thing since Greta Garbo) and a consequent relegation of others such as phenobarbitone and primidone. Available evidence suggests that carbamazepine has an anticonvulsant potency equal to that of phenytoin or phenobarbitone and produces fewer undesirable effects.

Phenobarbitone and primidone are now used less often because of the well-known risk of behaviour disturbance and possible interference with learning in young children. For the prophylaxis of febrile convulsions phenobarbitone is the only drug about which a large volume of information is available and, even so, the evidence about its effectiveness is conflicting.

Phenytoin is an effective anticonvulsant drug that has been in use for a long time and still has many devotees, but its

numerous side effects and the relatively small difference between subtherapeutic and toxic doses make it, in my opinion, inferior to carbamazepine.

Sodium valproate is undoubtedly a very effective drug and, so far, has shown remarkably little toxicity. Some would prefer it to phenytoin but I do not use it as a first-choice drug, preferring to keep it in reserve for the more resistant cases. Nausea and vomiting occasionally restrict its use.

Nowadays, no drug can be said to have had an adequate trial unless adequate serum concentrations have been achieved, and such measurements help to avoid an unwarranted rush into polypharmacy. There is no proof that in epilepsy two drugs are better than one. If it becomes necessary to use a second drug the first should be gradually withdrawn as soon as the epilepsy is controlled. Only exceptionally should it be necessary to use more than one drug in the long term. I offer the following as a practical guide.

Choice of drug

Drugs are listed in order of preference. Use only *one* drug if possible—controlled by estimating serum drug concentrations if necessary.

Major generalised seizures (grand mal)—carbamazepine or phenytoin,* sodium valproate, primidone, phenobarbitone, acetazolamide, or sulthiame.

*In adolescent girls particularly, carbamazepine might be preferred in view of the risk of gingival hyperplasia and hirsuties.

Suggested drug dosages

Drug	Proprietary name	Presentation†	Dose§ mg/kg/day	Average dose at age			Therapeutic serum concentration	
				2 years	7 years	Adult	mg/l	µmol/l
Phenobarbitone*		Tablets, 15, 30, 60, 100 mg Elixir, 15 mg in 5 ml	3-8	15 mg twice or thrice daily	30 mg twice or thrice daily	30-60 mg twice or thrice daily	15-35	60-150
Methylphenobarbitone	Prominal	Tablets, 30, 60, 200 mg	5-10	30 mg thrice daily	100 mg twice or thrice daily	1-200 mg thrice daily		
Primidone	Mysoline	Tablets 250 mg Suspension, 250 mg in 5 ml	10-30	62.5 mg thrice daily	125 mg thrice daily	250-500 mg thrice daily	Phenobarbitone 25-35 100-150 Primidone about $\frac{1}{2} \times \text{Phb}$	
Phenytoin†	Epanutin (Dilantin in USA)	Capsules, 25, 50, 100 mg Tablets, 30, 50, 100 mg Suspension, 30 mg in 5 ml	5-7	15-30 mg twice daily	50-100 mg twice daily	100-150 mg twice daily	10-20	40-80
Carbamazepine	Tegretol	Tablets, 200 mg Syrup, 100 mg in 5 ml	10-20	50 mg twice or thrice daily	100 mg twice or thrice daily	2-400 mg twice or thrice daily	4-10	17-42
Sodium valproate	Epilim	Tablets, 200 and 500 mg Syrup, 200 mg in 5 ml	25-50	1-200 mg thrice daily	2-400 mg thrice daily	3-600 mg twice or thrice daily	50-100	350-700
Clonazepam	Rivotril	Tablets, 0.5, 2.0 mg	0.1-0.15	0.5 mg thrice daily	1.0 mg thrice daily	2.0 mg thrice daily		
Nitrazepam	Mogadon	Tablets, 5 mg	0.5-0.75	2.5 mg thrice daily	5 mg thrice daily	10 mg thrice daily		
Sulthiame	Ospolot	Tablets, 50, 200 mg Suspension 50 mg in 5 ml	10-15	50 mg thrice daily	100 mg thrice daily	200 mg thrice daily		
Acetazolamide	Diamox	Tablets, 250 mg	20-50	125 mg twice daily	250 mg twice daily	500 mg twice daily		
Ethosuximide	Zarontin Emeside	Capsules, 250 mg Elixir, 250 mg in 5 ml	15-50	125 mg thrice daily	250 mg thrice daily	500 mg thrice daily	4-12	300-850

*About 20% of young children will develop a behaviour disturbance on phenobarbitone.

†Note that with phenytoin a small dosage increment may take the serum concentrations from the subtherapeutic to the toxic range. Changes of dose should therefore be cautious and preferably monitored with serum concentrations.

‡With liquid preparations, especially phenytoin, emphasise the importance of shaking the bottle to avoid uneven distribution of the drug.

§Always start with minimum dose and increase gradually.

Minor motor epilepsy—nitrazepam, clonazepam,† sodium valproate,† steroids, or ketogenic (MCT) diet.

Typical absences (petit mal)—ethosuximide, sodium valproate, troxidone, or acetazolamide.

Complex partial seizures (temporal lobe epilepsy) and other focal seizures—carbamazepine, phenytoin, primidone, or sodium valproate.

Infantile spasms—steroids, nitrazepam, or clonazepam.

- (c) Check for hypoglycaemia or acidosis and treat accordingly.
 (d) Diazepam, 100 mg in 500 ml saline or dextrose, or clonazepam, 6 mg in 500 ml saline or dextrose, by constant infusion. (Rate of infusion to be adjusted according to needs of patient.)
 (e) Muscle relaxants and general anaesthesia.

Further reading

A Textbook of Epilepsy, ed J Laidlaw and A Richens. London and New York, Churchill Livingstone, 1976.

Choice of dosage

Presentations and dosage are given in the table. In school-children it is almost always possible to avoid giving a dose during school hours.

Status epilepticus

Status epilepticus should be treated as follows:

- (1) Maintain aeration: oral airway; oxygen by face mask.
- (2) At home: intramuscular paraldehyde 0.15 ml/kg or 1 ml per year of age. (A plastic syringe may be used, provided the paraldehyde is drawn up immediately before injection.)
- (3) In hospital: always have laryngoscope and endotracheal tube ready.
 - (a) Diazepam, 0.3 mg/kg intravenously,
 - (b) Clonazepam, 0.02 mg/kg intravenously. (Can be repeated in 20-30 min if necessary.)
 - (c) Phenytoin, into drip side arm, 15 mg/kg over 15 min, then 2.5 mg/kg every 12 hours (check not having phenytoin routinely).
 - (d) Oral sodium valproate, 50 mg/kg/day via nasogastric tube.
- (4) (a) Combat brain oedema: mannitol 20% 7 ml/kg intravenously over 30 min; or dexamethasone intravenously 0.1 mg/kg immediately then 0.05 mg/kg every six hours.
 (b) Hyperpyrexia: tepid sponge, fan.

†Clonazepam and sodium valproate should not be used together because of the risk of drowsiness and of petit mal status.

WORDS When a new word is needed, a responsibility lies with the designer of the new coinage. There are instances where the design might have been done better. ISOTOPE (1912), from *G isos*, equal + *topos*, place, would have been better as homotope (*G homos*, same + *topos*) and so named because it occupies the same place in the Periodic Table of the elements. ATOPY (Coca, 1923) (*G atopos*, out of place, strange, odd; from *a-*, no + *topos*) was applied to a group of allergic diseases—for instance, eczema, asthma, and hay fever, some of which commonly occur in the same person or in a member of his family. Surely most diseases and syndromes, when first discovered or investigated, equally merit the term atopic on the basis of their being “strange.” Furthermore, when a disease is well understood and no longer strange it is no longer atopic. In any case the term was pre-empted by Sir Thomas More (1551) in his invention of Utopia, no place, that is, an imaginary, non-existent place. While on the TOPIC, topical application of a drug is well described as such because it is applied in the place where it is effective, rather than given systemically to be transported to the desired place by the blood stream. We used to say LOCAL (*L locus*, place), and we still speak of local application and local anaesthetic, but that term stuck before it was generally agreed that three syllables are better than two. And what about topics for discussion and topic teaching? Starting from the title of a work by Aristotle, *Ta Topika, Concerning Commonplaces*, considerations common to many kinds of subjects, thence by a long, tortuous, and barely recognisable route to its present meaning.