Personal Therapeutics

Angina

DAVID SHORT

British Medical Journal, 1978, 2, 939-940

In order to assess the condition four questions must be answered: (1) is the patient really suffering from angina; (2) if so, is it due to coronary disease; (3) are there any aggravating factors; and (4) are there any dangerous features?

Is the patient really suffering from angina?

Attacks of pain or oppression occurring anywhere in the torso—especially in the region of the upper or middle sternum—precipitated by effort or emotion and persisting for a few minutes are virtually diagnostic of angina. Attacks that do not have this precise relationship to effort or emotion are probably due to some less serious cause.

Is the angina due to coronary disease?

A diagnosis of angina implies myocardial ischaemia, but in about 7-10% of cases it is due to non-coronary cardiovascular disease.

Non-coronary causes of angina include aortic valve disease, hypertension, cardiomyopathy, mitral stenosis, and pulmonary hypertension. Occasionally no evidence of heart disease (or any other disease) can be found.

Are there any aggravating factors?

Extracardiac factors that aggravate angina include anaemia, thyrotoxicosis, obesity, smoking, chronic cholecystitis, duodenal ulceration, cervical spondylosis, and anxiety.

Are there any dangerous features?

Coronary arterial disease is notoriously unpredictable. Nevertheless, certain patterns of angina have a particularly serious prognosis—for example, crescendo course, recent onset (especially if abrupt), attacks of unprovoked rest pain, and failure to respond to medical treatment. (In 15-30% of patients angina subsides completely and may not return.)

General management

Remediable factors should be dealt with, and advice given on

how to avoid future attacks.¹ For cases with dangerous features (unstable angina) urgent specialist help¹ should be sought.

Drug treatment

GLYCERYL TRINITRATE

Glyceryl trinitrate is usually sufficient in mild cases. Tablets should be sucked or chewed to relieve an attack and also before doing anything known by experience to be likely to provoke angina.

Side effects—headache and syncope.

BETA-ADRENERGIC BLOCKING DRUGS

Beta-adrenergic blocking drugs greatly help most patients with angina—but not all. Start with a low dosage—for example, propranolol (Inderal) 10 mg twice or thrice daily; double dose every three or four days until relief is obtained or the heart rate falls below 60 beats/min.

Special problems

Cardiac enlargement or an abnormal electrocardiogram—Combine beta-blockade with a diuretic or digoxin or both.

Nocturnal attacks—Slow oxprenolol (Trasicor) may be most effective.

Bronchospasm—Beta-blocking drugs should be avoided or used with great care. Metoprolol (Betaloc, Lopressor) is probably the safest.

Diabetes—Beta-blocking drugs may produce troublesome hypoglycaemic side effects in diabetic patients. Atenolol (Tenormin) and acebutolol (Sectral) are probably the best choice.

OTHER DRUGS

Sorbide nitrate (Cedocard, Sorbitrate)

Sorbide nitrate has a more gradual and in particular a more prolonged effect than glyceryl trinitrate, its action persisting from three to six hours.

Dose—2.5 to 20 mg six-hourly. The tablets are more effective when chewed than when swallowed.

Side effect—headache, which can usually be avoided by reducing the dose; later a higher dose may be tolerated.

Perhexiline maleate (Pexid)

Perhexiline maleate is often strikingly successful in relieving angina that has proved resistant to all other medical treatment.

Dose—100-200 mg twice daily.

Cardiac Department, Aberdeen Royal Infirmary, Aberdeen AB9 2ZB

DAVID SHORT, MD, FRCP, consultant physician

Side effect—occasionally causes peripheral neuropathy and hepatic damage.

Nifedipine (Adalat)

Nifedipine is worth trying in patients with attacks of angina at rest—particularly those of the Prinzmetal type—who have proved resistant to other treatment.

Dose—10 to 20 mg eight-hourly. If a quick effect is needed, the capsule should be crushed.

Since angina often subsides spontaneously, particularly in the summer, it may not be necessary to continue drug treatment indefinitely. If the patient is getting no angina at all, the dose should be gradually reduced, and if the symptom does not return the drug should be stopped and kept in reserve.

Surgical treatment

The operation of saphenous vein bypass is often dramatically successful in relieving severe angina that has proved resistant to full medical treatment, and mortality is low in experienced hands. There is, however, no evidence that it alters the course of ischaemic heart disease or prolongs life.

Every patient who continues to suffer from disabling angina after all remediable factors have been corrected and the medical armamentarium has been fully explored should be seriously considered for operation, particularly if the electrocardiogram and chest radiograph show only slight myocardial damage.

Reference

- ¹ Short, D, *British Medical Journal*, 1976, 2, 98.

MATERIA NON MEDICA

Amid the alien corn

Being a lifelong insomniac I thought I had found the perfect vacation job when I applied for work on the night-shift of a well-known cereal manufacturer. The job lasted 12 weeks and was to be my last foray into the outside world before my clinical training in medicine. Only men worked on this shift, which stretched from 10-00 pm to 6-00 am, and on our arrival we took over the machines from the women with much flirtatious banter on both sides. My first impression, as I fought back the surprisingly deadening fatigue, was the deafening, relentless noise of the machinery and conveyor belts. The place teemed with men and mechanical monsters like a scene from Dante's *Inferno*. My "Virgil" was a junior manager who took me around the enormous building. Boats brought corn along the ship canal to the factory, where it was piped to the upper floors to be processed, checked, and mixed with various vitamin additives. Moving downwards the products were formed into familiar breakfast cereals, and after further checks, they were packeted, weighed on conveyor belts, and loaded into lorries for distribution.

I was in "quality" and therefore made regular checks on various samples; I sat for hours with a large silver spoon in my hand poised over a moving belt of cornflakes. I removed the burnt ones. Sometimes I weighed cartons, sealed packets, or fed damaged packets into a "destructor machine." Occasionally gigantic vats of heated cereal were pushed past my nose so that the smell of cooking cereal lingered on in my nostrils for hours. "Tea-breaks" at midnight, "lunch at 2-30 am soon became routine.

During my sojourn at the factory I rarely saw the sun and became tall, thin, and etiolated, not unlike a stick of celery. Some of my fellow workers had toiled there for years; one confided that he had 11 children. As his wife worked on the day-shift I marvelled at their obvious opportunism.

Driving to work at night, a time normally associated with rest and quietness, I was amidst a sea of men, all heading in the direction of the docks and factories. Even stranger was leaving in the morning, exhausted after the night's work, because then I passed crowds of people yawning at bus stops, their day yet to begin, as I was going home to glorious bed.—R E GOODMAN (general practitioner, Northenden, Manchester).

The rape of the Ar

As I sat in the train listening to the babble of women and trying to read my grammar I became aware of a mechanised voice announcing that the seven fawty-faw would leave from platfawn faw. I wept a little because of the decadence. Even the Frenchies, who never could pronounce a thee and discarded the effeminate aitch long ago, cling steadfastly to their virile ars, doubling and even trebling them before they let them go. But first, in parenthesis, and to mitigate the disappointment of feathered friends who have come so far with me, I ought to explain that rape is not always what it seems to be. The sort I am expostulating about is as in *Rape of the Lock*, by Mr Pope. An American friend of mine, who pronounces her ars quite nicely, tells me that the Pilgrim Fathers took plenty with them in the *Mayflower*,

together with the seed corn and other good things. And to this day they flourish and multiply, enough to make a bridge of kinship between her vast continent and my own West Country from which the pilgrims sailed. If those sturdy puritans now drove cars instead of automobiles I am sure they would call them cars and not cahs. I would love to visit them, and perhaps I will, now that the fez to New Yawk are so much cheapah.

It seems to me that the ar is an essential ingredient of fine oratory, as one may easily prove by reciting a line or two of Milton's "Late Massacre in Piedmont" with and without them, cringing under the whiplash of it when they are allowed their proper force and relapsing into apathy when they are not. I do not dispute the virility of the Chinese, who make do without ars and so, as every schoolboy of the pre-Williams era knew, are condemned to live on flied lice and such wretchedness. They are a special case and always will be, having been created to provide an exception to every rule, like wearing white at funerals and knowing about tea before they were supposed to. Nevertheless, my study of the language of another ancient people leads me to suspect that a country that throws away its ars, having already given away its empire, is well on the road to oblivion. It happens that some dust-besprinkled philological acquaintances of mine are currently peering into the depths of this phenomenon, and publication of their findings is awaited with the usual impatience. In the meantime it would be well for certain clever authors I have been reading lately, who mock the robust diction of our yeomanry, to curb their arrogance, or they may soon look the sillier for it. People tell me that it is impolite of me to say my ars, but I don't much care. I like them.—VIVIAN JONES (St Albans).

WORDS HOMOPHONES are words that sound the same but have different meanings. Some are even spelt the same (homonyms). The context usually conveys the right sense, and confusion rarely occurs. We know that men's underclothing is not being discussed if someone says, "His breath came in thick short pants." But take for example psychosis and sycosis. Which one does he mean? Why! A mental (p)syc(h)osis of course. Mental:chin or mental:mind? Pleural and plural. Secretary: "Did you say lobe or lobes?" Pedantic doctor: "I said the plural." Natal may refer to birth or buttocks. Fissure and Fisher; heel and heal. "Is the liver friable?" means one thing in the postmortem room and another (one hopes) in the kitchen. Genuine confusion can occur in England but not in Scotland between the spoken oral and aural. South of the border both sound like "awrel"; north of it they are pronounced respectively orral, with a short o, and something like owral. Sassenachs, please copy. Alternatively, why not adopt the Greek roots stomal and otic? Difficulty has also been experienced, south of the border, with "inattention" and "inner tension." One near-homophone is every doctor's nightmare, namely the close similarity, especially during quick speech, between hyper- and hypo-. It is just too bad that they have opposite meanings. Here too the situation could be avoided by adoption of the Latin roots super- and sub-. Does supertension sound funny? You would get used to it and in three months forget you ever said hypertension. And anyway, hypertension has a hybrid derivation (as if you cared!).