In My Own Time

Addison's disease

G I M SWYER

British Medical Journal, 1979, 2, 25-26

When I was a medical student Addison's disease was a well-recognised "classic," the first clinical description, never bettered, having been given by Thomas Addison of Guy's Hospital in a paper entitled: "On Anaemia: Disease of the suprarenal capsules," read to the South London Medical Society in 1849. In his treatise of 1855 "On the constitutional and local effects of disease of the suprarenal capsules" Addison also described what is now called pernicious anaemia, but he clearly distinguished this (Addison's anaemia), with its absence of skin pigmentation and prevalence among fat people, from the disease resulting from destruction of the "suprarenal capsules"—in which skin pigmentation was a striking sign and body wasting was more typical than corpulence. Nevertheless, interestingly, *anaemia* was the first of the many symptoms mentioned by Addison in his description, which begins: "The leading and characteristic features of the morbid state to which I would direct attention are, anaemia, general languour and debility, remarkable feebleness of the heart's action, irritability of the stomach, and a peculiar change of colour in the skin, occurring in connection with a diseased condition of the supra-renal capsules." I suspect that I, along with other medical students, was at least for a time puzzled and confused by these two forms of anaemia associated with Addison's name.

The association of Addison's disease with tuberculosis had long been well known, but opinions varied on the prevalence of tuberculous lesions in patients with Addison's disease. Figures of over 70% had been generally accepted, but by the late '30s and '40s less than half the cases were thought to be tuberculous and, with the introduction of antituberculosis drugs, the proportion fell dramatically, and the disease has become a good deal less common. As a result, Addison's disease is now usually ascribed to one of the fashionable group of autoimmune diseases. More rare causes of adrenal destruction were also recognised, such as malignant disease (which usually ran a rapidly fatal course), syphilis, amyloid, and haemochromatosis. Acute adrenal haemorrhage, again usually rapidly fatal and mainly occurring in children with severe sepsis (such as meningitis), was distinguished as the Waterhouse-Friedrichsen syndrome.

Biochemical changes

Low blood pressure and a reduced serum sodium concentration with a raised serum potassium concentration were clearly recognised features, but only in the '30s did Loeb demonstrate that excessive salt loss was responsible. Deficiency of "cortical hormone" was accepted as the cause, the adrenal medulla

playing little or no part in pathogenesis. At that time the study of adrenal physiology was hampered by the rapidity with which most animals died after cortical adrenalectomy and by the want of effective cortical extracts. Nevertheless, the effects of adrenalectomy on the kidney were recognised, together with changes in blood electrolyte concentrations, loss of water, muscle weakness, disturbance of carbohydrate metabolism, lowered intestinal absorption, inhibition of growth, and impaired resistance to stress of all kinds. Only at the end of the '20s were active cortical extracts produced, by Hartman, MacArthur, and Hartman; Rogoff and Stewart; and, finally, in a more potent form, by Swingle and Pfiffner—whose preparation, manufactured on a large scale, was first used for the successful treatment of Addison's disease by Rowntree and Greene in America and by Simpson in England in 1931. Soon after, the importance of increased salt intake was recognised. Meanwhile, working separately in Switzerland and in the USA, Reichstein and his collaborators and Kendall and his colleagues isolated a number of crystalline fractions from crude cortical extracts and characterised them chemically.

Of these fractions, desoxycorticosterone, in the form of the acetate (DOCA), was first used successfully in treating Addison's disease by Simpson in 1938, but it soon became apparent that this was not adequate in all cases. Its action was limited to its influence on salt and water metabolism, the disturbed carbohydrate metabolism remaining unaffected. Another group of cortical hormones—including corticosterone, 11-dehydrocorticosterone, and 11-dehydro-17-hydroxycorticosterone (cortisone)—were known to affect carbohydrate metabolism but not until about 1950 did cortisone become available for treatment.

The cause of pigmentation in Addison's disease was quite unknown in my student days, though theories (since shown to be untenable) were not lacking. Nevertheless, by the late '40s or early '50s it was realised that, though both Addison's disease and Simmonds's disease (hypopituitarism) entailed adrenal deficiency, in the latter, loss of skin pigmentation was often a feature. On the other hand, deficiency of adrenocorticotrophic hormone (ACTH) was part of Simmonds's disease, whereas in Addison's disease loss of the negative feedback effect of cortical hormones (the pituitary itself being intact) resulted in increased secretion of ACTH. Pigmentation, therefore, is related to excessive ACTH concentrations, as was further evidenced by its occurrence in some patients treated with ACTH for non-endocrine diseases. The separation of melanocyte-stimulating hormone (β -MSH) as the actual agent causing increased pigmentation from ACTH (with which it was closely associated) came later. With the development of refined techniques in protein chemistry, the structure of ACTH and β -MSH was defined and their chemical relationships were made apparent. ACTH (of human origin) is a polypeptide with 39 amino-acids, the full biological activity residing in the N-terminal 26 amino-acids β -MSH has an identical amino-acid sequence with the N-terminal 13 amino-acids of ACTH. The current view is that both are derived from a common prehormone β -lipoprotein, of 91 amino-acids, itself biologically inactive but being the precursor of ACTH, β -MSH, encephalins, and endorphins.

The greatest single advance

The availability of cortisone for clinical use in the early 1950s was undoubtedly the greatest single advance in treating Addison's disease that had so far been made. Its effectiveness by mouth was a happy bonus, since previously cortical extracts had had to be given by injection. Its action in restoring normal carbohydrate metabolism is completely reliable, thus eliminating the hypoglycaemic attacks to which some patients, hitherto treated with DOCA alone, were liable. Nevertheless, for treating acute adrenal insufficiency (Addisonian crises) intravenous administration of hydrocortisone (cortisol) remains the route of choice. Though the effect of cortisone on the negative salt balance is limited, most patients with Addison's disease are adequately controlled with cortisone alone, only a minority requiring additional salt-retaining steroid. With the introduction of 9 α -fluorohydrocortisone (fludrocortisone), effective by mouth in remarkably small doses (eg, 0.1 mg per day or less), the complete maintenance treatment of adrenal insufficiency by mouth became possible and with it, adrenocortical ablation as a practical therapeutic procedure. Thus within a matter of 20 years a hitherto inevitably fatal disease, of shorter or longer course, had been brought so effectively under control that surgeons could confidently induce it for therapeutic reasons, knowing that the previously lethal effect of their surgery could be eliminated by appropriate, relatively simple, oral steroid treatment.

Although the isolation and identification of adrenocortical steroids had begun in the mid '30s—and the number of such steroids had reached 28 by 1943—when all the known crystalline steroids had been removed from adrenocortical extract there remained an amorphous fraction which physiological tests showed had great sodium-retaining potency. Many attempts were made to isolate further compounds from the amorphous fraction, and considerable controversy developed over whether

all the known physiological functions of the adrenal cortex could be explained by the actions of the known cortical steroids or whether there was a separate hormone uniquely responsible for sodium and potassium metabolism. From 1950 onwards evidence supporting the existence of such a hormone began to accumulate. Using chromatographic methods, Simpson and Tait, of the Middlesex Hospital, and their colleagues succeeded in isolating from steroids in the blood in the adrenal vein a fraction with 56-75 times the salt-retaining activity of cortisone. This was provisionally called "electrocortin" and was produced on a large scale from beef adrenocortical extract by collaborative work at the Middlesex Hospital and in Switzerland. In 1954 Reichstein and his associates elucidated the constitution of electrocortin, showing it to be corticosterone-18-aldehyde and gave it its present name, aldosterone.

Research from the '50s onwards on adrenal physiology has been worldwide and on a vast scale and aided by powerful technological developments such as chromatography. Since the '60s radioligand assay has also enormously increased our knowledge of adrenal physiology. The pathways of biosynthesis of the adrenal steroids and the enzymes controlling the several steps have been elucidated, leading to an understanding of the pathogenesis of those forms of partial adrenocortical failure collected under the heading of congenital adrenal hyperplasia. The discovery of the intimate association—by way of aldosterone—of the adrenal cortex with cardiovascular renal physiology, including the renin-angiotensin system, has been the product of comparatively recent research. Such has been the pace of the revolution in my time in this part of medical science that few can truthfully claim to be fully au fait with all the modern developments; no doubt even Addison, were he a contemporary, would have found himself hard pressed in striving to remain the authority on "the constitutional and local effects of disease of the suprarenal capsules."

Is there any new treatment for post-traumatic osteodystrophy (Sudek's atrophy) of the foot? Does calcitonin help?

Porcine calcitonin inhibits bone absorption and if it is given very early on or in the transition from the first to the second stage the spontaneous burning pain and, after some weeks, the oedema disappear. It is just as effective as corticosteroids but has fewer side effects. It may also be administered when corticosteroids are contraindicated. In stage three it is ineffective. There is now a salmon calcitonin that is more effective, lasts longer, and is a better choice of drug.

In the light of the controversy over fats and atheroma what types of fat should be used for cooking and what types are used in manufacturing biscuits and cake?

The cook's principal concern should be to use recipes and methods of preparation that require less fat. Unfortunately, this key point has been lost sight of in the controversy over butter and margarine. Fats can be cut down in several ways: by cooking in ways that draw out fat from food, such as grilling, dry roasting, and when frying cannot be avoided the use of a non-stick pan requires much less fat than traditional methods; by substituting fish and chicken for fattier meats more often; by eating less of those foods that are rich sources of invisible fats such as cakes, biscuits, pastries, crisps, and chocolate; and by eating more dishes where bread, rice, pasta, or vegetables are the bulk of the meal rather than the accompaniment. Vegetable oils and soft margarines rich in polyunsaturated fatty acids should be used sparingly in cooking. Not all vegetable oils, however, are low in saturated fats, the exceptions being palm kernel oil and coconut oil, which are highly saturated. Soft margarines rich in PUFAs are sometimes found to be unsatisfactory in baking; low-fat spreads are also unsuitable for cooking because of their high water content. Provided total fat consumption is kept reasonably low, butter or cream may be eaten in moderation. On the other hand, those who are accustomed to a diet rich in fat and who are not prepared to reduce their fat intake should use vegetable oils and low-fat spreads or soft

margarines rich in polyunsaturated fatty acids wherever possible. So far as manufactured cakes and biscuits are concerned, which fat is used will depend on what is most cheaply available. No monitoring of the fatty acid composition of the mix is undertaken and so it is impossible to describe the nature of the fat in these products with any degree of certainty. Manufactured cakes, however, usually contain a little less fat than their home-made counterparts.¹

¹ Paul, A, and Southgate, D A T, In McCance and Widdowson's *The Composition of Food*. London, HMSO, 1978.

What effect does caffeine have on the cardiovascular system?

This depends on drug and patient factors. Caffeine, like other xanthine derivatives, such as theophylline and aminophylline, has stimulant actions on the myocardium. In pharmacological models caffeine increases the force of contraction, the heart rate, and cardiac output. Caffeine also has a direct vasodilator action on coronary, pulmonary, and systemic blood vessels. Xanthine derivatives, however, also have central nervous system effects, including stimulation of medullary, respiratory, vasomotor, and vagal centres. These central actions result in opposing cardiovascular influences, such as slowing of the heart rate and vasoconstriction. The effects of low doses of caffeine on pulse rate and blood pressure therefore vary, though with increasing amounts of caffeine ingested the stimulatory cardiovascular effects predominate. Deleterious effects depend on patient factors such as the presence and severity of associated cardiovascular disease—for example, arrhythmias or coronary artery disease; concurrent drug treatment such as β -adrenoceptor agonists; or of other risk factors like smoking. The report that individuals drinking more than five to six cups of coffee a day are about twice as likely to suffer myocardial infarction has been challenged.¹ A careful study found no independent association between coffee drinking and a first subsequent myocardial infarction.²

¹ Jick, J, et al, *New England Journal of Medicine*, 1973, **289**, 63.

² Klatsky, A L, Friedman, G D, and Siegelman, A B, *Journal of the American Medical Association*, 1973, **226**, 540.