health district and was glad that after the interview I was not faced with the decision of whether to accept the post.

The interview

The interviews are fairly standardised and as the candidates are usually interviewed in alphabetical order I have had the misfortune of always being the penultimate or last one interviewed. Surely drawing lots for the order of interview would be fairer? While the interviews are taking place is a good time for renewing acquaintances, picking up surgical gossip, and studying the opposition. There is usually one person who appears intent on digesting an impressive-looking book—though no page is turned throughout the afternoon—and there is always one who fills the room with tobacco smoke, much to the discomfort of the rest. Conversation tends to be lighthearted but each candidate noticeably quietens as his turn for interview approaches. After it he visibly relaxes but after the final interview the tension returns, conversation becomes more difficult, and the consumption of cigarettes increases.

In my limited experience the members of the interviewing panel have always been most courteous and the chairmen have gone out of their way to settle the candidate. Interviews tend to average 20 to 30 minutes each, though in one place no candidate had longer than 10 minutes. It is remarkable that a job for the rest of one's working life can be settled on such a short acquaintance, particularly as in this place there had been no previous trial by sherry. In my first interview the professor of surgery on

the panel had to be awakened to question me. Perhaps his embarrassment prevented him from looking at me throughout his questioning. The questions are fairly standard—though one friend was asked whether he was a homosexual, to which he replied that he had not realised that it was *de rigeur* for the appointment.

Courtesy to candidates

Several authorities do not acknowledge the receipt of application forms, and some fail to inform candidates that they have not been short-listed for interview. A little courtesy here would relieve the suspense and save many telephone calls requesting this information. Similarly, some authorities could be much quicker in reimbursing expenses incurred in attending interviews.

Any selection procedure is open to abuse and criticism and appointing consultant medical staff is no exception. It is somewhat reminiscent of an assault course and the more practice you have at it the better you become. I hope that my comments may be useful to the organisers of future appointments and to the candidates about to embark on the course. Regrettably, there are many who have had much more experience of this game than I and to them may I quote Rex Lawrie's advice, which he regards as infallible, "that the job they are eventually destined to get will be better than all those they did not."

(Accepted 22 June 1979)

Scientifically Speaking

Person-rem's and the future

BARBARA J CULLITON, WALLACE K WATERFALL

British Medical Journal, 1979, 2, 375-376

Washington, DC—Contemplation of the biological consequences of low-level ionising radiation has become so frequently written up that, in the United States anyway, average citizens are being asked to consider their futures in terms of "person-rem's." While such a unit of extrapolation may be handy for biostatisticians, it saps the accompanying number of relevance to the individual. Some opponents of nuclear power even suggest the latter is the purpose of person-rem's and other univivid ways of expressing risks and damage.

Two new reports along these lines have been released to the public. Each is a portion of a larger report to be issued later. Both bear directly on the kinds of events associated with the

accident on 28 March at the Three Mile Island nuclear-powered electrical generating station near Harrisburg, Pennsylvania. Both were assembled and issued under the auspices of the US National Academy of Sciences.

Reports on risks

One, subtitled "A Critical Review of the Literature," is *Risks Associated with Nuclear Power*. It was undertaken as a contribution to a huge and as yet unfinished study on nuclear and alternative energy systems. The other is the report of the committee on the Biological Effects of Ionising Radiation, known as the "BEIR report," and first issued in 1972. This 1979 version holds to much of the "linear hypothesis" advanced in 1972 to deny a threshold below which radiation doesn't damage life. But the new report has also managed to alienate several of its authors among the committee members because of one aspect or another of that hypothesis.

Neither report, at least in the summaries issued thus far, has material on the Three Mile Island mishap. The *Risks* report contains one footnote, added in proof, specifically mentioning that it was written before the accident. That footnote follows a paragraph beginning: "As regards risks of reactor accidents, the optimists cite some 200-300 reactor years of operation of

5026 Eskridge Terrace NW, Washington DC, 20016, USA

BARBARA J CULLITON, AB, news editor of *Science*, the weekly journal of the American Association for the Advancement of Science

WALLACE K WATERFALL, AB, senior professional associate and director, Office of Communications, Institute of Medicine, National Academy of Sciences

commercial power reactors in the United States alone—plus additional experience with naval reactors—without a significant accidental release of radioactivity, much less an identifiable injury to a member of the public; the pessimists point to the conceptual possibility of an accident that will ultimately entail many tens of thousands of deaths.”

Risks accompanying nuclear-power production, in conditions of less than catastrophic mishandling of the plant, are much the same matters of concern that occupy the BEIR committee: low level radiation. Hence before getting into the possible consequences of “large unplanned releases of radioactivity,” as the *Risks* report puts it, a great deal must be reviewed about radiation-effect measurement in various units of absorbed dose (rad), roentgen equivalent in man (rem), the quality factor that makes an exposure of 1 rem in x-rays climb to 10 rem if the radiation is alpha particles from a radionuclide, and the importance of the time over which a dose is delivered.

The *Risks* report, being a survey of published work, can finesse the linear hypothesis problem—and does, by reporting that various “respected” members of appropriate scientific groups do and do not believe that the plot of radiation dose against probability of carcinogenesis is a fairly straight line down to the origin. Nevertheless, after some hemming and hawing, the report tends to fix on “an augmented probability for ultimate cancer death equal to 1/5000th the dose equivalent in rem, if the body is uniformly irradiated and if one averages over all ages and other vital factors in the US population.” Put another way, a fatal case of cancer is to be expected from among 5000 persons exposed to a 1 rem whole-body dose of radiation—that is to say, 5000 person-rem.

On that yardstick, say the authors of the *Risks* summary, the natural background radiation in the US (cosmic-ray exposure, rock emissions, and the like), plus the almost equal exposure of the populace to medical irradiation, plus old fallout from weapons tests, would be regarded as responsible for 7000 cancer deaths of the yearly national total of 360 000. Those 7000 cancer deaths are identified as “unrelated to nuclear power.”

Assuming that nuclear power plants perk along uneventfully until the end of the century, growing in number according to projections that seemed likely until recently, the *Risks* report provides figures that can readily be multiplied to form an estimate that radiation from normal nuclear power plant operation will cause 1950 cancer deaths in the last quarter of this century. Indeed, one newspaper reporter made those multiplications, rounded off the fatalities to 2000, and was publicly rebuked by the chairman of the committee that issued the *Risks* report—mostly for failing to point out that the death rate was lower than that associated with generating electricity from coal, and that the total US cancer deaths would be about nine million by the year 2000.

One possible conclusion is that people who assess risks and make projections about phenomena as little understood as the biological effects of low-level radiation, must introduce so many hedges as to make impermissible the examination of a single consequence. The *Risks* report may, perhaps, be best excerpted for its assessment of the risk of reactor accidents: “the one whose range of uncertainty extends to the largest values. . . .”

Conflict of evidence

The BEIR report does not have to be so concerned with the source of low-level radiation as with its effects. Rather oddly, however, it pegs natural background radiation at 100 millirems per year. The *Risks* report said it was 80 mrem. An interagency task force of the Department of Health, Education, and Welfare said in March that it was 130 mrem. Such are the figures with which committees must deal.

For the risk of genetic effects of low-level radiation, the BEIR group, even though using a new method of estimating, came up with figures very similar to the 1972 report. One rem of parental exposure throughout the general population in the first genera-

tion is predicted to cause an increase of 5 to 75 additional serious genetic disorders per million live births. A 1-rem exposure received in each generation is estimated to result, at genetic equilibrium, in an increase of 60 to 1000 serious genetic disorders per million live offspring. The relative importance of this last bracket can be gauged from current estimates that the incidence of serious human disorders of genetic origin is about 107 000 per million live births.

On somatic effects, the BEIR committee lost its cohesiveness when it came to low-dose radiation that is also low in linear energy transfer—low-LET—as x-rays are. The somatic effects subcommittee seems to agree that the total carcinogenic effect for high-LET radiation essentially is proportional to dose in the low-exposure range. But, said a dissenting report by five of the 16 subcommittee members, “in contrast, the dose-effect relation for low-LET radiation is very unlikely to be linear” for low doses. The majority of the BEIR panel members, said the minority, was overestimating the cancer risk of low-dose low-LET radiation by 10 times or more. “There is far too little theoretical information to serve as a reliable guide for extrapolation,” said the minority.

What the minority was disagreeing with were extrapolations by the majority such as these:

(1) A lifetime risk of cancer incidence induced by low-LET radiation in a single whole-body exposure to 10 rads is increased 1.2 to 4.5% of the naturally occurring rate for women, and 0.8 to 3.2% for men. The mortality rate for that exposure is raised 0.5 to 2.7% for women, and 0.4 to 2.0% for men.

(2) A risk of cancer incidence from continuous lifetime exposure to 1 rad per year poses an increase above normal of 8.4 to 32.6% for women, and 5.2 to 17.9% for men, with the corresponding mortality estimates being 16.9% for women and 2.7% for men.

Most recently, the National Academy of Sciences has acknowledged that “a restatement of a small portion of the report may result in a more balanced expression of the differing interpretations,” which would seem to predict a rewrite of the already released new BEIR report conclusions.

Studies at Three Mile Island

Meanwhile, in the real world of the citizenry, the accident at Three Mile Island appears to have raised all sorts of issues. For the two million residents within 50 miles of the plant, Government officials are acknowledging the possibility of one “excess” cancer death, which would seem to indicate a possibility of the 5000 person-rem condition having been fulfilled. Studies planned for the area include a census of the 50 000 residents within five miles of the plant, in an effort to identify people who were there during the radioactive gas release, but have since left. A study will also be made of pregnant women within 10 miles of the plant, and of their infants during the first few months of life. Another investigation will be made of the health of the plant workers. The study expected to turn up some of the most interesting information is to be conducted by the National Institute of Mental Health: plumbing the strain on persons living in fear of radiation exposure.

Some sequelae of the reactor accident are less grim: a Government agency investigating the event reported that a plant operator whose instruments should have indicated a lack of cooling water couldn’t see the instruments: he was “a big man with a large belly that hung over the instrument panel.” The University of Mississippi supposedly found high levels of ²²Na in toads, frogs, and a deformed lizard taken from a site of underground nuclear tests in that state during the 1960s. A dozen families near the site were advised to leave their homes—until it was discovered that a university petri dish was contaminated with radioactive sodium. The Tennessee Valley Authority, a quasi-governmental agency whose power generating stations include three fueled by uranium, announced that it would subject the operators at those plants to intelligence tests.