SHORT REPORTS

Progressive depletion of plasma cholinesterase during daily plasma exchange

Suxamethonium is used extensively in anaesthesia and intensive care to facilitate rapid endotracheal intubation. It is hydrolysed by the plasma enzyme cholinesterase, and the duration of action of the drug is thought to be prolonged when the enzyme concentration is less than 20% of normal.¹ Since a single four-litre plasma exchange removed 64% of the plasma cholinesterase,² we have examined the effects of daily plasma exchanges on the circulating cholinesterase concentration.

Patients, methods, and results

Two patients who received daily plasma exchanges for the treatment of Goodpasture's syndrome (15 days) and Wegener's granulomatosis (8 days) were studied. Four-litre exchanges were undertaken with a Haemonetic Model 30 blood processor and the plasma replaced with plasma protein fraction (PPF). Venous blood was collected at the end of each exchange for determining the plasma cholinesterase concentrations.²

The figure shows that repeated daily plasma exchanges progressively decreased the cholinesterase value to that of the infused PPF (100-200 U/l). The increases in cholinesterase concentration observed in case 1 after exchanges 6 and 13 were associated with the administration of whole blood together with PPF to replace the plasma removed.

Effect of daily plasma exchanges on plasma cholinesterase concentration.

Comment

Regular plasma exchanges carried out at intervals of one to four weeks are an accepted and safe form of treatment for diseases such as the paraproteinaemias and hypercholesterolaemia. Intensive daily exchange has been introduced recently as part of the treatment of fulminating antibody-mediated and immune-complex diseases. There is little information available about the effects of this regimen on blood constituents other than a decrease in immunoglobulins and clotting factors together with a moderate defibrinogenation.³ Healthy donors undergoing daily plasma exchanges of only 1 litre for five days showed a mean decrease in total protein concentration of 18.4 g/dl.

We have shown that daily plasma exchange caused virtually complete removal of circulating cholinesterase in patients who started treatment with the low values of 1267 and 844 U/l (normal mean concentration 3637 U/l (SE of mean ± 226), $n=22$). We recommend that drugs metabolised by cholinesterase, such as suxamethonium and procaine-like esters, should not be used in patients undergoing

intensive daily plasma exchange so that their duration of action is not grossly prolonged.

We thank Professor D K Peters for permission to study the patients.

¹ Vickers, M D, Wood-Smith, F G, and Stewart, H C, *Drugs in Anaesthetic Practice*, 5th edn. London, Butterworths, 1978.

² Wood, G J, and Hall, G M, *British Journal of Anaesthesia*, 1978, **50**, 945.

³ Pinching, A J, *British Journal of Anaesthesia*, 1979, **51**, 21.

⁴ Kliman, A, *et al*, *Blood*, 1964, **23**, 647.

(Accepted 8 June 1979)

Department of Anaesthetics, Royal Postgraduate Medical School, London W12 0HS

JENNIFER L PATERSON, BSC, research technician

ELEANOR S WALSH, MB, FFARCSI, senior registrar

G M HALL, PHD, FFARCS, senior lecturer

Occupational handling of chemicals preceding Hodgkin's disease in men

In 1976 Olin¹ reported an increased death rate from Hodgkin's disease (HD) among Swedish chemistry graduates. We report an unduly high prevalence of occupations characterised by the handling of chemicals in men with HD.

Material, methods, and results

From tables describing the occupations of the working population of Sweden² we selected occupations indicating a high degree of exposure to chemicals (see table).

Patients—We examined the case records of all men aged 20-65 years (median 36) diagnosed and treated for HD during 1973-8 at this department and recorded their occupations at diagnosis, ages, and histological classifications. We thus studied 88 patients.

Controls—Control patients comprised three groups. In group A the case records of 100 consecutive men aged 20-65 years (median 43) with non-malignant disorders treated at the department of internal medicine in 1977 were examined and occupations recorded. In group B the case records of 100 consecutive men aged 20-65 years (median 39) treated at the department of allergology in 1977 were examined in the same way. In group C the occupations of 32 men in the same age group (median 52) with chronic leukaemias (10 with chronic lymphocytic and 22 with chronic myeloid leukaemia) seen during 1969-77 at the department of internal medicine were recorded. The recording of occupation was similar in all departments. Because only men aged 20-65 years were included in the study an occupation was obtained for all the patients and controls.

Of the 88 men with HD, 17 (19%) had occupations indicating that they handled chemicals (table). The corresponding figures in the two control groups with non-malignant disorders (A and B) were 2% and 3% respectively. This difference between the patients with HD and each of the control groups was significant ($P=0.0001$ and 0.0002 ; Fisher's exact probability test). Of the men with chronic leukaemias (group C), one (3%) had an occupation indicating handling of chemicals. This was also significantly different from the patients with HD ($P=0.02$). In Sweden in 1976³ about 64 000 out of 2.2 million working men (2.9%) had occupations listed in the table.

Only two out of 35 patients (6%) aged 20-30 years were occupationally exposed to chemicals, compared with eight out of 31 (26%) aged 31-50 years. The corresponding proportion for the patients aged over 50 years was seven out of 22 (32%). Histological classification did not differ significantly between the exposed and unexposed patients.

Comment

Of the men with HD, which was diagnosed between 20 and 65 years of age, about 20% had occupations indicating that they handled chemicals. In the three control groups the corresponding figure was only 2-3%, which agrees well with the prevalence (2.9%) of such