In My Own Time

General practice

C A H WATTS

British Medical Journal, 1979, 2, 1055-1056

I qualified in 1934 and was lucky enough to get two house appointments in my teaching hospital in surgery and medicine. I was paid £50 a year, all found, for this invaluable experience. In those days it was a common saying that the backbone of good general practice was sound obstetrics, and it was my ambition to become a GP expert in this specialty, but I found that I had to wait for three months before starting a job at the maternity unit. It was autumn and holiday locums were getting scarce, so when I was offered an assistantship which was to last until the end of the year I accepted the post with alacrity.

Thus it was that I found myself working in a mining community with incredibly meagre medical facilities. Before I finally settled down, I had worked as a locum or an assistant in several practices; some were good, others bad, but this one was quite the worst. I was so dismayed by the conditions that I would have left—but in those days one felt bound to honour any agreement, and the post was for only three months.

Primitive conditions

The surgery set-up was astonishing. There were rows of seats to accommodate about 60 patients, who all sat facing a long high bench reminiscent of a bank counter. Behind this contraption I stood between my two principals. Behind us was a smaller but similar bench, and this was presided over by a dispenser, with most of her wares on shelves at the back of the room. The doctor called for the next patient to come forward, and, having listened to the complaint, he turned to the dispenser to order the appropriate remedy. There was rarely any attempt to examine anybody, unless some obvious pathology like a septic hand or some surface with a rash was proffered. If the patient complained of a cough, one could lean across the bench and apply the stethoscope to the exposed manubrium sterni as a token of investigation. There was one examination room for the three of us. It was rarely used and, as a very junior doctor, I was most unpopular if I wasted time by occupying this inner sanctum. With this method of work, we got through a great many patients at each surgery session, and these took place twice a day, six days a week and on each Sunday morning. Even on Christmas morning the three doctors solemnly assembled to see patients. Once the surgery was over, I was on duty every third Sunday. I did have a half-day during the week which started after the morning surgery—once I had finished my visits. These usually numbered about 50, but there could be twice as many during an epidemic. I recollect that my record was 94. Visits were made on a bicycle, and, as the streets were not macadamised, this meant riding on the flagged

pavement. More than once I bowled someone over who had emerged too fast from her home, and once I fell off into the mud of the roadway when a housewife inadvertently threw a bucket of ashes over me as I was passing. Such incidents were cheerfully accepted as normal hazards of life.

Diphtheria was endemic, and every sore throat was viewed with suspicion. If there was any possibility of the disease, a throat swab was taken. I carried a few sterile tubes in my coat pocket. The specimen then had to be taken to a laboratory at the local hospital, and one collected a phial of antiserum. The streets of the town were then combed for the partner with the 20 ml syringe, as there was only one suitable instrument in the practice. If the possessor had already used it, one had to take it back to the surgery to sterilise it by boiling. This took time because one had to start from cold: the heavy glass syringe would crack if it was plunged into hot water. The whole process of caring for a possible case of diphtheria could take up an hour or more of a busy day, but it was well worth the trouble. The giving of serum was one of the few active treatments available to the family doctor. If it was given within 24 hours of the onset of the disease, the results were excellent. If the injection was delayed, there was every chance of serious complications, and a mortality of 20%. Both this fever and scarlatina were rife in the undernourished miners and their families, and patients suffering from these conditions were taken away in a yellow fever van to the local isolation hospital. Every community of any size had such an institution. It was separated from the township by a few green fields, and the victims were cared for until they had recovered or died. Many did die. The stay in hospital was for at least six weeks and could be twice as long. No visitors were allowed, and this was a terrible ordeal for the children.

Lobar pneumonia was not uncommon. It occurred in all age groups. Often fatal among the aged, it was called the old man's friend, or, alternatively, the captain of the men of death. The family doctor was impotent to help by way of drugs, his main task being to support the morale of both the patient and the family by frequent visits. Until the patient was out of the wood, we called from three to five times a day; such attention was greatly appreciated, especially when the visit was made last thing at night, or before the morning surgery. It was a wretched illness for all concerned. The incessant cough and pleuritic pains were aggravated by a devastating sense of weakness and an inability to sleep. With the more fortunate patients, there was a crisis about the seventh day. The temperature fell, and at last the patient was able to sleep, to wake up looking and feeling a great deal better. This experience suffered by family and doctor together forged a splendid bond between them, and this was one of the things that made general practice so well worth while.

Deaths and emergencies

The most dreaded of maladies was pulmonary tuberculosis. Some families were especially vulnerable to it, and it tended to

Ashby de la Zouch, Leicestershire LE6 5GZ

C A H WATTS, MD, FRCGP, retired general practitioner

strike at young people. The onset was insidious, and the course could be lingering or extremely rapid. I recall seeing a girl of 16 with her mother; she had missed a few periods. In those days it would have caused the greatest offence to probe into the possibility of pregnancy. The subject of sex was delicate, and had to be approached with the greatest tact. The girl looked pale and thin, and she was one of the few patients I took into the examination room for a check-up. Beyond her pallor there were no signs of serious disease, so she was given some iron medicine. A week later I was sent for as she had developed a tiresome cough, and had taken to her bed. She seemed thinner than ever and looked really ill. A careful examination of her chest disclosed nothing of note, but I did manage to secure a specimen of her sputum. The pathology report indicated that it was loaded with tubercle bacilli. She was dead in a month from what was called galloping consumption. I was conscience-stricken at not having suspected this trouble at my first consultation; amenorrhoea was high up on the list of symptoms for this dread disease. Even so, I realised that there was nothing I could have done.

There were always beds for the acute infectious diseases in the fever hospitals, but accommodation in a tuberculosis sanatorium was in short supply. Patients had to wait for months for a vacancy. My feelings of guilt were not allayed by the great gratitude of the family for all the care and attention I had lavished on the girl. Death among young people and children was no rarity. Lung cancer was never diagnosed. If it occurred, it went unrecognised. I remember one man in his 50s dying from what was called a pulmonary abscess. He certainly produced masses of sputum which was free of tubercle bacilli. With hindsight, I have since wondered if he could have had a bronchial carcinoma.

In this, my first real glimpse of an industrial general practice, I never saw any evidence of antenatal care. The district midwife delivered most of the babies, as only a minority could afford the luxury of having a doctor at the confinement; he was called in only if the midwife ran into difficulties. Few family doctors had any special training in the art of obstetrics; indeed many

newly qualified doctors went straight into practice after taking finals without any postgraduate hospital experience at all. At the expense of their first patients they learnt practical midwifery the hard way by trial and error, and some became real experts. The confinement could be protracted, a good deal of reliance being put on what was called "masterly inactivity."

I well remember one woman who was finally delivered successfully after a four-day labour. One night I was sent for by the local midwife to help in an obstetric case. She assured me that forceps were now indicated. The idea of operating without any experience terrified me. The only midder bag in the practice had complicated forceps that I had never seen before, and there was no chloroform. My courage failed me, and I sent for one of the principals. He was tactful and kind, and commended me for seeking his help. After a quick examination, the patient was turned on her side, and the nurse held up a leg. The forceps were deftly applied, there was a fearful scream from the mother, and the baby was born yelling its head off. I was most impressed by his skill, but I hated the brutality of the operation. This mother was successfully delivered. Many were not so lucky. If called on the GP felt honour bound to tackle the most daunting procedures like midcavity forceps, the breech birth in primips, or the manual removal of the placenta. Most doctors used chloroform as an anaesthetic, but like my principal, some felt it was safer and quicker without it.

The suffering of so many patients was so commonplace that the family doctor had to be tough to get on with many of his more interesting and rewarding tasks, such as setting a simple fracture, reducing a dislocation, or lancing a boil. He had to be cruel to be kind. The masses of working-class people in the '30s did not expect to be comfortable. Most went hungry, and their undernourished children all showed evidence of rickets. They were miserably cold in the winter, unless they were roasting in front of the coal fire in the kitchen. Pain and discomfort were accepted as part of life to be endured with stoicism. Successful treatment by the family doctor was accepted with gratitude, and the many failures were tolerated as inevitable, without rancour or recrimination.

A woman with one normal child has just given birth to an infant with a grossly deformed heart. What are the chances of her having another such infant?

The risk of recurrence after the birth of a child with a congenital heart malformation varies between 1% and 4%, depending on the type of malformation. Usually the commoner malformations have the higher risk of recurrence. Early prenatal diagnosis of congenital heart defects is not yet available.

Zetterquist, P, *A Clinical and Genetic Study of Congenital Heart Defects*. Sweden, The Institute for Medical Genetics of the University of Uppsala, 1972.
Nora, J J, McGill, C W, and McNamara, D G, *Teratology*, 1970, 3, 325.

Is breast-feeding contraindicated if the mother is taking warfarin?

There has been controversy about warfarin in breast milk. But the studies of L'E Orme *et al*¹ and de Swiet and Lewis² showed conclusively that either warfarin is secreted in such trivial amounts in breast milk that it could not possibly do any harm or that none at all passes into the breast milk. Phenindione, however, taken by a breast-feeding mother may harm the baby.

is a true depression, which must be recognised and treated accordingly. Most, often young and previously fit, have physical and psychological symptoms including mild recurrences of the original illness for which there is no satisfactory treatment. This puts a great strain not only on the patient and his or her relatives but also on the doctor, whose job it is to emphasise that the symptoms will disappear after perhaps as long as 6-12 months (it may help to give an approximate date—"about Christmas," "the end of the summer") and that they do *not* imply continuing damage to vital organs. It is best to keep such patients at work or otherwise occupied if possible. Tonic mixtures containing such things as iron, arsenic, and strychnine or tonic wines are a support to both patient, for whom they can be prescribed, and doctor.

What is the use and efficacy of alpha-chymotrypsin?

Alpha-chymotrypsin is a proteolytic enzyme obtained from the bovine pancreas by aqueous acid extraction of its precursor, chymotrypsinogen, and subsequent conversion to chymotrypsin. Use of this enzyme as an anti-inflammatory agent or in the treatment of ulcers is not based on any substantial evidence of its effectiveness.^{1 2} There may be a place for chymotrypsin in thoracic surgery where it has been used to promote liquefaction of secretions of the upper respiratory tract. In ophthalmology the zonulolytic dissection of the lens with chymotrypsin (a 1 in 5000 or 1 in 10 000 solution in a sterile diluent) has facilitated intracapsular cataract removal.³ Chymotrypsin is highly antigenic and severe allergic reactions have occurred after its intramuscular injection, so a sensitivity test before injection is advisable. Nausea, vomiting, diarrhoea, and skin eruption also occur after oral administration.

¹ Orme, M L'E, *et al*, *British Medical Journal*, 1977, 1, 1564.

² de Swiet, M, and Lewis, P J, *New England Journal of Medicine*, 1977, 297, 1471.

What treatment, if any, is advised for the sometimes prolonged mental and physical debility that occurs after "non-specific" virus infection?

I am not sure what is meant by "non-specific." Many specific virus infections are followed in a few patients by debility; the best known are influenza, hepatitis, and glandular fever. In a few patients there

¹ *The Medical Letter*, 1975, 17, 41.

² *Federal Register*, 1977, 42, 58739.

³ Kirsch, R E, *et al*, *Archives of Ophthalmology*, 1964, 72, 612.