Procedures in Practice

PHILIP H BEALES
M J AL-KHALED

LARYNGOSCOPY

Laryngoscopy is the method of examining the larynx from above by direct vision. It should always be preceded by an inspection of the exterior of the larynx in the neck, and of the movements on deglutition, when the larynx normally moves upwards. The larynx moves downwards on inspiration in laryngeal stenosis and is immobile in tracheal stenosis. Palpation of the larynx in the neck will disclose any broadening and tenderness that may indicate perichondritis of the thyroid cartilage.

Indications

Indications
Persistent hoarseness
Stridor
Thyroidectomy
Dysphagia
Earache
Contraindications
Acute inflammation of the throat causing trismus
Acute epiglottitis

Diagnosis—The commonest symptom of laryngeal disease is hoarseness or change in the normal voice, and if this persists for more than two weeks after medical treatment the larynx must be examined. Other indications include stridor and examination of the movements of the vocal cords in patients before thyroidectomy. In patients with dysphagia examination of the larynx may disclose a postcricoid carcinoma, and earache may be a result of referred pain from a carcinoma in the laryngeal area.

Treatment—Simple lesions such as cysts and vocal-cord nodules may be removed with special instruments by using the method of direct laryngoscopy, and biopsy of a suspicious lesion in the larynx is an important procedure. It must be remembered that treatment of early carcinoma of the vocal cords is curative.

Contraindications to laryngoscopy—Contraindications are few, but the procedure cannot be performed in acute inflammations of the throat that give rise to trismus—for example, a peritonsillar abscess. Direct laryngoscopy is inadvisable in acute epiglottitis, as it could lead to a spread of inflammatory oedema to the glottis.

Indirect laryngoscopy

This is the method of examining the larynx from above with a laryngeal mirror. It is a simple procedure that is within the capabilities of every doctor who is willing to practise, and should be performed more often than it is.

Instruments—A laryngeal mirror or mirrors and a spirit lamp, or hot water if a spirit lamp is not available; gauze swabs for holding the tongue; and a light source, which may be a battery-operated head lamp or a head mirror using reflected light that may come from an electric bulb or a good torch.

Procedure—The patient is seated in front of the examiner, who sits at about the same height. The light is directed on to the lips of the patient, who is asked to open his mouth and protrude his tongue as far as possible. The tongue is grasped between the thumb and middle fingers of the examiner's left hand through the gauze swab. It is important not to pull the tongue too far forward or the frenum will impinge on the lower teeth and cause pain. The index finger of the left hand is rested on the upper teeth to steady the hand.

The mirror is warmed over the spirit lamp, the flame being directed on to the glass surface. After the temperature has been tested on the examiner's cheek the mirror, held in the right hand, is introduced into the mouth and placed at the back of the uvula without touching the tongue, as this would smear the surface of the mirror. Steady pressure is maintained on the soft palate, and the light is directed on to the mirror. A reflection of the interior of the larynx will be seen, and the various structures may be examined in sequence by tilting the mirror.

The base of the tongue, epiglottis, valleculae, aryepiglottic folds, false cords, true cords, and upper tracheal rings can all be seen in turn. The mobility of the cords can be tested by asking the patient to say "ee," when the glottis should close. The mirror may have to be removed and warmed again and the examination repeated before the full information is obtained.

5% Cocaine solution
or
Amethocaine lozenge

Difficulties of indirect laryngoscopy—Some patients have an overactive "gag" reflex, and examination is difficult unless the soft palate is sprayed with an anaesthetic solution such as 5% cocaine or an amethocaine lozenge is sucked before the examination. It is difficult to examine children by indirect laryngoscopy, but sometimes the larynx may be seen by placing the mirror almost horizontally against the hard palate instead of the soft palate, so that the gag reflex is not excited. The anterior commissure of the larynx is the most difficult part to see, and complete examination of the piriform fossae is not possible by indirect laryngoscopy.

It should be remembered that the structures of the larynx are seen in a mirror, so that the anterior part points away from the examiner and the right-hand structures are seen on the left.

Direct laryngoscopy

In direct laryngoscopy the larynx is looked at from above through a tube with an integral light source; there is no distortion or shortening of the structures, as occurs when a mirror is used, and the image is not reversed.

Surgical instruments may be passed down the laryngoscope and limited surgery carried out, and accurate biopsy is possible. It is the method of choice in children.

Contraindications—Contraindications are few and would include injuries and disease of the cervical spine; severe trismus; and appreciable laryngeal obstruction, when a tracheostomy should be done first.

Method—Under general anaesthesia with a hard, narrow-bore catheter inserted through the mouth into the trachea the patient rests on his back with his head flexed against his neck and his teeth protected by a guard. When the muscles are relaxed the laryngoscope is inserted, and the structures in the vicinity of the larynx and the larynx itself are inspected so that all the structures are seen. The laryngoscope can be fixed in position by suspension attachments to the chest of the patient or the head of the operating table, allowing the surgeon two hands to perform endolaryngeal surgery.

Micro-laryngoscopy—This method was first described by Kleinsasser in 1968. A special wide laryngoscope is used, held in position by a suspension system. The larynx is examined with the Zeiss operating microscope, which gives a magnified binocular view of the larynx. Fine instruments are available for accurate surgical treatment of such lesions as polyps, singers' nodes, etc, and accurate biopsy of suspicious lesions is possible. This is now the method of choice for treating lesions of the larynx endoscopically.

Complications of direct laryngoscopy—These are few and include damage to the teeth and laceration of the lips and pharyngeal wall.