The First Year of Life

H B VALMAN

WEANING

Weaning is the process of getting the baby used to eating foods other than milk and using a spoon and cup. For the first four months of life the baby will need only milk and additional drinks of boiled water. New foods should start to be introduced at about 4 to 6 months. If solid foods are introduced too early the baby may become too fat; if they are introduced too late there may be problems with chewing. Once introduced, solid foods should be given regularly and in gradually increasing quantities. As the amount of solid food increases the number or size of milk feeds should decrease.

Mothers need a small wide plastic teaspoon with no sharp edges, a small plastic cup or bowl, a feeding beaker, and a cotton or pelican-type bib. The feeding equipment should be kept in a sterilising solution. If a metal spoon is used it should be scalded in boiling water and then allowed to cool before being used.

Firstly, the baby has to be taught to take food from the spoon rather than just sucking. He should also be given plenty to drink to replace the milk: at least 100 to 150 ml of water or diluted fruit juice each day. Most fruit *squashes* are unsuitable for babies.

4 to 6 months

6-8 am	Milk
10 am	Milk Baby cereal
2 pm	Milk Fruit or vegetable puree
6 pm	Milk
10 pm	Milk

At 4 to 6 months the baby will usually be having breast or bottle feeds at 6-8 am, 10 am, 2 pm, 6 pm, and 10 pm. Solid foods should be introduced by being fed to the baby before the mid-morning or lunchtime feeds—usually a baby cereal or a fruit or vegetable purée.

Of the many cereals available the high-protein varieties should be avoided. The mother should start by using one made from rice and should mix half to one teaspoonful of cereal with one to two tablespoonfuls of milk and feed it to her baby from the spoon. Sugar should not be added. At lunchtime the baby should be fed with fresh vegetables, such as potato or carrot, or fruit such as apple. The vegetables should be boiled and the fruit stewed and then liquidised, mashed into purée or sieved. Feeding should start with half or one teaspoonful before or during the lunchtime feed and gradually increase to about four teaspoonfuls a day.

At about five months mothers should start to give a wider variety of foods. Banana mashed with a fork is a favourite of many babies. Egg yolk, other cereals, soups, milk puddings, and strained baby foods should all be tried, with the emphasis on savoury flavours rather than sweet ones.

Preparing food

Home-prepared foods are best. The mother knows what is in them and they can be prepared easily and cheaply as she can use a small portion of food cooked for the rest of the family. Many families have an electric liquidiser, but a hand blender is equally effective. If there is a deep-freeze time can be saved by preparing puréed food in bulk and individual portions can be stored for up to a month.

6 to 7 months

6	am	Milk feed
9-10	am	Cereal + milk, or Scrambled or boiled eggs Milk feed
1-2	pm	Minced /pureed meat or fish, pureed vegetables and potato, gravy Diluted fruit juice or milk feed
3-4	pm	Dilute fruit or water
5-6	pm	Fruit puree / mashed banana Custard or milk pudding Milk feed

At 6 to 7 months a greater variety of tastes and textures should be introduced. A little finely minced chicken, lamb, beef, or liver may be given mixed with a potato purée. White fish can be boiled with milk and then, once the bones have been removed, mashed in with some potato or vegetable purée. Grated or cottage cheese may be given, mixed in with other foods, and lightly boiled or scrambled eggs should be introduced.

The timetable of feeds should also start to change at this age, and the number of milk feeds should fall.

Babies often reject food. This may be because they are not used to a new taste or texture. But it may also be because they are thirsty or because the food is too hot, or simply because they want to attract attention. Mothers should be told to keep trying with new foods and to give them in different ways. If the baby becomes constipated he should be given more fruit and vegetables and water.

About 8 months

The baby can now sit in his high chair at the table with the rest of the family. Two-course meals at lunch and teatime should be introduced—for example, a savoury food followed by a fruit purée. Also food should be minced less finely, and potatoes mashed rather than puréed. The baby should also be given hard foods such as a crust of bread or sponge finger biscuit which he can hold in his hand and this will encourage him to chew. These should always be given under supervision to make sure that he does not choke. After 6 to 7 months ordinary cows' milk can be given to the baby and he should be encouraged to take it from a beaker or cup. In homes with good hygiene there is no need to boil the milk if it is kept in a refrigerator. Infants receiving ordinary cows' milk should have supplementary vitamin D until the age of 2 years.

9-12 months

+ Vitamin D

At this stage the baby should be able to eat the same food as the rest of the family, though it will have to be cut up or mashed. Each day the baby should drink a pint of milk, or less if he eats some cheese or yoghurt; two small portions of meat, fish, poultry, or eggs; some fruit, fresh fruit juice, or vegetables; some cereals—for example, a cereal at breakfast and half a slice of bread at tea; and about 15 g of butter or margarine.

Dr H B Valman, MD, FRCP, is consultant paediatrician, Northwick Park Hospital and Clinical Research Centre, Harrow.

Weighing and measuring

Being undressed and weighed is often the most worrying part of the visit to the child. It is essential that he should be completely undressed, although an allowance can be made for a napkin or pants. Suitable scales which are checked regularly are an essential piece of equipment, and for small infants a pan type with a dial is convenient. The weight in both kg and lbs can be read simultaneously, and the mother will often insist on having the weight in lbs. For older infants and children there is a versatile type of scale which has a pan, a seat, and a platform.

In very young infants it is easier to measure head circumference than length. The measurement should be made, using a paper or fibreglass tapemeasure, around the occipitofrontal circumference (the largest circumference). Measurements can then be plotted on a growth chart together with weights.

The height of infants aged under 2 years is measured on a special measuring board, and accurate results can be obtained only when there are two dedicated measurers present. One has to hold the infant's head against the top board while the other brings up the footboard to the child's feet while stretching him out. When the child is old enough to stand a special stadiometer can be used to measure height, but careful attention to detail is necessary for reproducible results.

The pictures have been taken to show the apparatus rather than how to use it, so as not to distress the child.