problem of supply. Possibly the development centres that have contributed so much will try to meet demands throughout Britain, or the matter may be handed over to industry. As is so frequently the case with new technological developments in the Health Service, no single organisation has clear responsibility for initiating and seeing through the creation of a seat-support for the disabled. A team approach may be the counsel of perfection, but who is to assemble and lead the team?

These difficulties recognised, there is little doubt that the development engineering skills are available to provide adequate seating for the disabled. Whether use is made of these skills will doubtless depend on the determination of clinicians and the allocation of all-too-scarce resources.

¹ Biological Engineering Society. *Seating systems for the disabled. Proceedings of a conference held at the Derbyshire Royal Infirmary, October 27, 1978.* London: Biological Engineering Society, 1979.

² Motloch, WM, quoted by Johnson GR. In: *Seating systems for the disabled. Proceedings of a conference held at the Derbyshire Royal Infirmary, October 27, 1978.* London: Biological Engineering Society, 1979:5.

³ Ewing MR, Garrow C, McHugh N. A sheepskin as a nursing aid. *Lancet* 1961;iii:1447-8.

⁴ Milligan JL. Expanded polyether seat moulds. In: *Seating systems for the disabled. Proceedings of a conference held at the Derbyshire Royal Infirmary, October 27, 1978.* London: Biological Engineering Society, 1979:38.

Doctors in Malta

No end is yet in sight to the dispute between the Maltese government and its doctors.¹ In 1977 laws were introduced requiring all newly qualified doctors to spend two years in Government hospitals and clinics; and the government also assumed control of the island's medical council, so taking away from the medical profession the right to regulate standards. When Maltese doctors protested they were locked out of their hospitals, which have since been staffed by doctors from Libya, Pakistan, and eastern Europe.

This month the Maltese government has extended its powers to cover all premises used for the practice of medicine, so making independent practice there virtually impossible. Throughout the dispute the BMA has supported the Medical Association of Malta and has assisted Maltese medical students in completing their interrupted studies in Britain.

Doctors and other health professionals here can help by refusing invitations to teach or examine in Malta until normal relations are resumed between the island's government and its doctors.

¹ BMA Council. Annual Report 1977-8. *Br Med J* 1978;ii:1065-87.

Regular Review

Do drugs change the course of rheumatoid arthritis?

V WRIGHT, R AMOS

Interpretation of the efficacy of the drug treatment of rheumatoid arthritis is made difficult by the problems of assessing the progress of the disease. Many of the early methods were laudable attempts to introduce order into chaos, and some are still in common use but are not assessments of the rheumatoid process itself; they are either subjective (pain score, early morning stiffness) or semi-subjective (articular index, grip strength) measures of symptoms or of function.

More recent methods of assessment are frequently based on laboratory measurements and are both more objective and reproducible; and they can be shown to alter coincidentally with clinical improvement or deterioration. Methods such as thermography¹ and isotope uptake² measure some of the signs of inflammation, but there is no evidence that these are any more fundamental to understanding the progression of the disease than the earlier, less objective measures. Similarly, measurement of acute-phase proteins or of the erythrocyte sedimentation rate (ESR)^{3,4} reflects secondary phenomena rather than fundamental changes. Nevertheless, they have some important advantages.

Estimation of the titres of rheumatoid factor in the blood is useful in population studies, a poor prognosis generally being associated with high values; but the test is of limited predictive value in individual patients; recognition of IgG rheumatoid

factors not detected by conventional means may improve its clinical value. In general, tests of immunological function are still in their infancy and are frequently difficult to interpret. The serum concentration of histidine is consistently lower than normal in patients with rheumatoid arthritis, but this finding has not received widespread attention in the assessment of the disease.⁵⁻⁷

Probably the best method to detect unequivocal progression of rheumatoid arthritis is radiological assessment. While the presence of erosions does not necessarily indicate poor function, advancing bony destruction must contribute to increasing deformity and further disability. Moreover, the appearance of new erosions is clear evidence of a continuing attack on the joints by the inflammatory process. Nevertheless, radiological assessment is not always straightforward: radiographs can provide only retrospective evidence; with advancing disease further bony damage becomes increasingly difficult to assess; in some circumstances (in certain occupations, for example) erosions may increase in size in the absence of active inflammation, suggesting that mechanical factors can contribute to damage once it has been started by any method. Finally, evidence (or lack of it) of the progression of erosions in some joints does not mean that they are progressing (or not progressing) in all joints.

If, therefore, drug treatment is to alter the outcome of

rheumatoid arthritis it should not only slow or halt radiological damage but also reduce towards normal the objective laboratory indices of active disease. Of all the laboratory tests available, only the erythrocyte sedimentation rate and measurement of C-reactive protein and possibly rheumatoid factor have been correlated with radiological progression.⁸ Measurement of the ESR and acute-phase reactants is easy, and this may distinguish those drugs that might be expected to have an effect on the outcome of rheumatoid arthritis from those that almost certainly do not.

Non-steroidal anti-inflammatory drugs—None of the available non-steroidal drugs alter the outcome of rheumatoid arthritis. Many of the older ones have been shown to be ineffective in reducing the ESR and concentrations of C-reactive protein⁴; and the occasional claims that the newer ones may do so are unsubstantiated. Indeed, relief of pain may make the disease worse. Almost all these drugs produce their symptomatic benefit to a greater or lesser extent by inhibiting prostaglandin synthesis—an action that possibly exacerbates the process of joint destruction.⁹

Gold, D-penicillamine, and chloroquine, in contrast, all have the ability to reduce the ESR and concentrations of the acute-phase proteins.^{10–18} They reduce symptoms effectively despite their lack of intrinsic analgesic activity, but they share the property of a slow onset of action, usually delayed for eight to 12 weeks. When beneficial they are usually far more effective than the non-steroidal anti-inflammatory agents. All three drugs are capable of lowering concentrations of immunoglobulin and of reducing the Rose-Waaler titre.

In some cases there is direct evidence that a reduction may be expected in the progress of bony damage. Early trials¹⁰ showed gold treatment to be symptomatically effective but to have no effect on erosions. More recently, however, serial radiography during five to six years¹⁹ showed a less severe progression of destructive changes in patients treated with a full gold regimen than in those who discontinued the drug early because of toxicity. The earlier that treatment had been started the better the result. Though the design of this study was unusual and the two groups of patients were not entirely comparable, the results confirm an earlier report by Sigler *et al*²⁰ that gold slowed the rate of appearance of new erosions.

D-penicillamine, introduced for the treatment of rheumatoid arthritis because it can dissociate macroglobulins and possibly therefore rheumatoid factors,¹³ shares many of the therapeutic hallmarks of gold. It usually lowers the ESR in parallel with clinical improvement; but, though most measures of rheumatoid disease, whether subjective or objective, improve during treatment with D-penicillamine, so far there is no direct evidence to suggest that it halts erosive damage. Early results in Leeds suggest that D-penicillamine may be unique among the commonly used long-term agents in its potential capability of restoring serum concentrations of histidine to near normality.⁷

Chloroquine has been shown to be symptomatically effective and to lower both the ESR and the titre of rheumatoid factor, but there is no evidence that radiological progression is delayed.

Levamisole, dapsone, and sulphasalazine—With the exception of sulphasalazine (originally introduced for the treatment of rheumatoid arthritis²¹; abandoned, at least in Britain; and recently reintroduced²²), these drugs are recent additions to antirheumatic treatment. Levamisole is undoubtedly clinically effective and lowers the ESR, but is too toxic to be recommended for routine use at present.^{23–25} Dapsone and sulphasalazine improve some patients' symptoms over a period of at

least several weeks while concurrently reducing the ESR and concentrations of acute-phase proteins.²⁶ Results with sulphasalazine are particularly encouraging.

Corticosteroids—Most trials have shown that corticosteroids regularly produce rapid symptomatic improvement and substantially lower the ESR, but that by the end of one year there is little difference between patients treated with aspirin and corticosteroids.^{27–31} This generalisation, however, hides the clinical observation that corticosteroids can halt aggressive or rapidly advancing disease—whereas aspirin cannot. The deterioration that commonly occurs after the initial good response may well reflect reduction of steroids by the physician to avoid toxic effects, whereas if the disease is to be kept under control the dosage needs to be increased.

Early reports from the combined Medical Research Council and Nuffield Foundation studies consistently showed less radiological progression in patients treated with corticosteroids than in those given aspirin, though the difference did not reach statistical significance. After 20 weeks of treatment with gold or prednisolone similar changes are found in the ESR and C-reactive protein.³² But whereas almost every patient on corticosteroids will show this sort of objective improvement those treated with gold may either obtain an even longer-lasting remission or no beneficial effect whatsoever.

One of the early trials of treatment with prednisolone over two years did suggest that the progress of the disease may have been slowed; nevertheless, a rise in the sheep-cell agglutination titre was disquieting,³³ since such a change may be associated with a poorer prognosis. Moreover, the initial dose of prednisone was 20 mg daily—unacceptably high for long-term treatment—and five patients developed major complications. Most clinicians would prefer not to go over 7.5 mg daily, and there is no evidence that this dose inhibits the development of erosions.

Local injections of steroids may also be dangerous. Steroid arthropathy, in which a Charcot-like picture occurs, is more frequent in those patients who have the greatest symptomatic benefit.³⁴ This form of treatment may confer on the joint a degree of function beyond its true capacity; in these circumstances pain is protective.

On balance, there is no strong evidence to suggest that corticosteroids in conventional dosage are regularly capable of altering the course of rheumatoid arthritis.

Cyclophosphamide and azathioprine—Cytotoxic drugs are not widely used, despite reports of improvement in the clinical features of the disease activity (though the ESR has generally not fallen greatly^{35–36}). Isolated claims have been made that the appearance of new joint erosions can be delayed both by cyclophosphamide and by azathioprine.³⁷ The drawback of these drugs is their high general toxicity.

Conclusions—Some drugs in some patients appear to alter profoundly the clinical picture of rheumatoid arthritis and many or all of its laboratory accompaniments, suggesting that they could be influencing the natural history of the disease. Unfortunately, no drug is effective for all patients and, even when a good clinical response has been achieved, relapse may ultimately occur despite continued treatment. Moreover, for some patients treatment may produce only an improvement and not a complete clinical remission. Yet others will abandon treatment because of drug toxicity. All these factors will tend to obscure the very good results achieved for a minority.

Claims that drug treatment does more than temporarily reduce the rate of progression of the disease would be rash on the evidence available, yet most rheumatologists can recall some patients with established active rheumatoid arthritis who

achieved long-lasting remissions while having a particular drug; to dismiss this as coincidence may be to overlook important evidence. We are not yet able to predict which patients will respond to which drugs and who will suffer toxic reactions, though this position may be changing with more understanding of the histocompatibility system.³⁸ If the rate of dropout from trials due to drug toxicity could be reduced we could more easily assess the results.

For the future, one priority should be to identify the characteristics of patients who respond to each drug and to compare responders and non-responders. Another approach may be comparison of correlation matrices of clinical and

biochemical variables to decide whether drugs have a long-term effect.¹⁸ With careful radiological assessment over a long period (a few months is inadequate) we should be able to give a definitive answer to the vexed question of whether drug treatment can alter the outcome of rheumatoid arthritis.

V WRIGHT

Professor of rheumatology,
University of Leeds

R AMOS

Senior registrar in rheumatology,
General Infirmary at Leeds

- ¹ Bacon PA, Collins AJ, Ring FJ, Cosh JA. Thermography in the assessment of inflammatory arthritis. In: MIV Jayson, ed. *Clinics in rheumatic diseases. Diagnosis and management*. London: WB Saunders, 1976;21: 51-66.
- ² Dick WC, Grennan DM. Radioisotopes in the study of normal and inflamed joints. In: MIV Jayson, ed. *Clinics in rheumatic diseases. Diagnosis and management*. London: WB Saunders, 1976;21: 67-76.
- ³ McConkey B, Crockson RA, Crockson AP. The assessment of rheumatoid arthritis. A study based on measurements of serum acute-phase reactants. *Quart J Med* 1972;41:115-25.
- ⁴ McConkey B, Crockson RA, Crockson AP, Wilkinson AR. The effect of some anti-inflammatory drugs on the acute-phase proteins in rheumatoid arthritis. *Quart J Med* 1973;42:785-91.
- ⁵ Gerber DA. Low free serum histidine concentration in rheumatoid arthritis a measure of disease activity. *J Clin Invest* 1975;55:1164-73.
- ⁶ Gerber DA, Gerber HG. Specificity of a low free serum histidine concentration for rheumatoid arthritis. *J Chron Dis* 1977;30:115-27.
- ⁷ Pickup ME, Dixon JS, Lowe JR, Wright V. Serum histidine in rheumatoid arthritis: changes induced by anti-rheumatic drug therapy. *J Rheumatol* (in press).
- ⁸ Amos RS, Constable TJ, Crockson RA, Crockson AP, McConkey B. Rheumatoid arthritis: relation of serum C-reactive protein and erythrocyte sedimentation rates to radiographic changes. *Br Med J* 1977;ii: 195-97.
- ⁹ Morley J. Prostaglandins and lymphokines in inflammation. *Rheum Rehab* 1978; suppl:18-24.
- ¹⁰ Research Sub-committee of the Empire Rheumatism Council. Gold therapy in rheumatoid arthritis. Report of a multi-centre controlled trial. *Ann Rheum Dis* 1960;19:95-119.
- ¹¹ Popert AJ, Meijers KAE, Sharp J, Bier F. Chloroquine diphosphate in rheumatoid arthritis: A controlled trial. *Ann Rheum Dis* 1961;20:18-35.
- ¹² Hamilton EBD, Scott JT. Hydroxychloroquine sulfate (Plaquenil) in treatment of rheumatoid arthritis. *Arthr and Rheum* 1962;5:502-12.
- ¹³ Jaffe IA. The treatment of rheumatoid arthritis and necrotizing vasculitis with penicillamine. *Arthr and Rheum* 1970;13:436-43.
- ¹⁴ Multicentre Trial Group. Controlled trial of D(-)penicillamine in severe rheumatoid arthritis. *Lancet* 1973;ii:275-80.
- ¹⁵ Co-operating Clinics Committee of the American Rheumatism Association. A controlled trial of gold salt therapy in rheumatoid arthritis. *Arthr and Rheum* 1973;16:353-8.
- ¹⁶ Day AT, Golding JR, Lee PN, Butterworth AD. Penicillamine in rheumatoid disease: A long-term study. *Br Med J* 1974;ii:180-3.
- ¹⁷ Dixon A St J, Davies J, Dormandy TL, et al. Synthetic D(-)penicillamine in rheumatoid arthritis. Double-blind controlled study of a high and low dosage regimen. *Ann Rheum Dis* 1975;34:416-21.
- ¹⁸ Dixon JS, Pickup ME, Lowe JR, Hallett C, Lee MR, Wright V. Discriminatory indices of response of patients with rheumatoid arthritis treated with D-penicillamine. *Ann Rheum Dis* in press.
- ¹⁹ Luukkainen R, Kajander A, Isomaki H. Effect of gold on progression of erosions in rheumatoid arthritis: Better results with early treatment. *Scand J Rheum* 1977;6:189-92.
- ²⁰ Sigler JW, Bluhm GB, Duncan H, Sharp JT, Ensign DC, McCrum WR. Gold salts in the treatment of rheumatoid arthritis. A double-blind study. *Ann Int Med* 1974;80:21-6.
- ²¹ Svartz N. A. Therapeutic results in rheumatic polyarthritis. B. Therapeutic results in ulcerative colitis. C. Toxic manifestations in treatment with sulfanilamide preparations. *Acta Med Scand* 1942;110:577-98.
- ²² McConkey B, Amos RS, Butler EP, Crockson RA, Crockson AP, Walsh L. Salazopyrin in rheumatoid arthritis. *Agents and Actions* 1978;8:438-41.
- ²³ Runge LA, Pinals RS, Lourie SH, Tomar RH. Treatment of rheumatoid arthritis with levamisole: a controlled trial. *Arthr and Rheum* 1977;20: 1445-48.
- ²⁴ Multicentre Study Group. Levamisole in rheumatoid arthritis. A randomised double-blind study comparing two dosage regimens of levamisole with placebo. *Lancet* 1978;ii:1007-12.
- ²⁵ Anon. Levamisole—a cautionary note. *Lancet* 1979;ii:291-2.
- ²⁶ McConkey B, Davies P, Crockson RA, Crockson AP, Butler M, Constable TJ. Dapsone in rheumatoid arthritis. *Rheum and Rehab* 1976;15:230-4.
- ²⁷ Joint Committee of the Medical Research Council and Nuffield Foundation on clinical trials of cortisone, ACTH, and other therapeutic measures in chronic rheumatic diseases. A comparison of cortisone and aspirin in the treatment of early cases of rheumatoid arthritis. *Br Med J* 1954;ii:1223.
- ²⁸ Joint Committee of the Medical Research Council and Nuffield Foundation on clinical trials of cortisone, ACTH, and other therapeutic measures in chronic rheumatic diseases. A comparison of cortisone and aspirin in the treatment of early cases of rheumatoid arthritis. *Br Med J* 1955;ii:695.
- ²⁹ Joint Committee of the Medical Research Council and Nuffield Foundation on clinical trials of cortisone, ACTH, and other therapeutic measures in chronic rheumatic diseases. Long-term results in early cases of rheumatoid arthritis treated with either cortisone or aspirin. *Br Med J* 1957;ii:847.
- ³⁰ Empire Rheumatism Council. Multi-centre controlled trial comparing cortisone acetate and acetyl salicylic acid in the long-term treatment of rheumatoid arthritis. Results up to one year. *Ann Rheum Dis* 1955;14: 353-70.
- ³¹ Empire Rheumatism Council. Multi-centre controlled trial comparing cortisone acetate and acetyl salicylic acid in the long-term treatment of rheumatoid arthritis. Results of three years treatment. *Ann Rheum Dis* 1957;16:277-89.
- ³² McConkey B, Davies P, Crockson RA et al. Effects of gold, dapsone and prednisone on serum C-reactive protein and haptoglobin and the erythrocyte sedimentation rate in rheumatoid arthritis. *Ann Rheum Dis* 1979;38:141-4.
- ³³ Joint Committee of the Medical Research Council and the Nuffield Foundation on clinical trials of cortisone, ACTH, and other therapeutic measures in chronic rheumatic diseases. A comparison of prednisolone with aspirin or other analgesics in the treatment of rheumatoid arthritis. *Ann Rheum Dis* 1959;18:173-86.
- ³⁴ Chandler GN, Wright V. Deleterious effect of intra-articular hydrocortisone. *Lancet* 1958;ii:661-3.
- ³⁵ Mason M, Currie HLF, Barnes CG, Dunne JF, Hazelman BL, Strickland ID. Azathioprine in rheumatoid arthritis. *Br Med J* 1969;ii:420-22.
- ³⁶ Urowitz MB, Hunter T, Bookman AAM. Azathioprine in rheumatoid arthritis: Double-blind study comparing full dose to half dose. *J Rheum* 1974;1:274-81.
- ³⁷ Currie HLF, Harris J, Mason RM et al. Comparison of azathioprine, cyclophosphamide, and gold in treatment of rheumatoid arthritis. *Br Med J* 1974;iii:763-6.
- ³⁸ Panayi GS, Wooley P, Batchelor JR. Genetic basis of rheumatoid disease: HLA antigens, disease manifestations, and toxic reactions to drugs. *Br Med J* 1978;ii:1326-28.