CORRESPONDENCE

Bladder cancer as a prescribed industrial disease C A Veys, MD; D Leon, BA.....	Vitamin D supplements in Asian women J G Bissenden, MRCP; O G Brooke, MD, and I R F Brown, MRCPATH.....	Mebendazole and hydatid disease A L Macnair, MB.....
1053	1054	1055
Systematic review of the benzodiazepines E S Snell, FRCP.....	Polypharmacy in rheumatic diseases R C Hamdy, MRCP, and others.....	Lumbar puncture C E Blogg, FFARCS; C G Clough, MRCP, and J M S Pearce, FRCP; J K Hasler, FFARCS..
1053	1054	1055
Kidney biopsy J R T Gabriel, FRCP; M C Bateson, MD; R G Hughes, FRCS.....	Trimethoprim resistance in Finland H E Busk, MD, and B Korner, MD.....	Awareness in general anaesthesia R H Whitaker, FRCS; B Williams, FRCOG; I F Russell, FFARCS; A R Ryan, FFARCS; J B Skinner, FFARCS.....
1054	1054	1056
	Examination of abortuses A J Barson, MD.....	
	1055	

Bladder cancer as a prescribed industrial disease

SIR,—The paper by Sheena M Somerville and others (23 February, p 540) is a timely reminder to clinicians to probe more often into the retrospective life-long occupational history of their patients who present with a neoplasm of the urinary tract. In reply to their question in their follow-up letter (22 March, p 867) asking how the present state of affairs might be improved, I can offer one suggestion—albeit emanating from the other end of the spectrum.

Stoke-on-Trent has had since 1965 a coroner's notification scheme for all patients certified as dying from bladder cancer, whether this was directly related to the cause of death or not. This has been fully described elsewhere.¹ One advantage of the scheme is that it increases doctors' awareness of the relationship between disease and the working environment. Since 1965 I have attended many inquests in order to establish whether this relationship was, in fact, causal; and in 17 instances it was. One rodent controller was exposed to the rodenticide antu containing residual beta-naphthylamine; three firelighter makers to carcinogenic aromatic amines emanating from the hot creosote oil over which they leaned, breathing fumes as they dipped the wooden faggots; and 13 ex-rubber workers were exposed to similar aromatic amines, which contaminated certain antioxidants used in rubber processing up to the end of 1949, when they were promptly withdrawn. All 17 cases were accorded the right to apply for industrial death benefit, and were encouraged to do so.

The Stoke-on-Trent firelighter makers may feel aggrieved at being left out of Somerville and Davies's otherwise comprehensive list; but as doctors we might feel equally concerned at our failure to recognise and help those who gave up their health and even life in the normal pursuance of their job.

CHARLES VEYS

Industrial and Community Health Research Unit,
North Staffs Medical Institute,
Stoke-on-Trent ST4 7NY

¹ Veys, CA. *Br J Ind Med* 1974;31:65-71.

SIR,—We support the view expressed by Sheena Somerville and her colleagues in their letter (22 March, p 876) that there is serious underclaiming of benefits for prescribed

disease No 39—papilloma of the bladder. St Peter's Hospitals have run a claims service for patients with bladder cancer since 1977. We too have found that about one in 20 patients interviewed is eligible for benefit. This is well in excess of the proportion that would be expected on the basis of the national experience.

In Britain over the period 1967-8 to 1971-2 there was a total of 127 new awards of injury or disablement benefit made in cases of occupational bladder cancer.¹ Virtually all the awards were to men. In the corresponding five-year period 1967-71 an average of 5500 new cases of bladder cancer in men was registered annually in Britain. One would therefore expect an average of less than one man in 200 with bladder cancer to be eligible for benefit. The considerable difference between this proportion derived, albeit rather crudely, from national aggregate figures and the proportion of St Peter's and Royal

Marsden cases found eligible for benefit is an indication of the serious extent of underclaiming. Some of this difference may be due to the cases seen at our two hospitals not being completely representative of cases elsewhere. It is unlikely, however, that this could account for much of the discrepancy.

To improve this situation the Institute of Urology, in conjunction with others, is developing a questionnaire to assist clinicians in the crucial step of initially identifying those people most likely to be eligible for benefit. To complement this screening process, carried out at a local level, there is also a need to establish further specialist services to deal with and give expert advice on individual claims. These would perhaps best be organised on a regional basis.

DAVID LEON

Community and Environmental Studies Unit,
St Peter's Hospital, London WC2

¹ Hansard, House of Commons, 31 January 1978, col III.

Systematic review of the benzodiazepines

SIR,—In the wake of alarmist and exaggerated statements on the benzodiazepines in the media and in Parliament a publication of recommendations by the Committee on the Review of Medicines (29 March, p 910) should be welcomed by doctors seeking expert and responsible guidance. However, we fear that in several respects the official response is unsatisfactory and we wish to draw attention to these in your columns.

The format of the statement in the style of data sheets will be confusing. Statutory data sheets already exist on these products and are contained in the Association of the British Pharmaceutical Industry *Compendium*. These are the official data sheets to guide prescribing by the profession and promotion by the industry. The Medicines Act 1968 requires data sheets to be consistent with product licences but the latter cannot be changed other than through the licensing authority, which acts on recommendations from, among others, the Committee on the Review of Medicines. Manufacturers have the right of appeal against decisions of the licensing authority and

this association deplores the publication of recommendations by the Committee on the Review of Medicines before the manufacturers have had an opportunity to exercise this right. It would be much more reasonable if any issues between manufacturers and the licensing authority could be settled and any amendments to licences and data sheets completed before statements from either party are published.

It would have been less confusing to have published the recommendations in general terms—as in the commentary by the Committee on the Review of Medicines. By itself this would have met the situation despite some of the details which are open to question. Furthermore the "Guidelines for data sheets" were drawn up without the normal consultations to be expected and were finalised without full examination of the available evidence from manufacturers. Companies have an unrivalled data source on their own products, accumulated both nationally and internationally. Failure to take such information fully into account can easily lead to imperfect decisions.

One example which casts doubt on the scientific validity of the guidelines is the comment "Benzodiazepines are not suitable for use in combination products." This extreme and unjustifiable pronouncement effectively abolishes all future research in this field (and who knows what clinical benefits may await present or future benzodiazepines coupled with some other known or unknown compound?), and in respect of current products it casts a judgment without allowing debate.

ERIC S SNELL

Association of the British Pharmaceutical Industry,
London W1R 6DD

Kidney biopsy

SIR,—The very clear description of the technique of renal biopsy (23 February, p 547) worries me because the authors imply that the procedure could be used in many hospitals. There are two reasons why this should not be so.

Firstly, to maintain competence in the procedure one needs to perform at least two biopsies each month. General medical wards are unlikely to provide sufficient patients for a single member of a firm to achieve or maintain expertise in kidney biopsy. Secondly, it is unfair to ask a pathologist to make a reasonable diagnosis unless he is reporting on at least 30 renal biopsies annually. In addition, light microscopy alone even in the best hands is inadequate for diagnosis. Immunofluorescent or immunoperoxidase studies and electron microscopy are essential for full morphological analysis. These techniques are only possible with the help of highly trained and skilled technical staff. Kidney biopsy is therefore a team procedure and every hospital cannot have such teams.

ROGER GABRIEL

Department of Renal Medicine,
St Mary's Hospital,
London W2 1NY

SIR,—It is a great pity that recent contributions on needle biopsy of the kidney (23 February, p 547) and liver (15 March, p 776) should recommend a procedure at variance with that proposed by the manufacturers of the Tru-Cut needle. The instructions supplied with the needles were amended to improve safety, and operators should consult and comply with them.

The disposable Tru-Cut needle was introduced and became popular without formal comparison with the reusable Menghini needle. It is very much more expensive, and certainly for liver biopsy confers no advantage.¹

MALCOLM C BATESON

University Department of Medicine,
Ninewells Hospital and Medical School,
Dundee DD1 9SY

¹ Bateson MC, Hopwood D, Duguid HLD, Bouchier IAD. *J Clin Path* 1980;33:131-3.

SIR,—I noted your correction (8 March, p 700) to the table of contraindications to kidney biopsy (23 February, p 547); however, I feel that some of the haematological data still require comment.

A prothrombin time of greater than or equal to 16 seconds may be a contraindication, although a safer limit would be "not greater than 2-3 seconds longer than the time for

control plasma" (which is from a pool of normal plasma samples and which may give times as short as 11 seconds, depending on the thromboplastin used in the test). Recording the result of the test plasma as a ratio of the control value is a recommended method.¹

The expression of the platelet count $\times 10^9/l$ [added in the subediting—Ed, *BMJ*] is unfortunate when the conventional factor is $\times 10^9/l$,^{2,3} the contraindication then becoming a platelet count under $100 \times 10^9/l$. One useful simple screening test in renal disease is the bleeding time,⁴ which helps to detect abnormal platelet function (the major factor contributing to the haemostatic defect of uraemia). Abnormalities of platelet retention (in glass bead filters) and platelet factor 3 availability (decrease often associated with abnormal platelet aggregation) have been found in a variable percentage of uraemic patients by most investigators.⁵ In general, however, these defects are observed in patients with clinically noteworthy bleeding and usually disappear after peritoneal dialysis or haemodialysis.⁶

R G HUGHES

Royal Marsden Hospital,
London SW3 6JJ

¹ Poller L. *Association of Clinical Pathologists Broad-sheet* 1970;No 71.

² Baron DN, Broughton PMG, Cohen M, Lansley TS, Lewis SM, Swinton NK. *J Clin Pathol* 1974;27:590-7.

³ ICSM, IFCC, WAPS. *Br J Haematol* 1972;23:787-8.

⁴ Ivy AC, Nelson D, Bucher G. *J Lab Clin Med* 1940;26:1812-22.

⁵ Weiss HJ. In: Williams WJ, Beutler E, Ersler AJ, Rundles RW, eds. *Haematology*. New York: McGraw Hill Book Co, 1972:1171-5.

⁶ Rabiner SF. *Prog Hemostasis Thromb* 1972;1:233-50.

Vitamin D supplements in Asian women

SIR,—Dr O G Brooke and others (15 March, p 751) report a better calcium status in babies born to Asian women whose diets were supplemented by calciferol 1000 IU/day during pregnancy. The recommendation is therefore to give calciferol to pregnant Asian women. Before such a recommendation is adopted on a wide scale further information must be provided. If allocation to the trial were on a random basis, why at the end of the trial were there 67 women in the control group and 59 in the treatment group? Patients were excluded from the trial for reasons such as pre-term deliveries and congenital malformation. Was there a higher incidence in these complications in babies born to mothers receiving the calciferol supplement, and if so can an explanation be provided?

J G BISSENDEN

East Birmingham Hospital,
Birmingham B9 5ST

* * * We sent a copy of this letter to the author, whose reply is printed below.—Ed, *BMJ*.

SIR,—We thank Dr Bissenden for his interest and respond to his questions as follows:

(1) In all, 135 women entered the trial. They were allocated at random by the hospital pharmacy on the basis of their hospital numbers, and there were initially 63 in the treatment group and 72 in the control group. Simple statistical analysis shows that there is a 5% chance that the size of one of the groups could be less than 56 or greater than 79. Our group sizes were thus well within these limits.

(2) Three women in the control group and two in the treatment group withdrew, either

because they moved to other areas before delivery or because they decided not to continue with the trial. Other exclusions were as follows: in the control group there was one preterm delivery and one neonatal death due to intrapartum anoxia and meconium aspiration; in the treatment group one woman was excluded because she developed gestational diabetes; and one infant had spina bifida. There is thus no evidence of a higher incidence of complications in the treatment group.

O G BROOKE
I R F BROWN

St George's Hospital Medical School,
London SW17 0RE

Polypharmacy in rheumatic diseases

SIR,—We were pleased to read in your leading article (8 March, p 666) "Polypharmacy in rheumatic diseases" that the plasma half lives of drugs tend to be longer in older patients. We wish more pharmaceutical companies took notice of this statement, and would like to emphasise the need to establish the pharmacokinetic profile of various drugs in the older population before launching the sales of these drugs. It is well established that the older population is the major consumer of drugs,^{1,2} in particular the antirheumatic agents, and yet when one screens the literature one is appalled by the paucity of information regarding the biological handling of these drugs by elderly patients. In fact, most of these parameters seem to be measured only in young healthy volunteers. Is it reasonable to extrapolate the findings to the older population, when it is well known that as the individual ages the intestinal mucosal surface area decreases,³ the blood supply to the gastrointestinal tract decreases,⁴ the hepatic and renal clearances become impaired,⁴ the binding of drugs to plasma protein seems altered,^{5,6} and the number of receptor sites on the target organs may well also decrease?⁷

We feel that clinicians should insist on knowing the pharmacokinetic profile of the various drugs they prescribe to their elderly patients and we look forward to the time when this information will become an integral part of the work done on any drug before it is marketed.

R HAMDY
N H PERERA
M WHITWORTH

Department of Clinical Gerontology,
St John's Hospital,
London SW11 1SP

¹ Crooks J, Shepherd A, Stevenson I. *Health Bulletin* 1975;33:222-7.

² Williamson J. *Practitioner* 1978;220:749-55.

³ Warren PM. *Lancet* 1978;ii:849-50.

⁴ Richey DP. In: Goldman R, Rockstein M, Sussman M, eds. *The physiology and pathology of human ageing*. New York: Academic Press, 1975.

⁵ O'Malley K, Judge T, Crooks J. In: Avery GS ed. *Drug treatment—principles and practice of clinical pharmacology and therapeutics*. Sydney: Adis Press, 1976:123-42.

⁶ Stevenson IH, Shepherd AM. *Br J Clin Pract* 1978; suppl 2: 67-71.

⁷ Roth GS. *Fed Proc* 1979;38:1910-4.

Trimethoprim resistance in Finland

SIR,—Mr P Huovinen and Professor P Toivanen reported (12 January, p 72) that in 1978, after five years' use of plain trimethoprim in the Turku area, the incidence of trimethoprim-resistant *Escherichia coli* isolated from urines was 11% and 23% in outpatients and inpatients respectively. By contrast, the inci-