function in putative immune complex disease.^{12, 13} Some of our patients were found by Lockwood *et al*¹² to have defective splenic function during their initial disease, which was reversed by a plasma exchange regimen. The effect of infection on splenic reticuloendothelial function and its relation to the reappearance of circulating immune complexes in infection-provoked relapse clearly warrant further study. A role for infection in the pathogenesis of Wegener's granulomatosis is further suggested by the observation that 10 of the 18 patients studied had a history of chronic suppurative infection of the respiratory tract, which long predated the vasculitic illness (A J Pinching *et al*); four of the five patients with infection-provoked relapse had such a history.

Another possible mechanism for infection-provoked relapse is the enhancement of quiescent disease by the acute-phase or cellular response to infection: a limited amount of immunopathogenetic initiator (for example, circulating immune complex) may cause more tissue injury as a result of the provision of an increased supply of inflammatory mediators.

The phenomenon of infection-provoked enhancement of allergic tissue injury is seen in conditions of widely differing pathogenesis including autoantibody-mediated diseases such as antiglomerular basement-membrane disease¹ and myasthenia gravis, immune complex disease such as systemic lupus erythematosus,² renal graft rejection,³ and in this study Wegener's granulomatosis. Infection may also exacerbate disease in asthma and infantile eczema. Whether infection exacerbates these varying allergic diseases by a common mechanism is not known. Clearly further studies of this clinically and theoretically important phenomenon are required.

We are grateful to the clinical virology department for carrying out the virological procedures.

References

- 1 Rees AJ, Lockwood CM, Peters DK. Enhanced allergic tissue injury in Goodpasture's syndrome by intercurrent bacterial infection. *Br Med J* 1977;iii:723-6.
- 2 Dubois EL. *Lupus erythematosus*. 2nd ed. Los Angeles: University of South California Press, 1974.
- 3 Simmons RL, Weil R, Tallent MB, Kjellstrand CM, Najarian JS. Do mild infections trigger the rejection of renal allografts? *Transplant Proc* 1970;2:419-23.
- 4 David DS, Millian SJ, Whitsell JC, *et al*. Viral syndromes and renal homograft rejection. *Ann Surg* 1972;175:257-9.
- 5 Briggs JD, Timbury MC, Paton AM, Bell PRF. Viral infections and renal transplant rejection. *Br Med J* 1972;iv:520-2.
- 6 Lopes C, Simmons RL, Mauer M, Park B, Najarian JS, Good RA. Role of virus infections in immunosuppressed renal transplant patients. *Transplant Proc* 1973;5:803-8.
- 7 Dossetor JB, Schlant J, Olson LA, Zabriskie JB. Streptococcal infection as a cause of hyperacute renal allograft rejection. *Transplant Proc* 1978;10:483-7.
- 8 Byrd LH, Tapin L, Cheigh JS, Aronian J, Stenzel KH, Rubin AL. Association between *Streptococcus faecalis* urinary infections and graft rejection in kidney transplantation. *Lancet* 1978;ii:1167-9.
- 9 Fussell BA, Lockwood CM, Scott DM, Pinching AJ, Peters DK. Value of immune complex assays in diagnosis and management. *Lancet* 1978;ii:359-63.
- 10 Ewan PW, Jones HA, Rhodes CG, Hughes JMB. Detection of intrapulmonary haemorrhage by carbon monoxide uptake: a new method and its application in Goodpasture's syndrome. *N Engl J Med* 1976;295:1391-6.
- 11 Fauci AS, Haynes BK, Katz P. The spectrum of vasculitis. *Ann Int Med* 1978;89:660-76.
- 12 Lockwood CM, Worrledge S, Nicholas A, Cotton C, Peters DK. Reversal of impaired splenic function in patients with nephritis or vasculitis (or both) by plasma exchange. *N Engl J Med* 1979;300:524-30.
- 13 Frank MM, Hamburger MJ, Lawley TJ, Kimberly RD, Plotz PH. Defective reticulo-endothelial system Fc-receptor function in systemic lupus erythematosus. *N Engl J Med* 1979;300:518-23.

(Accepted 8 August 1980)

SHORT REPORTS

Recurrent abdominal pain associated with digoxin in a patient undergoing maintenance haemodialysis

In addition to its inotropic effect on the heart digitalis is a direct peripheral vasoconstrictor.¹ In particular, cardiac glycosides cause splanchnic vasoconstriction and reduced mesenteric blood flow in dogs,² monkeys,³ and variably in patients with congestive heart failure.⁴ There is a well-recognised association of administration of digitalis with intestinal infarction,⁵ but digitalis has not been associated before with chronic intestinal ischaemia or abdominal angina.

Case report

A 55-year-old obese white man was admitted to hospital because of abdominal pain. He had been maintained on thrice weekly haemodialysis for two and a half years because of renal failure caused by nephrosclerosis. He had a history of duodenal ulcer, hiatus hernia, myocardial infarction, and angina pectoris. His medications included digoxin 0.25 mg every other day and nitroglycerin 0.4 mg sublingually as needed.

Soon after starting haemodialysis he began to have severe recurrent, sustained, central, non-radiating abdominal pain brought on by physical exertion and relieved after 10-20 minutes by lying supine. The pain often occurred after dialysis, especially when he had required fluid removal in excess of two litres and had become hypotensive. The pain was never related to meals; and he had had no weight loss, diarrhoea, melaena, or rectal bleeding. Radiographs of the upper gastrointestinal tract and gall bladder were normal.

He was pale, sweating, and in great distress. The blood pressure was 100/60 mm Hg, and results of cardiac and respiratory examinations were normal. There was mild tenderness in the hypogastrium with no rebound

tenderness, and the bowel sounds were active. Results of Hemocult test of the stool were negative. His haemoglobin concentration was 13.3 g/dl, packed cell volume 0.4 (40%), and peripheral blood leucocyte count $6.1 \times 10^9/l$ ($6100/mm^3$). There was no biochemical or electrocardiographic evidence of new myocardial injury. A plain film of the abdomen showed a non-specific gas pattern and heavily calcified abdominal aorta and iliac vessels.

We suspected that the pain was due to mesenteric vascular disease, and abdominal angiography was planned after a trial of withholding treatment with digoxin. This lasted for the next five months, and though the dialysis regimen was unaltered the patient had no further abdominal pain. To test whether digoxin was the provocative agent the patient was rechallenged with digoxin, and recurrence of abdominal pain after dialysis was noted after 1.0 mg. Digoxin was again withdrawn, and he has since remained free of pain.

Comment

Intestinal angina is a syndrome of postprandial abdominal pain and weight loss associated with minimal physical findings. We believe that this case represents a variation of this syndrome and that the abdominal pain was a result of intestinal ischaemia, since it was induced by factors that either reduce cardiac output or cause splanchnic vasoconstriction—namely, contraction of the intravascular volume by rapid fluid removal during ultrafiltration-haemodialysis and physical exertion. Since digitalis acts as a direct splanchnic vasoconstrictor,^{2,3} relief of this patient's pain after discontinuing digoxin and the recurrence when rechallenged support the belief that the pain represented intestinal ischaemia that was exacerbated by digoxin.

A syndrome of digoxin-induced abdominal pain has not been reported in patients undergoing dialysis, many of whom take digitalis, have calcified aortas, and often undergo the stress of rapid fluid removal. To explain why this patient experienced pain and apparently similar patients do not is difficult. An additional predisposing factor for this patient was his packed cell volume, which was unusually high

for a patient with renal failure. Lowered blood viscosity, which usually accompanies the anaemia of chronic renal failure, may prevent sluggish mesenteric blood flow in other patients undergoing dialysis who are subjected to similar haemodynamic stresses.

Digitalis should be considered as a possible cause when evaluating obscure abdominal pain in patients undergoing haemodialysis, who may benefit—and be rendered free of pain—when digitalis is withheld.

- ¹ Ross J, Waldhausen JA, Braunwald E. Studies on digitalis: direct effects on peripheral vascular resistance. *J Clin Invest* 1960;**39**:930-6.
- ² Harrison LA, Blaschke J, Phillips RS, Price WE, Cotten MD, Jacobson ED. Effects of ouabain on the splanchnic circulation. *J Pharmacol Exp Ther* 1969;**169**:321-7.
- ³ Shanbour LL, Jacobson ED, Brobmann GF, Hinshaw LB. Effects of ouabain on splanchnic hemodynamics in the rhesus monkey. *Am Heart J* 1971;**81**:511-5.
- ⁴ Ferrer MI, Bradley SE, Wheeler HO, Ewson Y, Preisig R, Harvey RM. The effect of digoxin in the splanchnic circulation in ventricular failure. *Circulation* 1965;**32**:524-37.
- ⁵ Gazes PC, Holmes CR, Moseley V, Pratt-Thomas HR. Acute hemorrhage and necrosis of the intestines associated with digitalization. *Circulation* 1961;**23**:358-64.

(Accepted 3 June 1980)

Departments of Medicine, Veterans Administration Medical Centre, and Downstate Medical Centre, Brooklyn, New York 11209, USA

MARTIN FEINROTH, MB, PHD, fellow in nephrology
MARY V FEINROTH, MB, fellow in nephrology
A PETER LUNDIN, MD, assistant professor of medicine
ELI A FRIEDMAN, MD, FACP, professor of medicine
GEOFFREY M BERLYNE, MD, FRCP, professor of medicine

Arthritis preceding fulminant ulcerative colitis and responding to colectomy

Although it has long been recognised that ulcerative colitis may be complicated by bouts of arthritis,^{1,2} the clinical course of "colitic arthritis" has only recently been characterised.³ Large joints are usually affected in a flitting and asymmetrical way, most commonly in chronic ulcerative colitis, and in a series of 269 patients never preceded the onset of bowel symptoms. Colectomy invariably results in prolonged remission of arthritic symptoms.⁴ We describe a patient with colitic arthritis whose joint symptoms preceded bowel symptoms by five years and in whom the onset of colitis was acute and fulminant.

Case report

A 19-year-old woman presented in December 1978 with a four-year history of recurrent pain in her shoulders, elbows, wrists, and knees. Her general health was otherwise good with no bowel symptoms. There was tenderness over the ulnar heads and discomfort on moving the shoulders and knees but no other abnormality. Radiographs were normal. Results of routine haematological studies and tests for antinuclear factor were normal. The presumptive diagnosis was seronegative rheumatoid disease. She was given a six months' course of chloroquine phosphate and naproxen and remained fairly well over the next 15 months with only one exacerbation of soreness and stiffness of the right shoulder, which responded to a local hydrocortisone injection.

In January 1980 she became generally unwell with lethargy, weakness, and an exacerbation of pain and stiffness in her large joints. She developed bowel symptoms for the first time with lower abdominal pain and loose, watery bowel motions up to eight times daily, streaked with mucus and blood. Several stool cultures were sterile, and symptomatic treatment had little effect. On admission to hospital she was pale and dehydrated with a temperature of 38.5°C, though her pulse and blood pressure were stable. There was no obvious swelling or inflammation around her large joints, but they were painful and stiff. She had generalised abdominal tenderness, particularly in the left iliac fossa, and bowel sounds were pronounced. Abdominal radiography showed a grossly dilated colon. Findings of routine haematological and serological studies were again normal. Her HLA type was subsequently reported as A2, B17, BW44, BW4 but not B27. Radiographs of her sacroiliac joints were normal.

Conservative treatment with intravenous fluids and high-dose corticosteroids produced a temporary initial response. Five days after admission,

however, she developed severe abdominal pain with guarding and rebound tenderness, and bowel sounds disappeared. Abdominal radiography confirmed perforation of the bowel. At operation the whole colon was distended, inflamed, and oedematous and there were perforations in the caecum and sigmoid colon. She underwent total colectomy and terminal ileostomy. Since the rectum had looked relatively normal on sigmoidoscopy it was conserved. Histological examination showed extensive ulceration with focal crypt abscess formation, the appearances being typical of severe ulcerative colitis.

The patient made an excellent recovery, and five months later remained entirely free of joint symptoms without anti-inflammatory drugs.

Comment

The pattern of joint disease, exacerbation of symptoms with the onset of colitis, and dramatic response to colectomy confirm that this patient had colitic arthritis.³ The unique features, however, were the long history of joint symptoms preceding the onset of colitis and the acute fulminant mode of onset of colitis. Significant colitis was probably not present during the five years when no bowel symptoms existed, but perhaps the relatively simple investigation of sigmoidoscopy with rectal biopsy should be considered in recurrent undiagnosed seronegative arthropathy. It might then be possible to offer specific treatment for colitis with resultant control of joint symptoms and to hope to prevent severe colitis.

Requests for reprints should be sent to Dr David M Fraser.

- ¹ Bargen JA. Ten years' experience in treatment of chronic ulcerative colitis. *Transactions of the American Gastroenterological Association* 1933;**36**:49-60.
- ² Hench PS, Bauer W, Boland E, et al. Rheumatism and arthritis. Review of American and English literature for 1940. *Ann Intern Med* 1941;**15**:1002-108.
- ³ Wright V, Watkinson G. The arthritis of ulcerative colitis. *Br Med J* 1965;iii:670-5.
- ⁴ Brooke BN. Outcome of surgery for ulcerative colitis. *Lancet* 1956;iii:532-6.

(Accepted 15 July 1980)

Milesmark Hospital, Rumbungwell, Dunfermline KY12 9NR

DAVID K McCULLOCH, MRCP, registrar
DAVID M FRASER, MRCP, consultant physician

Dunfermline and West Fife Hospital, Dunfermline

ALLAN L TURNER, FRCS, consultant surgeon

Successful pregnancy in severe chronic renal failure not requiring dialysis

Chronic renal failure in women is usually associated with impaired fertility¹ and a high incidence of miscarriage.^{2,3} In women with creatinine clearance below 10 ml/min successful pregnancy not requiring dialysis has not been recorded. We report such a case.

Case report

A 31-year-old Asian housewife had three full-term normal deliveries in 1968, 1969, and 1974, during each of which her albuminuria was + + +. During the third pregnancy creatinine clearance was 133 ml/min. She was normotensive during all three pregnancies. In June 1979, however, she presented to her GP with headache, and her blood pressure was 220/120 mm Hg. Oxprenolol 320 mg daily reduced the blood pressure to 160/85 mm Hg by October. In November 1979 (at 17 weeks of pregnancy) her blood pressure was 170/100 mm Hg. Her haemoglobin concentration was 7.6 g/dl, blood urea and creatinine concentrations 27 mmol/l (162.7 mg/100 ml) and 500 µmol/l (5.7 mg/100 ml) respectively, proteinuria 3 g/day, and creatinine clearance 7.6 ml/min. She was admitted to hospital and received a 40 g protein "no added salt" diet, transfused with 2 units of blood, and prescribed only methyldopa 250 mg twice daily. Blood urea and creatinine concentrations rose slowly, though blood pressure and fetal growth were initially satisfactory.

On 12 February 1980 she was transferred to the Manchester Royal