PRACTICE OBSERVED

Emergencies in the Home

Anaphylaxis and drug reactions

R L HERBERT

Type I hypersensitivity reactions result from the patient forming abnormal amounts of IgE antibodies in response to antigen exposure. The antibody attaches itself to the surface of the mast cells, which in response to antigen challenge break down and release histamine, 5-hydroxytryptamine, and slow-reacting substance of anaphylaxis from their granular contents. The reactions (also referred to as atopic or immediate) may vary from an irritating pimple through distressing bronchospasm to the extreme of circulatory collapse and death.

Managing mild hypersensitivity

This short paper is intended to deal with the immediate treatment of anaphylaxis in the home. Before dismissing the nuisance end of the scale, however, I want to make some observations and recommendations. Allergic rhinitis and urticaria often provoke reactions varying from discomfort to annoyance—not only from the public, but also sadly from many doctors. Patients with ugly and irritating rashes that cause them to behave like pariahs, students trying to sit final university examinations, and bus drivers working through floods of blinding tears and nasal fluid merit more than just antihistamines. While many patients respond well to antihistamines with no apparent sedation, my own opinion is that this is often a false state of security. The sedation produced may often be insidious, only being apparent after the exam has been failed or the child killed in a traffic accident. Nasal, ocular, and inhalable preparations of disodium cromoglycate and beclomethasone used either separately or in combination have made life much more tolerable for the atopic patient. A single intramuscular injection of either methylprednisolone

acetate or triamcinolone diacetate is often justified at the onset of the symptoms. The adrenal suppression, which can be proved only by sophisticated biochemistry, although a scientific reality is clinically a myth that may be safely ignored for all practical purposes. For those who still have reservations, however, it has recently been shown that many patients respond well to a single intramuscular injection of 120 units of corticosteroid. Even 5 mg-tablets of prednisolone on a daily reduced dosage (six the first day down to one on the sixth day and nothing on the seventh), which gives almost certain relief of symptoms with no interference in mental or adrenal state, should not in my opinion be withheld when the severity of the symptoms and the circumstances of the patient merit it. Picture, if you will, a bride and groom on the greatest day of their lives in a village church in England sometime in June trying to maintain their dignity while under attack by every pollen grain our countryside can muster.

Nevertheless, the mainstay of treatment must be accurate identification of the offending allergen and either their eradication from the environment or desensitisation. Repeated desensitisation courses or maintenance courses will often be required for a satisfactory result. This should still be the treatment of choice, however, and the previously-discussed treatments should be reserved for those patients who either present too late for desensitisation or fail to respond to it. Recently, the task has been made even easier with the availability of the Phadmac radioallergen sorbent test, which is capable of identifying IgE antibodies to specific allergens in 3.0 ml of blood. But for most of us careful history taking backed up with skin testing will suffice. It is also interesting to note that the same drug company now manufactures bee and wasp venom vaccines. There are five deaths each year in the British Isles from anaphylaxis to wasp and bee stings (not to mention the rest of the world), which is sufficient justification for costly research and production.

Desensitisation is a simple procedure even though time consuming. So long as basic precautionary principles are adhered to—that is, a syringe of 0.5 ml of adrenaline 1 in 1000

Chalford Health Centre, London E4
R L HERBERT, MA, CC, general practitioner

782

BRITISH MEDICAL JOURNAL VOLUME 282 7 MARCH 1981

drawn up ready, and the recommended injection regimen strictly adhered to—it has only minimal danger. If there should be a reaction at one dose then returning to the previous last dose that produced no reaction before advancing further through the regimen will almost always avoid disaster. In the past 15 years my desensitisation of about 50 patients each spring has resulted in only one case of anaphylaxis.

Severe anaphylaxis

While severe anaphylaxis is rare it is nevertheless a real and ever-present risk, which has little regard for fitness or age. Pooch Bear is credited with having said, "Accidents are funny things; you don't know they are going to happen until they do"; so often is the case with anaphylaxis. The availability at all times of the knowledge and materials with which immediately to intervene is mandatory.

In the first instance, the basic principles of the care of a shocked patient with increasing circulatory collapse apply as always. Fumbling about in a medical case or drugs cabinet searching for little-used drugs that when found may be out of date will often guarantee failure. By dialling 999 and passing the responsibility to the ambulance service you invoke a time-consuming series of events that you and your patient cannot afford. Your patient's circulatory system is progressively deteriorating, and soon the peripheral circulation will have shut down totally. Unless an intravenous cannula is inserted before this happens then the expertise of the hospital is at best hindered and at worst completely negated. Establishment of an intravenous infusion with either saline or Hartmann's solution before your patient's veins totally collapse is mandatory. It is equally essential to keep a cautious watch on the patient's cardiac state and airway if he becomes unconscious. If your patient, while you are cannulating his antecubital vein with him lying on his back, dies from anoxia resulting from an inadequately maintained airway, your hospital colleagues and the coroner are unlikely to be impressed. Finally, should the patient's heart stop despite an intravenous infusion, appropriate drugs, and airway maintenance external cardiac massage and mouth-to-mouth respiration must be started and must be continued until inside the hospital. Defibrillation and positive-pressure ventilation by those of us who have the necessary equipment would, of course, increase the chance of success, but this is rarely available.

Fortunately, most cases of anaphylaxis do not progress to such extremes and once the intravenous infusion is flowing the doctor can relax a little and make his choice of drugs.

Drugs for anaphylaxis

If the patient is not unconscious a very useful preparation, often forgotten, is sublingual isoprenaline in tablet form. Its speed of absorption will often give valuable time—the circulation will be maintained just long enough for the intravenous infusion to be set up. An immediate subcutaneous injection of 0.5 ml adrenaline 1 in 1000 if it is already prepared in a syringe and not in a bottle in a case or your cupboard will be equally effective. An 8 mg bolus of chlorpheniramine maleate into the infusion is the treatment of choice (4 mg in my experience is almost homeopathic); sedation such as that there is may serve you well in relating an often terrified patient. In such a potentially serious condition, however, I believe that the use of more powerful preparations than the antihistamines is not progress to such extremes and once the intravenous infusion is flowing the doctor can relax a little and make his choice of drugs.

treatment may also be of benefit. As a final gesture I suggest that the injection of 0.5 ml of adrenaline 1 in 1000 into the site of the original injection (if that was the causative factor) may slow down further absorption of the allergen and restrict the severity of the attack.

Treating in the home can be as effective as in hospital

I was asked to write on the treatment of anaphylaxis in the home. To me the title implies that there is a difference in treatment in the home as opposed to hospital. Although this is often so, I would submit that this is morally wrong and need not be so. Apart from the practical considerations of equipment and the great cost and physical size, there is no valid reason to justify the implied lesser treatment. The basic ideas of medical intervention in life-threatening illness or injury are the same, in my opinion, irrespective of the venue. To deny a patient in extremis medical expertise simply because he is either at home or in the street and enforce transport to hospital is to increase the risk of death.

While I accept that it is difficult to set up a drip singlehandedly, it is not impossible, nor is it impossible to deal with a cardiac arrest singlehandedly. With the advent of trained ambulance crew in intubation and infusion techniques there is now an ever increasing chance of such a team arriving at your patient's 999 call. The training programme, it is hoped, will gain in impetus so that in the near future all 999 calls will be responded to by such a team. In the past, as now, general practitioners have set themselves apart from the ambulance service, especially the men on the ground. We now see, however, a new phase developing where together ambulance crew and general practitioners may function as a team and apply skills previously only thought possible in the hospitals. Our moral and NHS contractual obligations are to give medical aid to those who need, where and when they need. We must go to those who cannot come to us. A whole new era lies before us; we can grasp it and have a primary care team par excellence with general practitioner and ambulance crew equally skilled and working as a powerful unit.

Too many primary care problems, however, are passed on to the secondary care physicians in hospital with neither intervention nor support; very often resulting in their transfer to the ultimate care physician in the mortuary.

My thanks to Mrs Linda Herbert for patiently typing this paper.

This series on emergencies in the home will continue in a few weeks.

Searching for a practice

When you're interviewed for a practice your wife is also interviewed. The standard pattern is that you drive up, you are greeted by the doctor, and then your wife is whisked away to meet his wife and then she meets the other wife. I always remember in 1954 a large practice in the North and being interviewed in a fabulous house. Very impressed, I thought, "This is it, this is what we want." With the partners there we couldn't talk to each other, but when we had tea together my wife was one side of the room and I was on the other, and we'd look at each other, trying to signal. I was trying to say, "I think it's all right," but she was very cold and I didn't know whether she was just fed up with me or what. Anyway, when we got into the car and drove home, she explained that she'd managed to spend half an hour with the junior partner's wife and all the misery and the exploitation in the partnership had come out. This girl had just poured her heart out to my wife. We thought no more about that job.—KEN DICKINSON, general practitioner, Birmingham

We will be pleased to consider similar interesting stories about general practice.—Ed, BMJ

BRITISH MEDICAL JOURNAL VOLUME 282 7 MARCH 1981

783

Medical Records

Practical problems of improving records

DAVID STUART

"What is written without effort is in general read without pleasure." Samuel Johnson, *Johnson's Miscellany*, Vol II, p 309.

The average family doctor reading articles about improving records may well be forgiven for having a sigh of despair, tinged with envy of those few who, with superhuman energy, have pressed on and completed some ambitious scheme of recording, and who then insist on pricking his conscience by writing about it. For his own part, he is inheriting some years ago over 2000 envelopes, some unopened by a medical pen, some held together by brown rotting Sellotape and containing a miscellany of cards he cannot read, and a bunch of letters, a few of which contain vital information about his patient, but most recording normal blood tests or unimportant casualty attendances of yesterday. His partners regard any scheme for improving things with the usual mixture of enthusiasm and suspicion of anything that might require more work, and his secretary and receptionist already have enough on their plate, thank you. . . .

What to do first

There are several prerequisites for any systematic approach to the problem.

(1) Start off by defining the objective of an assault on the records. Who is going to benefit and in what way? Is the objective to improve the usefulness of the notes in everyday practice, or is it related to some other end—for example, audit, medico-legal considerations, or even making good recording a passing preoccupation? Do not expect an exercise without knowing what benefit you expect, or else you may labour mightily and achieve nothing. For instance, one sees age-old registers constructed with painstaking effort but then, because the partnership concerned is not conducting research or audit and the county council computer is already recalling patients for cancer and immunisation.

(2) Everyone in the practice must know about, and agree to, any action that is proposed—this means ancillary staff as well as the partners.

(3) The time spent on a project needs to be identified and organised. Who is going to do a job? Will time be spent regularly on it, or will it be done in spare moments? If the latter, has a target of achievement been set?

(4) Be prepared to use unskilled or semi-skilled labour for the routine work—the doctor himself will have enough skilled sorting out to do anyway.

How to proceed

Once a partnership has actively decided to effect some improvement in records, a number of objectives present themselves. What might

Great Yarmouth, Norfolk
DAVID STUART, MA, MRCGP, general practitioner

be regarded as essential or merely desirable must be a matter for individual choice, and the list that follows is my own, in my personal order of importance.

(A) *Get continuous cards and letters put into date order*—a basic task requiring no special skill. If existing ancillary staff cannot cope with the task in reasonable time, extra staff may be employed temporarily. At least one practice has used the Youth Opportunities Employment Scheme for this purpose. The partnership must decide: (a) *What order to have the cards in.* Will the most recent face forwards or backwards? Will the letters be behind the continuation cards or vice versa?

(B) *How the continuation sheets should be attached to each other.* We push a hole in the top left corner and use treasury tags. Some prefer stapling the left edge. If left unattached, the sheets tend gradually to fall into chaos again.

(C) *Wash out and destroy unnecessary continuation cards and letters.* This may be done by skilled ancillary staff or a nurse, though some doctors think that it should be done only by them. One can attach the task alphabetically, but the alternative is to select first the records that need weeding out most (usually the "fat files"). In this case one needs to get into the habit of picking out one or more daily, and the envelopes that have been dealt with need to be identified by a mark (for example, a coloured sticker) unless another stick is being done at the same time that will identify the sorted envelope (for example, filling in a summary card).

(D) *Adopt colour coding for important diseases or facts.* The colours originally adopted by the Royal College of General Practitioners are now widely used, and a suitable sticker on the front of the envelope saves one from ignoring a hypersensitivity or that the patient is a known diabetic. Coloured stickers take hardly any time to put on and are open to some refinement of the doctor's own classification. I have diagonally halved stickers to indicate either a family history of the disease concerned or a predilection potential such as an isolated high blood pressure reading or mildly abnormal glucose tolerance. If a disease register is to be kept, the sticker may be ticked to indicate that the patient's name has been entered.

The RCGP list is: **RED**—hypertension; **BLUE**—hypertension; **BROWN**—diabetes; **YELLOW**—epilepsy; **GREEN**—tuberculosis; **BLACK**—attested suicide; **WHITE**—long-term maintenance treatment.

(E) *Look at the layout of the notes.* Minor changes in layout can improve legibility greatly, for example: (a) Is the writing legible? If not, can it be improved by deliberate self-training? Competent, if not beautiful, scripts are not difficult to learn.¹ If a doctor's handwriting is hopelessly illegible, having the records typed is the only remedy, though only time-wasting.

(F) *Box in diagnoses using colours to distinguish diagnosis from treatment registers.*

(G) *Leave one-line gaps between consultations.*

(H) *Have a system for transcribing out-of-hours notes to the main record.* Some practices have the notes typed in, though we prefer to use the special addressed paper supplied by Winthrop Laboratories.

(I) *Have some system for identifying current treatment.* Nothing is more irritating to one's partner or locum than to be confronted by treatment initiated at various times in the past and requiring, for confirmation, a hunt through the records. The least one can do is to box round the current treatment so that the most recent "box" represents the current regimen. In the end it is better to use a separate drug card, identifiable by colour (yellow being a frequent choice). This should not only show treatment but should also systematically identify drugs that may be repeated without consultation (and if so, for how long).

784

BRITISH MEDICAL JOURNAL VOLUME 282 7 MARCH 1981

(F) *Have a summary card in the notes listing briefly and chronologically the problems in the patient's past life that presently, or might in future, affect his health or longevity.* The problems may be physical, psychological, or social. Anyone who thinks he knows his patients well enough not to need a card of this kind should try picking out a fat file, listing the past problems he knows of, and then go through the file constructing a formal list—the unremembered items may be quite frightening. In any event one's partners and locums have little or no knowledge of a patient's past life, and a summary card of important events can be a godsend during an emergency call in the middle of the night.

(G) *Start registers.* If general practice is to maintain its identity and credibility there will be a need for doctors to look at their own performance in caring—to audit what they are doing. So there is a need to construct registers of the patients suffering from specific conditions that the doctor might wish to audit. This again is a personal choice, though I suggest the following:

(a) *Conditions requiring long-term maintenance treatment*—for example, hypertension, diabetes, thyroid disease.

(b) *Patients with conditions requiring long-term follow-up*—for example, gastroenteritis, anemias or contraceptive management, immunisation.

Pitfalls in Practice

Employment law

I: Why you need to know about it

NORMAN ELLIS

General practitioners who employ staff have certain legal responsibilities, but many of them are unaware of these. Whenever I meet a group of general practitioners I usually find that more than half of them have not provided written contracts to their staff, even though the law requires this to be done. This does not mean that no contract actually exists. A verbal (unwritten) contract exists when an offer of a job is made and accepted. And this contract can last for years on a verbal and unwritten basis, capable of being interpreted and enforced by the courts. But verbal contracts, although superficially attractive for their flexibility and apparent simplicity, may lead to serious problems if differences of opinion arise. This is why successive governments, both Conservative and Labour, have introduced legislation to promote written employment contracts.

Surprisingly but understandably the employment relationship is so often left undocumented. The general practitioner has little time to devote to the task of writing down the various details of a contract—for example, sick pay, holidays, pensions, and disciplinary and grievance procedures—and it often seems preferable to allow these to evolve "naturally" as the working relationship develops. A written contract is not only troublesome to prepare, but might also appear to be a happy way of introducing into it excessive formality and inflexibility.

(C) *Conditions in which the doctor has a special interest or knowledge.*

Good records are hard work, but any effort put into them will prove rewarding in both saved time and improved clinical effectiveness. Most of the changes advocated can be done in your own time or in spare moments, but remember to keep the initiative going; otherwise changes may take an age to accomplish. I try to try to myself when picking up an envelope, "Are these good enough?" or, "What will my partners think if they have to see this patient tonight?" If the answers are *No and Terrible*, I know what I should be doing.

I am grateful to my partner, Dr Paul Davies, for many helpful suggestions incorporated into this article.

Reference

¹ *Book of Scripts*. Revised ed. Harmondsworth: Penguin Books, 1968.

Because working relations between the doctor and his staff are close, the formal written contract is seen as an intrusion, bringing the formality of the law into what are essentially personal human relations. This reason is also given to explain why formalities are so often avoided. But the neglect of the written contract is also surprising. Irrespective of whether it is written out of a legal contract of employment does exist, and therefore the law already has a major part to play. The value of documentation is simply to state in writing the agreements and understandings that already define relations between the doctor and his staff. There are many other contractual relations—for example, partnership agreements, car purchase, television rental, bank loans, etc.—which no doctor would consider entering into without proper documentation spelling out in detail the agreement with the other party. But the employment contract may entail far greater liabilities and costs than even a rental or purchase agreement if it should turn sour. Yet the rights and obligations of this contract are often left to verbal understandings and agreements, which may so easily lead to misunderstandings and disagreements.

A simple precaution

I will not attempt to describe the plethora of employment law. This would be fruitless; the general practitioner has enough to worry about without trying to become an expert on employment matters. Instead, I advise general practitioners on a course of action that is simple to follow and which should help him to avoid most pitfalls.

British Medical Association, BMA House, Tavistock Square, London WC1H 9JP
NORMAN ELLIS, MA, PhD, under secretary and senior industrial relations officer