PRACTICE OBSERVED

Emergencies in the Home

Anaphylaxis and drug reactions

R L HERBERT

Type I hypersensitivity reactions result from the patient forming abnormal amounts of IgE antibodies in response to antigen exposure. The antibody attaches itself to the surface of the mast cells, which in response to antigen challenge break down and release histamine, 5-hydroxytryptamine, and slow-reacting substance of anaphylaxis from their granular contents. The reactions (also referred to as atopic or immediate) may vary from an irritating pimple through distressing bronchospasm to the extreme of circulatory collapse and death.

Managing mild hypersensitivity

This short paper is intended to deal with the immediate treatment of anaphylaxis in the home. Before dismissing the nuisance end of the scale, however, I want to make some observations and recommendations. Allergic rhinitis and urticaria often provoke reactions varying from discomfort to annoyance—not only from the public, but also sadly from many doctors. Patients with ugly and irritating rashes that cause them to behave like pariahs, students trying to sit final university examinations, and bus drivers working through floods of blinding tears and nasal fluid merit more than just antihistamines. While many patients respond well to antihistamines with no apparent sedation, my own opinion is that this is often a false state of security. The sedation produced may often be insidious, only being apparent after the exam has been failed or the child killed in a traffic accident. Nasal, ocular, and inhalable preparations of disodium cromoglycate and beclomethasone used either separately or in combination have made life much more tolerable for the atopic patient. A single intramuscular injection of either methylprednisolone

acetate or triamcinolone diacetate is often justified at the onset of the symptoms. The adrenal suppression, which can be proved only by sophisticated biochemistry, although a scientific reality is clinically a myth that may be safely ignored for all practical purposes. For those who still have reservations, however, it has recently been shown that many patients respond well to a single intramuscular injection of 120 units of corticosteroid. Even 5 mg-tablets of prednisolone on a daily reducing dosage (six the first day down to one on the sixth day and nothing on the seventh), which gives almost certain relief of symptoms with no interference in mental or adrenal state, should not in my opinion be withheld when the severity of the symptoms and the circumstances of the patient merit it. Picture, if you will, a bride and groom on the greatest day of their lives in a village church in England sometime in June trying to maintain their dignity while under attack by every pollen grain our countryside can muster.

Nevertheless, the mainstay of treatment must be accurate identification of the offending allergen and either their eradication from the environment or desensitisation. Repeated desensitisation courses or maintenance courses will often be required for a satisfactory result. This should still be the treatment of choice, however, and the previously-discussed treatments should be reserved for those patients who either present too late for desensitisation or fail to respond to it. Recently, the task has been made even easier with the availability of the Phadebas radioallergo sorbent test, which is capable of identifying IgE antibodies to specific allergens in 30 ml of blood. But for most of us careful history taking backed up with skin testing will suffice. It is also interesting to note that the same drug company now manufactures bee and wasp venom vaccines. There are five deaths each year in the British Isles from anaphylaxis to wasp and bee stings (not to mention the rest of the world), which is sufficient justification for costly research and production.

Desensitisation is a simple procedure even though time consuming. So long as basic precautionary principles are adhered to—that is, a syringe of 0.5 ml of adrenaline 1 in 1000

Chalford Health Centre, London E4
R L HERBERT, MA, CC, general practitioner

782

BRITISH MEDICAL JOURNAL VOLUME 282 7 MARCH 1981

drawn up ready, and the recommended injection regimen strictly adhered to—it has only minimal danger. If there should be a reaction at one dose then returning to the previous last dose that produced no reaction before advancing further through the regimen will almost always avoid disaster. In the past 15 years my desensitisation of about 50 patients each spring has resulted in only one case of anaphylaxis.

Severe anaphylaxis

While severe anaphylaxis is rare it is nevertheless a real and ever-present risk, which has little regard for fitness or age. Pooch Bear is credited with having said, "Accidents are funny things; you don't know they are going to happen until they do"; so often is the case with anaphylaxis. The availability at all times of the knowledge and materials with which immediately to intervene is mandatory.

In the first instance, the basic principles of the care of a shocked patient with increasing circulatory collapse apply as always. Fumbling about in a medical case or drugs cabinet searching for little-used drugs that when found may be out of date will often guarantee failure. By dialling 999 and passing the responsibility to the ambulance service you invoke a time-consuming series of events that you and your patient cannot afford. Your patient's circulatory state is progressively deteriorating, and soon the peripheral circulation will have shut down totally. Unless an intravenous cannula is inserted before this happens then the expertise of the hospital is at best hindered and at worst completely negated. Establishment of an intravenous infusion with either saline or Hartmann's solution before your patient's veins totally collapse is mandatory. It is equally essential to keep a cautious watch on the patient's cardiac state and airway if he becomes unconscious. If your patient, while you are cannulating his antecubital vein with him lying on his back, dies from anoxia resulting from an inadequately maintained airway, your hospital colleagues and the coroner are unlikely to be impressed. Finally, should the patient's heart stop despite an intravenous infusion, appropriate drugs, and airway maintenance external cardiac massage and mouth-to-mouth respiration must be started and must continue inside the hospital. Defibrillation and positive-pressure ventilation by those of us who have the necessary equipment would, of course, increase the chance of success, but this is rarely available.

Fortunately, most cases of anaphylaxis do not progress to such extremes and once the intravenous infusion is flowing the doctor can relax a little and make his choice of drugs.

Drugs for anaphylaxis

If the patient is not unconscious a very useful preparation, often forgotten, is sublingual isoprenaline in tablet form. Its speed of absorption will often give valuable time—the circulation will be maintained just long enough for the intravenous infusion to be set up. An immediate subcutaneous injection of 0.5 ml adrenaline 1 in 1000 if it is already prepared in a syringe and not in a bottle in a case or your cupboard will be equally effective. An 8 mg bolus of chlorpheniramine maleate into the infusion is the treatment of choice (4 mg in my experience is almost homeopathic); sedation such as that there is may serve you well in relating an often terrified patient. In such a potentially serious condition, however, I believe that the use of more powerful preparations than the antihistamines is not progress to such extremes and once the intravenous infusion is flowing the doctor can relax a little and make his choice of drugs.

treatment may also be of benefit. As a final gesture I suggest that the injection of 0.5 ml of adrenaline 1 in 1000 into the site of the original injection (if that was the causative factor) may slow down further absorption of the allergen and restrict the severity of the attack.

Treating in the home can be as effective as in hospital

I was asked to write on the treatment of anaphylaxis in the home. To me the title implies that there is a difference in treatment in the home as opposed to hospital. Although this is often so, I would submit that this is morally wrong and need not be so. Apart from the practical considerations of equipment and the great cost and physical size, there is no valid reason to justify the implied lesser treatment. The basic ideas of medical intervention in life-threatening illness or injury are the same, in my opinion, irrespective of the venue. To deny a patient in extremis medical expertise simply because he is either at home or in the street and enforce transport to hospital is to increase the risk of death.

While I accept that it is difficult to set up a drip singlehandedly, it is not impossible, nor is it impossible to deal with a cardiac arrest singlehandedly. With the advent of trained ambulance crew in intubation and infusion techniques there is now an ever increasing chance of such a technique surviving transport to your 999 call. The training programme, it is hoped, will gain in impetus so that in the near future all 999 calls will be responded to by such a team. In the past, as now, general practitioners have set themselves apart from the ambulance service, especially the men on the ground. We now see, however, a new phase developing where together ambulance men and general practitioners may function as a team and apply skills previously only thought possible in the hospitals. Our moral and NHS contractual obligations are to give medical aid to those who need, where and when they need. We must go to those who cannot come to us. A whole new era lies before us; we can grasp it and have a primary care team par excellence with general practitioner and ambulance crew equally skilled and working as a powerful unit.

Too many primary care problems, however, are passed on to the secondary care physicians in hospital with neither intervention nor support; very often resulting in their transfer to the ultimate care physician in the mortuary.

My thanks to Mrs Linda Herbert for patiently typing this paper.

This series on emergencies in the home will continue in a few weeks.

Searching for a practice

When you're interviewed for a practice your wife is also interviewed. The standard pattern is that you drive up, you are greeted by the doctor, and then your wife is whisked away to meet his wife and then she meets the other wife. I always remember in 1954 a large practice in the North and being interviewed in a fabulous house. Very impressed, I thought, "This is it, this is what we want." With the partners there we couldn't talk to each other, but when we had tea together my wife was one side of the room and I was on the other, and we'd look at each other, trying to signal. I was trying to say, "I think it's all right," but she was very cold and I didn't know whether she was just fed up with me or what. Anyway, when we got into the car and drove home, she explained that she'd managed to spend half an hour with the junior partner's wife and all the misery and the exploitation in the partnership had come out. This girl had just poured her heart out to my wife. We thought no more about that job.—KEN DICKINSON, general practitioner, Birmingham

We will be pleased to consider similar interesting stories about general practice.—Ed, BMJ

BRITISH MEDICAL JOURNAL VOLUME 282 7 MARCH 1981

783

Medical Records

Practical problems of improving records

DAVID STUART

"What is written without effort is in general read without pleasure." Samuel Johnson, *Johnson's Miscellany*, Vol II, p 309.

The average family doctor reading articles about improving records may well be forgiven for having a sigh of despair, tinged with envy of those few who, with superhuman energy, have pressed on and completed some ambitious scheme of recording, and who then insist on pricking his conscience by writing about it. For his own part, he is inheriting some years ago over 2000 envelopes, some unopened by a medical pen, some held together by brown rotting Sellotape and containing a miscellany of cards he cannot read, and a bunch of letters, a few of which contain vital information about his patient, but most recording normal blood tests or unimportant casualty attendances of yesterday. His partners regard any scheme for improving things with the usual mixture of enthusiasm and suspicion of anything that might require more work, and his secretary and receptionist already have enough on their plate, thank you. . . .

What to do first

There are several prerequisites for any systematic approach to the problem.

(1) Start off by defining the objective of an assault on the records. Who is going to benefit and in what way? Is the objective to improve the usefulness of the notes in everyday practice, or is it related to some other end—for example, audit, medico-legal considerations, or even making good recording a passing preoccupation? Do not expect an exercise without knowing what benefit you expect, or else you may labour mightily and achieve nothing. For instance, one sees age-old registers constructed with painstaking effort but then find that because the partnership concerned is not conducting research or audit and the county council computer is already recalling patients for research and insurance.

(2) Everyone in the practice must know about, and agree to, any action that is proposed—this means ancillary staff as well as the partners.

(3) The time spent on a project needs to be identified and organised. Who is going to do a job? Will time be spent regularly on it, or will it be done in spare moments? If the latter, has a target of achievement been set?

(4) Be prepared to use unskilled or semi-skilled labour for the routine work—the doctor himself will have enough skilled sorting out to do anyway.

How to proceed

Once a partnership has actively decided to effect some improvement in records, a number of objectives present themselves. What might

Great Yarmouth, Norfolk
DAVID STUART, MA, MRCGP, general practitioner

be regarded as essential or merely desirable must be a matter for individual choice, and the list that follows is my own, in my personal order of importance.

(A) *Get continuous cards and letters put into date order*—a basic task requiring no special skill. If existing ancillary staff cannot cope with the task in reasonable time, extra staff may be employed temporarily. At least one practice has used the Youth Opportunities Employment Scheme for this purpose. The partnership must decide:

(a) *What order to have the cards in.* Will the most recent face forwards or backwards? Will the letters be behind the continuation cards or vice versa?

(b) *How the continuation sheets should be attached to each other.* We push a hole in the top left corner and use treasury tags. Some prefer stapling the left edge. If left unattached, the sheets tend gradually to fall into chaos again.

(c) *Find out and destroy unnecessary continuation cards and letters.* This may be done by skilled ancillary staff or a nurse, though some doctors think that it should be done only by them. One can attach the task alphabetically, but the alternative is to select first the records that need weeding out most (usually the "fat files"). In this case one needs to get into the habit of picking out one or more daily, and the envelopes that have been dealt with need to be identified by a mark (for example, a coloured sticker) unless another task is being done at the same time that will identify the sorted envelope (for example, filling in a summary card).

(C) *Adopt colour coding for important diseases or facts.* The colours originally adopted by the Royal College of General Practitioners are now widely used, and a suitable sticker on the front of the envelope saves one from ignoring a hypersensitivity or that the patient is a known diabetic. Coloured stickers take hardly any time to put on and lead to some refinement of the doctor's own clinical thinking. I have diagnosed half-stickers to indicate either a family history of the disease concerned or a predilection potential such as an isolated high blood pressure reading or mildly abnormal glucose tolerance. If a disease register is to be kept, the sticker may be ticked to indicate that the patient's name has been entered.

The RCGP list is: **RED**—hypertension; **BLUE**—hypertension; **BROWN**—diabetes; **YELLOW**—epilepsy; **GREEN**—tuberculosis; **BLACK**—attenuated suicide; **WHITE**—long-term maintenance treatment.

(D) *Look at the layout of the notes.* Minor changes in layout can improve legibility greatly, for example:

(a) Is the writing legible? If not, can it be improved by deliberate self-training? Competent, if not beautiful, scripts are not difficult to learn.¹ If a doctor's handwriting is hopelessly illegible, having the records typed is the only remedy, though only time-wasting.

(b) How in diagnoses using colours to distinguish diagnosis from treatment registers.

(c) Leave one-line gaps between consultations.

(d) Have a system for transcribing out-of-hours notes to the main record. Some practices have the notes typed in, though we prefer to use the special addressed paper supplied by Winthrop Laboratories.

(E) *Have some system for identifying current treatment.* Nothing is more irritating to one's partner or locum than to be confronted by treatment initiated at various times in the past and requiring, for confirmation, a hunt through the records. The least one can do is to box round the current treatment so that the most recent "box" represents the current regimen. In the end it is better to use a separate drug card, identifiable by colour (yellow being a frequent choice). This should not only show treatment but should also systematically identify drugs that may be repeated without consultation (and if so, for how long).

784

BRITISH MEDICAL JOURNAL VOLUME 282 7 MARCH 1981

(F) *Have a summary card in the notes listing briefly and chronologically the problems in the patient's past life that presently, or might in future, affect his health or longevity.* The problems may be physical, psychological, or social. Anyone who thinks he knows his patients well enough not to need a card of this kind should try picking out a fat file, listing the past problems he knows of, and then go through the file constructing a formal list—the unremembered items may be quite frightening. In any event one's partners and locums have little or no knowledge of a patient's past life, and a summary card of important events can be a godsend during an emergency call in the middle of the night.

(G) *Start registers.* If general practice is to maintain its identity and credibility there will be a need for doctors to look at their own performance in caring—to audit what they are doing. So there is a need to construct registers of the patients suffering from specific conditions that the doctor might wish to audit. This again is a personal choice, though I suggest the following:

(a) Conditions requiring long-term maintenance treatment—for example, hypertension, diabetes, thyroid disease.

(b) Patients with conditions requiring long-term follow-up—for example, gastroenteritis, anemias or contraceptive management, immunisation.

Pitfalls in Practice

Employment law

I: Why you need to know about it

NORMAN ELLIS

General practitioners who employ staff have certain legal responsibilities, but many of them are unaware of these. Whenever I meet a group of general practitioners I usually find that more than half of them have not provided written contracts to their staff, even though the law requires this to be done. This does not mean that no contract actually exists. A verbal (unwritten) contract exists when an offer of a job is made and accepted. And this contract can last for years on a verbal and unwritten basis, capable of being interpreted and enforced by the courts. But verbal contracts, although superficially attractive for their flexibility and apparent simplicity, may lead to serious problems if differences of opinion arise. This is why successive governments, both Conservative and Labour, have introduced legislation to promote written employment contracts.

Surprisingly but understandably the employment relationship is so often left undocumented. The general practitioner has little time to devote to the task of writing down the various details of a contract—for example, sick pay, holidays, pensions, and disciplinary and grievance procedures—and it often seems preferable to allow these to evolve "naturally" as the working relationship develops. A written contract is not only troublesome to prepare, but might also appear to be a happy occasion spoiled by introducing into it excessive formality and inflexibility.

British Medical Association, BMA House, Tavistock Square, London WC1H 9JP
NORMAN ELLIS, MA, PhD, under secretary and senior industrial relations officer

(C) Conditions in which the doctor has a special interest or knowledge.

Good records are hard work, but any effort put into them will prove rewarding in both saved time and improved clinical effectiveness. Most of the changes advocated can be done in your own time or in spare moments, but remember to keep the initiative going; otherwise changes may take an age to accomplish. I try to try to myself when picking up an envelope, "Are these good enough?" or, "What will my partners think if they have to see this patient tonight?" If the answers are *No* and *Terrible*, I know what I should be doing.

I am grateful to my partner, Dr Paul Davies, for many helpful suggestions incorporated into this article.

Reference

¹ *Book of Scripts*. Revised ed. Harmondsworth: Penguin Books, 1968.

Because working relations between the doctor and his staff are close, the formal written contract is seen as an intrusion, bringing the formality of the law into what are essentially personal human relations. This reason is also given to explain why formalities are so often avoided. But the neglect of the written contract is also surprising. Irrespective of whether or not there is a legal contract of employment does exist, and therefore the law already has a major part to play. The value of documentation is simply to state in writing the agreements and understandings that already define relations between the doctor and his staff. There are many other contractual relations—for example, partnership agreements, car purchase, television rental, bank loans, etc.—which no doctor would consider entering into without proper documentation spelling out in detail the agreement with the other party. But the employment contract may entail far greater liabilities and costs than even a rental or purchase agreement if it should turn sour. Yet the rights and obligations of this contract are often left to verbal understandings and agreements, which may so easily lead to misunderstandings and disagreements.

A simple precaution

I will not attempt to describe the plethora of employment law. This would be fruitless; the general practitioner has enough to worry about without trying to become an expert on employment matters. Instead, I advise general practitioners on a course of action that is simple to follow and which should help him to avoid most pitfalls.

Firstly, I would advise any general practitioner who has not already done so to consider providing written contracts of employment to his ancillary staff. Of course, I would also advise the doctor to exercise great caution before rushing into the surgery with a bunch of written contracts. Such a gesture can be easily misunderstood, creating, rather than avoiding, problems.

Secondly, if care is taken in preparing the contract it will take full account of existing employment law and clarify many matters where there is uncertainty on both sides. The draft contract, which will appear in a later article, takes full account of existing statute law. It even takes account of health and safety legislation by including a policy statement on health and safety at work. (The Health and Safety at Work Act requires every employer of five or more staff to prepare a written statement of his general policy, organisation, and arrangements for health and safety at work.)

Finally, the BMA's regional offices can provide specialist advice and help to BMA members on how to introduce written contracts among ancillary staff. This service has already been widely used by members, and many general practitioners have already made use of the draft contract provided in a subsequent article.

Why is there so much employment law?

When I discuss employment law with doctors and explain its implications for general practitioners I am very often asked why there is so much employment law. My answer refers to the various influences that have contributed to its growth. (About 10 new Acts of Parliament have been implemented since the mid-1960s, establishing over 20 new rights for the individual employee.) Successive Conservative and Labour Governments have tried to influence the behaviour of unions by devising laws that would encourage them to act more reasonably and responsibly. As a quid pro quo for restrictions on closed shop agreements—for example, picketing, secondary boycotts, certain union—management have provided legal ground rules covering such matters as dismissal, employment rights, and health and safety at work. Three important episodes illustrate the "trade off" between these two sets of objectives: *In Place of Strife* and Mrs Barbara Castle's proposals for legislation; *Fair Deal at Work* and the 1970 Conservative Government's Industrial Relations Act; and the 1974 Labour Government's Employment Protection Act.

But this plethora of legislation is not merely one side of an equation designed to contain the challenge that unions pose for governments. Governments have also believed that the more formal and equitable approach to industrial relations and personnel matters required in these enactments would quickly reduce the number of disputes by improving industrial relations. Of course, there were other influences—comparisons with established international standards, for example, the Equal Pay Act and Sex Discrimination Act; the influence of legislation on other subjects such as the Race Relations Act; and the commitment of governments to a climate of increased job security, for example, the Redundancy Payments Act and Industrial Training Act—that would encourage employers to relax restrictive practices and accept changes in working practices and new technology.

The law has been designed, however, to alter the personnel practices and policies of large organisations where unions are already well established. The special position of the small employer, whose circumstances are considerably different, has been mostly ignored by legislators, but the 1980 Employment Act has made some very important concessions. Many of these ground rules—the rights of the employee and the duties of the employer—apply irrespective of the size and resources of the organisation or firm.

It should be remembered that most of this law has the support of both major political parties. Both have contributed to

its growth and both now accept it as a permanent feature of the industrial relations environment. On a few issues such as the closed shop, picketing, and strike ballots there is a main divergence of opinion between Conservative and Labour governments. Such a divergence has become a focus for people who have preferred to ignore the legislation have done so on the wholly erroneous assumption that the tide would rapidly recede after the change of government in May 1979. In fact, quite the reverse seems to have occurred: the body of employment legislation is still growing, as is evident from the 1980 Employment Act.

What significance does legislation have for the small employer?

Because governments have been heavily criticised for ignoring the problems of the small employer, some thought has now been given to the significance of the legislation. The previous Government made a survey of the problems of the small employer, which showed that the legislation had had a very limited impact. Only one small employer in 10 had had a case of unfair dismissal brought against him in the past two years. The survey also showed that small employers have very few disputes and problems. When asked, "What would you say have been the main difficulties you have faced in the past year in running your business?" employment legislation and health and safety regulations were hardly mentioned.

These findings could be said to vindicate those who have preferred to ignore the legislation and carried on regardless, believing that too much fuss has been made about the risks for the small employer, and that this survey gave a realistic perspective to it. This view may be acceptable to many employers who prefer to remain ignorant of the legislation (though ignorance of the law is no defence if legal proceedings should arise), but there are good reasons for being more cautious. Firstly, the survey's evidence was gathered about three years ago, and it is reasonable to assume that the climate of industrial relations has continued to change since then; more employers are becoming aware of the employment rights, and the process of unionisation is now penetrating general practice at an increasing rate. About eight unions actively compete for members—among them the Society of Medical Technicians, NALGO, NUPE, COHSE, MATSA, (GAWU) clerical and executive section, and ACTS (TGWU) clerical and executive sections—as the effects of changes in Health Service industrial relations spill over into this field.

I have focused on the need for a written employment contract because this offers an easy remedy for the many doctors who prefer to keep employment law at a safe and decent distance. In a later article I will provide a draft contract that should suit the circumstances of most general practitioners, together with advice on how it might be introduced.

Avoid litigation by good management

I must emphasise that the overriding aim should be to avoid litigation. When doctors ask for help with problems concerning their ancillary staff they often mistakenly look for legal solutions to problems that are caused simply by their failure to manage their staff effectively. For example, one doctor asked if the law could help him to persuade his receptionist to provide him with

*APEX, Association of Professional, Executive, Clerical and Computer Staff; ASTMS, Association of Scientific, Technical and Medical Staff; NALGO, National and Local Government Officers' Association; NUPE, National Union of Public Employees; COHSE, Confederation of Health Service Employees; MATSA, Medical Association of Technicians; ACTS, Association of Clerical and Technical Staff; TGWU, General Transport and General Workers' Union; TGWU, General Transport and General Workers' Union.

Beyond the Surgery

General practitioner as social anthropologist

Cecil G. HELMAN

Most social anthropologists these days do their research in remote areas of the non-industrialised world. They live for a year or two in tiny jungle villages or isolated settlements studying small-scale tribal societies. Gradually, by inquiry and observation—the process of "participant observation"—they build up a picture of the tribe's culture, its customs and rituals, its political and economic structure, and they learn to view the world through tribal eyes. Later they return to their universities laden with ideas and artefacts and try to relate their findings to more complicated Western societies in which they live.

As an anthropologist my "tribe," as it were, is somewhat less exotic: it is my own patients and those of neighbouring GPs on the fringes of north-west London. There are no tropical forests here, no parkland squawking in the jungle, no crocodiles, no bamboo huts or cooking fires. Only rows of semi-detached houses containing suburbanites, with mortgages and marital problems, neat front gardens, and a Main in every driveway. But even in this sedate setting there are rich pickings for the medical anthropologist with an interest in medical folklore.

Medical anthropology

Medical anthropology is a fairly new branch of social anthropology. It is the study of how people in different groups or cultures explain the causes of illness, the types of treatment they believe in, and whom they turn to if they do get ill.¹ The subject includes more than a list of exotic medical practices from faraway lands; any culture's theory of illness and its treatment are greatly influenced by its view of the world in general and of man in particular. Illness is only a special type of misfortune, part of a wider spectrum of adversities which, in some societies, are all said to have the same cause. For example, when one has a traffic accident, loses money on the stock exchange, and gets a violent toothache all in the same day one may ascribe it to something called "a streak of bad luck," as though there were a single causal force or agent; similarly in other societies all these bad events happening together may be taken as proof that the victim has been "bewitched" by some malevolent person. Medical anthropologists study these various beliefs and how they affect patients' behaviour; they compare the differing views of health and illness of different societies and also of different groups (say, doctors and patients) within one society. Problems of communication between general practitioners and their patients in an English suburb² are as much of interest to the modern medical anthropologist as, say, some exotic childbirth ritual in Outer Mongolia.

My own interest in the subject began several years ago. I remember, in Africa, seeing a grossly infected wound on a patient's leg. Among the purulent matter were a number of

London N16 3JR

Cecil G. HELMAN, MB, ChB, general practitioner

small dark bristles. I found out later that the wound was originally a dog bite and that these bristles were hairs cut off the offending animal. The patient, it appeared, had taken quite literally that old saying: "The hair of the dog that bit you." He had caught the dog, cut off its hair, and rubbed it into the wound with highly purulent result! This bizarre incident—and many others—led me gradually to medical anthropology. I had always been fascinated by folklore, old myths and legends, and curious superstitions, especially those relating to health and disease. But to look at these beliefs and their clinical significance required a more academic approach. Two years after my housemanship, the local locums, and a short spell as a ship's doctor in the Med, I enrolled as a full-time student at University College London and graduated in social anthropology. For almost seven years I have been a full-time GP, but have been also since 1977 an honorary research assistant in anthropology at University College, and a co-founder of the British Medical Anthropology Society.

I find that general practice, even full-time, offers an excellent setting for medical anthropological research, in my case into the clinical significance of folk beliefs about illness and health.³ The two types of data gathered by anthropologists—"hard" or statistical data and "soft" or personal observations and open-ended interviews—can both be gathered by an astute GP working in any area where there is a wealth of folklore beliefs about illness.

The importance of these beliefs is that they affect the behaviour of patients, and of those around them. One classic study on this theme was done by Dr Louella Snow, of Michigan State University. She analysed the folk beliefs about menstruation and fertility of working-class women attending a city clinic in the United States. She found a variety of odd ideas about the uterus and its function, which clearly had an effect on the women's health. For example, some women viewed the uterus as a hollow organ that is tightly closed between menstrual periods, while it slowly fills up with tainted blood, then opens to release this blood at the time of the period. This belief led these women to assume that they were most likely to get pregnant just before, or just after a period "before it is tightly closed again." As a result they could not see the logic of taking contraceptive precautions at other times of the month. They also believed that a woman could not contract venereal diseases during pregnancy "because the uterus is closed and the germs cannot enter."

Colds, chills, and germs

A prime aim of medical anthropology is the improvement of health care, and the beliefs and behaviour of patients obviously may have important clinical implications. So far I have conducted and published two research projects on this theme.⁴ The first⁵ was a detailed examination of folk beliefs about "colds," "chills," and "fevers," to find out the concepts and premises that lay behind the old aphorism "Feed a cold, starve a fever." Over four years I asked several hundred patients, as

a cup of tea twice a day, morning and late afternoon. (At the time I met the doctor he was actually making tea twice daily for his receptionist.) This doctor had not established contact with a manager 14 years earlier when he first contacted that his receptionist should no longer undertake this task as part of her job only a few days after she started work. At the time he did not like to say anything because it would cause unpleasantness. And he hoped that the difficulty would improve. But the question of who made the tea was just a symptom; it was the problem of their working relations that had been unsatisfactory from the outset, and these had deteriorated further over the years. The law as such had little to offer this doctor in the way of a remedy. The problem had been going on for a long time, too long, and it needed more than legal action to put it right.

This example is typical of many cases where BMA advice is sought. It illustrates some of the advantages of a more formal approach to the employment of staff. It shows how a firm foundation for good working relations. The written contract also describe the disciplinary procedure that would be involved if performance in the job was unsatisfactory.

The prescription of a more formal approach to such matters, relying on properly agreed and documented contracts, might be

regarded as oversteering to something of minor importance. But this is simply not so. The relation between the doctor and his receptionist may seriously undermine his efficiency as a practitioner. And where there is only a small number of staff, the problems are often more difficult to deal with.

In my experience many of the problems of general practitioners are caused by poor selection procedures when staff are first appointed, inadequate supervision, and a reluctance to act decisively when staff are found to be unsatisfactory. Of course, for the general practitioner, who employs only a few staff it is pointless to recommend formal procedures drawn from the practices of large organisations for the recruitment, appointment, and discipline of staff. But at least one of the most basic and commonsense principles should be applied to these matters. The general practitioner should take great care to recruit the right staff; they should serve a probationary period (say, nine months) with effective procedures for assessing their performance, and the general practitioner should not hesitate to take firm action if their performance fails to meet the standard required.

This is the third of six articles on employing staff.

Another time we went to a seaside town to see a practice. A smooth doctor said to me, "I'm no businessman," which is a classical start. "I'm no businessman," he said, "but it would help you a lot if you purchased the house containing the practice. Because I don't think you can afford the main surgery, but I'll rent the accommodation to you." We went to see the beach surgery, which was a ghastly little suburban house. He was going to London. He was a single-handed general practitioner and had no right to do this at all because it was an executive council vacancy, which if he left would be advertised. What he was going to do was to take on a partner, and once he'd got him installed he'd move up to London. The condition would be that once you were in he would resign and you'd move into the next.

We went for another interview with three doctors in a nursing home. They had a list of only about four or five thousand, and I wondered how on earth three doctors could survive. One doctor was leaving and he was the usual remark that you had to buy his house and he would leave. They had a list of only about four or five thousand, and I wondered how on earth three doctors could survive. One doctor was leaving and he was the usual remark that you had to buy his house and he would leave. They had a list of only about four or five thousand, and I wondered how on earth three doctors could survive. One doctor was leaving and he was the usual remark that you had to buy his house and he would leave.

Clinical Corrie

All was a Chaucer legend who spent the war years in Jersey, under German occupation. During that time, when he was in his 20s, he had a right orchidectomy for what he was told was a malignant tumour of the testis. I first saw him as a patient in the 1960s, when he took on the job of wardens at one of the local old people's homes. Ever since his operation his scrotum had been very sensitive to "bryophobus-like" appearance. The penis was almost buried in the swollen disease. I arranged for him to see a surgeon in London, but no treatment was offered.

Six or five years after I had first seen him he came to see me with a smile on his face. Two days earlier he had been to a foot and mouth clinic. The doctor had said to him, "You've got a foot and mouth disease, but don't worry, it's not serious, it's just a virus. The back fence to get a better view—and had lost his oedema. In some way, whatever was obstructing the flow had been unblocked. I never found out why—ERNEST CANNON, general practitioner, London.

We will be pleased to consider for publication other interesting clinical observations made in general practice.—Ed, BMJ.

Cold hands make lighter work

Dr Romeo was a super doctor: kind, considerate, conscientious, and hardworking. But even he had his limits; he had a particular woman who was always sending for him, and in those days when a patient called you had to go. He really got sick of her. One night—a cold, miserable night—he received a wretched phone call from her. She had abdominal pain, and Dr Romeo knew that she had what we'd now call irritable bowel syndrome. He went out, but as he did so he thought, "I can't stand this." It was icy outside and he had put on all his heavy clothes, but he relented the weather and drove to her house with his hand out of the window. When he finally reached her house, he rang the bell and was let in by her husband (who was equally fed up). He rushed upstairs, threw his coat and hat, and he was in the room in a flash. He said, "Where is the pain?" The shock cured her of abdominal pain.—KEN DICKINSON, general practitioner, Birmingham.

Searching for a practice

In 1956 it was hard to find a partnership in general practice, and with the current numbers of general practitioners it is likely to become harder again in the next few years. In 1956 we were all much the same, just as trainees are now: we were all ex-army, had all old obese patients, and had all done traineeships. We were exploited by poor pay, long duties, having to cope with all the obstacles, and the problems of house purchase. My experience of 1956 might be of interest to present-day trainees as they may well experience similar things.

My wife and I had a routine: every Sunday we would go through the BMA, select jobs, write out applications, and send them off enclosing a hand-written curriculum vitae. Every Sunday we would go and see the interviewers, and we were never offered a job, but sometimes we were. On Monday, having been rejected for the job, we would have been told by a bottle of wine.

Once an advertisement appeared for a practice in an attractive country town in Herefordshire. We went there and a pleasant and delightful Irish doctor showed us his home. It quickly became clear, however, that his aim was to sell this house, which was an old rectory standing by a beautiful church. We couldn't understand what this was all about: why he was offering to sell his house? We found that he was leaving the practice to his wife, who was a very good one, and was coming to purchase his house, and at that moment his partners didn't even know he was going. When I rang up the partnership the following week to discuss some of the terms they were horrified.

There was a doctor at Graston who did the same sort of thing. The senior partner put in an advertisement but the junior partner always managed to intercept him and tell him that he was not to buy his house, because they really wanted this partnership to go to my house and say quite honestly unless I sell my house for the proper price I'm not leaving. And the senior partner, of course, wanted him out and was quite prepared to accept this sort of thing happening.

well as locally recruited surgery staff, a number of key questions such as, "What is a cold?" "Is a chill?" "Is a fever?" "What do you mean by a germ?" and so on. At first these questions drew puzzled frowns ("He's an odd 'un", that doctor; doesn't even know what a germ is!") but gradually the patients got used to it, and were often surprised by the extra attention.

By this ethnographic technique I built up a picture of patients' lay beliefs about these conditions. The research revealed a host of unexpected and exotic beliefs. For example, "germs" the vast majority of patients have never seen "a germ" nor have they any perceptual evidence for its existence. For most of them "germs" remain a hypothesis, a folk theory of causality, much like the "spirits" and to cause illness in some Third World societies. The stated attributes of "germs" were as follows: they are living, invisible, malevolent entities. They have no free existence in nature, and exist only in or among people. All "germs" are "bugs," and the distinction is made between "germs," "bugs," and "viruses." They are thought of as occurring in a cloud of tiny particles, or as a tiny invisible "insect." Germs causing gastrointestinal symptoms seem to be more "insect-like," and "germs" causing respiratory symptoms seem to be more "bug-like." Personalities which are expressed in, and can be recognised by, the symptoms they cause. ("I've got that germ, doctor, you know—the one that gives you the dry cough and the watery eyes"—or "the one that gives you diarrhoea, and makes you bring up.") Germs seem to be single in their effect; you are only attacked by one germ at a time ("I've got a germ, doctor"). They select their victims randomly, and therefore the victim is blameless and can mobilise caring relatives around him.

This is very different from "colds" and "chills," which are byproducts of one's personal battle with the environment, and usually result from doing something "fally," such as "going out in the cold after a hot bath," or "standing in the rain after a haircut." The "germs" enter the body by its orifices, the natural breaks in its defences, then move on to attack other parts of the body ("It began as a sore throat, but then went down to my tummy") and causes a fever. Sometimes fevers occur in a "dry" form—for example, pyrexia with a dry cough—sometimes in a "wet" form, where the pyrexia is accompanied by excess body fluid in the form of phlegm, mucus, saliva, diarrhoea, or sweat; the aim of folk therapy seems to be to move the patient from the "dry" to the "wet" state and thereby expel the "germs." Fluids are often used to effect this expulsion; patients with "dry" coughs are "coughed" their cough "loosen" so that they can "cough it off my chest," and so continue to demand cough lozenges. Wilks⁶ has noted that an estimated six million gallons of cough mixtures are prescribed every year under the National Health Service; could it be that part of the demand for these mixtures is that they are seen by patients as somehow "loosening" and then "washing out" the offending germ from the patient's chest?

Always go late at night

I once had a patient who was grossly overweight, lived in filthy circumstances, had a few cars, and a transport driver who came on a Thursday with diarrhoea. His family read on the Friday and said he'd still got diarrhoea and some vomiting. I said, "Well, carry on with the health and medicine, but I shall know in the morning if it's not serious and I'll come and see him." And about 2 o'clock in the morning they rang up and said he'd got a lot more diarrhoea—he was trying to go to the toilet two or three times every hour. I said, "Right, I'll move him to the fever hospital; he's obviously got acute gastroenteritis."

The discharge letter said, "Dear Dr Dickinson, Thank you very much for admitting this man. When we opened him we found an acute peritonitis."

It is logical from the patient's perspective to demand antibiotics when they are told they only have "a virus." In the folk model no differentiation is made between bacteria and viruses; both are "germs" that must be washed out or killed in situ by antibiotics. In this case, as in others, the value of in-depth medical anthropology is that one can get clues to the underlying premises governing patients' health behaviour, and also to their attitudes to prescribed medical treatment. My second study⁷ dealt with this theme—that is, patients' perceptions of their psychotropic drugs. It revealed a wealth of lay beliefs about these drugs and their supposed effects, which bear little relation to those of the medical profession. Once again I have found that viewing medical treatment and diagnosis from the patient's perspective can be a valuable experience.

Making sense of a dangerous world

Lay views of illness deserve our serious study because they are patients' ways of making sense—however imperfectly—of an unpredictable and dangerous world, where disease and death strike in an apparently random way. Hence all the superstitions about "good luck" and "bad luck," about "knocking on wood," and all the other "just in case" remedies, from which even doctors are not immune. I think there may well be a partial return to these "old wives' tales" and self-medication as a result of the increased prescription charges, as well as a sense of disillusionment with overtechnical medicine.⁸ If this drift back does occur then the medical anthropologist is well placed to analyse what has happened, and to help design a better health service for the future.

In the meantime, ask the next patient to come in, please....

References

- Helman CG. What is medical anthropology? *BMJ* 1980; 1:155-63.
- Lundy D, ed. *Culture, disease and healing: studies in medical anthropology*. London: Collins Macmillan, 1977.
- Helman CG. "Feed a cold, starve a fever": folk models of infection in an English suburban community, and their relation to medical treatment. *Medical Anthropology*, 1979; 8:107-37.
- Helman CG. Lay and medical attitudes to illness. *BMJ* 1980; 1:151-3.
- Stanton G, Webb B. *Going to see the doctor: the consultation process in general practice*. London: Routledge and Kegan Paul, 1975.
- Helman CG. Patient expectations of psychotropic drugs. *J R Coll Gen Pract* 1981; 31:107-12.
- Snow LF, Johnson SM. Modern day ancestral folk: some clinical implications. *JAMA* 1977; 237:278-9.
- Wilks E. The treatment of cough in general practice. *Prescriber's Journal* 1974; 1:107-12.
- Helman CG. An anthropological view of the modern hospital. *BMJ* 1980; 1:137-41.

His appendix had obviously perforated two days earlier. The lesson I learnt from this is really very simple: if ever you get a call after eleven at night, go; it's always possible. Whenever the message is you go, because patients never ring after midnight unless it's genuine. I felt as guilty then when I only went out on my own—once or five times over the years—as they were still grateful to me if I said they could stay on my list. I hadn't the courage to say no. And the cross I bore to go to the long drive to the hospital, it explained my guilt.—KEN DICKINSON, general practitioner, Birmingham.

We will be pleased to consider similar interesting stories about general practice.—Ed, BMJ.