problem as a long-term effect of treatment with guanethidine or thioridazine hydrochloride. A high proportion of our patients receiving the α_1 -blocking agent indoramin described a similar phenomenon. Failure of ejaculation with indoramin is probably due to its action on transmission at the short adrenergic neurones of the male reproductive tract, although further studies are indicated, including studies of urine obtained before and after masturbation, to determine whether retrograde ejaculation occurs.

Although a few reports¹⁻⁵ have mentioned sexual dysfunction in a small number of patients, the high incidence of failure of ejaculation without disturbance of erection and with the sensation of orgasm has not been emphasised. Two-thirds of our patients suffered from failure of ejaculation while taking the fairly low daily dose of 60 mg indoramin, and this emphasises the need to question patients about sexual performance when they are being treated with this or other α -adrenergic-blocking agents. This finding may also support the hypothesis that this group of drugs may be developed for use as reversible male contraceptives.⁴

We thank Dr C Mawdsley for allowing us to study his patients. The drugs were supplied by Wyeth Laboratories.

Requests for reprints should be addressed to Dr B Pentland, University Department of Medical Neurology, Royal Infirmary, Lauriston Place, Edinburgh.

¹ Lewis PJ, George CF, Dollery CT. Clinical evaluation of indoramin, a new antihypertensive agent. *Eur J Clin Pharmacol* 1973;6:211-6.

² Bianco S, Griffin JP, Kamburoff PL, Prime FJ. Prevention of exercise-induced asthma by indoramin. *Br Med J* 1974;iv:18-20.

³ Wainscott G, Volans GN, Wilkinson M, Faux GA. Indoramin in prevention of migraine. *Lancet* 1975;iii:32-3.

⁴ Kedia KR, Markland C. The ejaculatory process. In: Hafez ESE, ed. *Human semen and fertility regulation in men*. St Louis: C V Mosby, 1976: 497-503.

⁵ Faerchtein I, Roque AF, Kastansky I, Campos JC, Puppini S. A placebo controlled trial of the α -blocker, indoramin, in the treatment of arterial hypertension. *Curr Med Res Opin* 1976;3:675-84.

(Accepted 9 February 1981)

Royal Infirmary, Edinburgh

BRIAN PENTLAND, MRCP, lecturer in medical neurology
DOROTHY A ANDERSON, MSC, MPS, principal pharmacist, Lothian drug information centre
JULIAN A J H CRITCHLEY, MRCP, lecturer in therapeutics and clinical pharmacology

Scombrototoxic atrial flutter

Scombrototoxic fish poisoning has been recognised as a major form of ichthyosarcotoxism in many parts of the world.¹ Scombridae include tuna, bonito, and mackerel, and poisoning occurs after ingestion of fish that have formed a histamine-like toxin. Once produced this toxin is remarkably resistant to further preparation; to prevent its formation immediate packing at 0°C is essential as soon as possible after the fish have been caught. If fish are allowed to remain at room temperature tissue concentrations of histamine rise rapidly, possibly due to the action of bacteria on available histidine residues.

Case report

A 60-year-old man was admitted with a four-day history of palpitation that had started after a large meal that had included smoked mackerel. Within two hours he had developed flatulence, upper abdominal pain, and nausea, followed by weakness, pulsating headache, a sensation of fear, itching of the scalp, increased frequency of micturition, and rapid regular palpitation. After 48 hours he had sought advice and was referred to us. He was in atrial flutter with a ventricular rate of 150/minute; blood pressure was 160/90 mm Hg, with no evidence of cardiac failure. Heart sounds were normal. Routine blood investigations, including blood count, erythrocyte sedimentation rate, cardiac enzymes, urea, electrolytes, and thyroid function tests, were all normal. Chest radiography showed normal heart size and no pulmonary congestion. He reverted spontaneously to sinus rhythm the next day. Arrhythmia did not recur during follow-up of 18 months, which included four 24-hour ambulatory electrocardiograms taken in relation to similar meals.

Of 10 people who had also eaten the mackerel at the dinner, three had developed headache and itching of the scalp and six had felt generally unwell

for a few hours. We were unable to obtain a fish from the same batch for analysis.

Comment

Our patient developed classical symptoms of scombrototoxicity immediately after eating smoked mackerel. Many of the features of this condition are similar to those encountered in histamine poisoning and may be relieved by antihistamine drugs (H_1 -receptor blockers).¹

Though histamine concentrations in affected fish often exceed 100 mg/100 g,¹ well-documented episodes of scombrototoxicity have been reported where no excess of histamine was found, and non-scombrotoxic fish, such as herring, may have high histamine concentrations without producing symptoms. Thus the toxin is probably a more stable histamine-like compound.¹

In scombrototoxicity the clinical presentation often includes a rapid weak pulse and palpitation; Gilbert *et al*¹ found six episodes of palpitation among 30 incidents of poisoning, but electrocardiograms were not recorded. Histamine is released in large quantities in anaphylactic reactions, and its toxic effects² on the myocardium may cause some of the deaths associated with this condition. In severely affected cases electrocardiograms have occasionally been recorded and have disclosed arrhythmias, which may be supraventricular—for example, atrial fibrillation.³ In our patient the histamine-like toxin may have induced atrial flutter, which persisted after the other effects of the toxin had disappeared.

In most cases of scombrototoxicity palpitation may be sympathetically mediated, as a result of the peripheral effects of the histamine-like compound, and hence readily abolished by H_1 -antagonists. Direct stimulation of the heart via H_2 -receptors, however, may activate a magnesium-dependent channel,⁴ which may give rise to arrhythmia, as in this patient. In view of the potentially serious complications of histamine-receptor stimulation, the cardiac rhythm of patients with scombrotoxic poisoning should be observed before treatment with histamine antagonists is started.

Nearly 30 years ago atrial flutter was observed after a "heavy seafood meal."⁵ Perhaps closer scrutiny of dietary factors may help explain some paroxysmal arrhythmias.

¹ Gilbert RJ, Hobbs G, Murray CK, Cruikshank JG, Young SEJ. Scombrototoxic fish poisoning: features of the first 50 incidents to be reported in Britain (1976-9). *Br Med J* 1980;281:71-2.

² Peters GA, Horton BT. Continuous intravenous administration of histamine: effect on the electrocardiogram and serum potassium. *Am Heart J* 1944;27:845-58.

³ Booth BH, Patterson R. Electrocardiographic changes during human anaphylaxis. *JAMA* 1970;211:627-31.

⁴ Späh F, Fleckenstein A. Evidence of a new, preferentially Mg-carrying, transport system besides the fast Na and the slow Ca channels in the excited myocardial sarcolemma membrane. *J Mol Cell Cardiol* 1979; 11:1109-27.

⁵ Hejtmancik MR, Herrmann GR, Bradfield JY. Atrial flutter. I. Clinical aspects. *Am Heart J* 1950;40:884-90.

(Accepted 12 February 1981)

Ealing Hospital, Southall, Middlesex UB1 3HW

LESZEK BORYSIEWICZ, MB, MRCP, medical registrar

Royal Postgraduate Medical School, London W12 0HS

DENNIS KRIKLER, MD, FRCP, consultant cardiologist

ALEXANDER. It is called Alisander, Horse-parsley, and Wild-parsley, and the Black Pot-herb; the seed of it is that which is usually sold in apothecaries' shops for Macedonian Parsley-seed. It is usually sown in all the gardens in Europe, and so well known, that it needs no farther description. It flowers in June and July; the seed is ripe in August.

It is an herb of Jupiter, and therefore friendly to nature, for it warms a cold stomach, and opens a stoppage of the liver and spleen; it is good to move women's courses, to expel the afterbirth, to break wind, to provoke urine, and helps the stranguary; and these things the seeds will do likewise. If either of them be boiled in wine, or being bruised and taken in wine, is also effectual against the biting of serpents. And you know what Alexander pottage is good for, that you may no longer eat it out of ignorance but out of knowledge. (Nicholas Culpeper (1616-54) *The Complete Herbal*, 1850.)