PRACTICE OBSERVED

Law and the General Practitioner

Prescription medicines

STUART CARNE

Under the Medicines Act 1968 all medicinal products fall into one of the following categories:

- General sale list medicines, which may be sold at any shop (and in a few instances from automatic vending machines);
- Pharmacy medicines, which may only be sold by a registered pharmacist;
- Prescription only medicines, which may only be sold by a registered pharmacist on the prescription of a doctor, dentist, or veterinary surgeon (midwives, chiropodists, ophthalmic opticians, and physiotherapists may also prescribe certain of these drugs); or dispensed by such practitioners.

Under the Misuse of Drugs Act 1971 certain "dangerous or otherwise harmful" drugs are placed in the controlled drugs list. They may only be dispensed on prescription and are subject to additional safeguards against misuse (see below).

Writing a prescription

While in theory a prescription for a general sale list medicine or a pharmacy medicine could be written in any form understandable to the pharmacist, in practice all prescriptions written by a registered medical practitioner should conform to the regulations governing prescription only medicines. The prescription must:

- Be typed or handwritten in an indelible ink.
- Give the address and usual signature of the prescribing doctor.
- Give the date on which the prescription is signed or, in the case of a postdated prescription, the date on which it becomes valid.

Goldhawk Road, London W15 8EJ
STUART CARNE, *senior, general practitioner*

valid. Except for certain repeat items, the prescription will remain valid for six months from that date.

— Give some indication—for example, the qualifications—of the prescriber, indicating that he is a registered medical practitioner (or dentist, etc.). National Health Service (NHS) prescriptions form (FP10) have the doctor's name stamped on them and this suffices for identification.

— Give the name and address of the patient and, if under 12 years, the age. (This need only be given to the nearest month for infants under one year or to the nearest year for those older.)

By tradition a prescription also contains the instructions ("Signa") the pharmacist is to put on the label. These are usually written as abbreviations: *ps*—to take in the sumendum, post cibum (three times a day after meals). Previous editions of the *British National Formulary* gave the quantity of non-controlled drugs the pharmacist was to supply if the doctor omitted the amount on his prescription. The latest (1981) edition omitted this. The pharmacist is now encouraged to use his judgment in determining what quantity would be appropriate in such circumstances. It is generally held to be good prescribing to include this information on every prescription, however, either as the quantity—for example, 20 tablets of an antibiotic or 100 ml of a cough linctus—or the number of days the remedy is to be taken. Form FP10 has a space on it for the doctor to write the number of days for which the drug is to be taken. If there is more than one item on the form the use of this box may achieve quite an economy in the physician's writing time as well as a possible reduction in waste because one drug has been prescribed in a greater proportion relative to another. (NB: This "number of days" formula cannot be applied to local applications where the amount used per day is variable; nor may it be used for controlled drugs.)

Provided the signature is that of the prescribing practitioner

*Signa is Latin for label. It may also be an abbreviation for signetur (let it be labelled) or signetur (let them be labelled). The advantage of the abbreviation sign is that it covers all three possibilities.

drug may be repeated without the patient being seen is also stated on the card. When he needs a further supply of one or more items on the card, the patient sends his card to the surgery indicating the items he wants and calls at the surgery to collect his prescription either from the receptionist or, sometimes, from a rack in the waiting room. In some practices repeat prescriptions are posted, though the patient is usually expected to supply a stamped-addressed envelope. If the number of repeats is restricted the patient will be notified when he collects his "final" repeat that on the next occasion he must arrange to see the doctor.

In practices that do not control the number of repeats and a patient is entitled to collect his prescriptions without seeing the doctor there is a risk that some patients will go on taking medication they no longer need or take them in doses which are either too big or too small.

In many practices filling in repeat prescriptions is left to a nurse, receptionist, or secretary. Before signing it, a doctor checks the items against the patient's medical record card or repeat prescription form, or both. There is anecdotal evidence that in some practices the receptionist or secretary is given a supply of blank, signed NHS prescription forms, which she fills in when a patient requests a repeat prescription. (Some receptionists are also said to prescribe for "first events"—for example, cough mixture for a cold or antibiotics for a sore throat—working to a protocol prepared by the doctor, but not having his diagnosis or prescription checked.) There is evidence that there are many errors in prescriptions written by receptionists (and allegedly checked by the doctor).¹ How many of these errors are important and how many are due to clerical errors, such as spelling mistakes, which nevertheless leave the prescriber's intent beyond doubt—is not clear, and further studies are needed.

It has been argued that certain items might be supplied on the NHS by a pharmacist without a doctor's prescription. This would certainly save the doctor's time, but there are snags. From the financial point of view it is difficult to say what control the government would have on this expenditure. At present, by monitoring every general practitioner's prescribing costs once or twice a year a degree of control of these costs can be maintained; what analogous control could there be for prescribing by the pharmacist? Secondly, except for dispensing doctors, the person writing the prescription (the doctor) is not the person being paid for dispensing the drug (the chemist). Even in the case of dispensing doctors, these costs can be monitored against those of non-dispensing doctors.

From the clinical point of view it can be argued that the pharmacist's training does not equip him to make a diagnosis, which most physicians would regard as a prerequisite to prescribing. Nevertheless, pharmacists have always offered advice and often they select a medicine for their customers. ("Can I have something, please, for my 2-year-old son's cough?") To an extent this practice is limited by the fact that such medicines have to be paid for, and in my opinion the cost does provide some control. If it ever came to a choice between the doctor's recommendation or a qualified pharmacist deciding what is to be prescribed then the case for the pharmacist prescribing may be more difficult to oppose.

Drug trials

A doctor is entitled in the course of his practice to compare any two (or more) standard treatments. For example, he may compare the use of aspirin with paracetamol in relieving pain; or one *British National Formulary* cough linctus with another *BNF* cough linctus to see which is the most effective under specific conditions.

With a new product a clinical trial is only permitted if the drug is to be tested for a condition listed in the terms of its licence. For example, a new antihistaminic drug may have been approved for use in osteoarthritis. Any physician or group of

physicians could legitimately test that product in patients suffering from osteoarthritis but not in patients suffering from, say, gout. Nevertheless, an individual doctor could carry out a therapeutic trial on an individual patient. Thus, using the example above, a doctor may prescribe the drug for one or more of his patients on an individually named basis to see if it will relieve that patient's gout. If he wanted to conduct a trial in several patients suffering from gout, in which the patients would not receive the drug on an individually named basis, the doctor would have to get (from the DHSS) a Clinical Trial Certificate to obtain a supply of the drug from the manufacturer for use in the trial. Alternatively he could apply to the DHSS for an exemption from holding a Clinical Trial Certificate. This would usually be granted unless the proposed trial posed some hazard, the degree of which could be assessed only by examination of data, such as toxicity or bioavailability. Such an exemption permits a manufacturer to supply the drug for a purpose other than as stated in the Drugs Licence.

So far as a general practitioner is concerned it is more likely that if he is involved in a trial, it would be as a member of a team rather than as an organiser of the trial. In such cases, he must ask himself:

— Is the design of the trial satisfactory? (He may have very limited personal knowledge or experience of these matters, and in practice he will have to go on the reputation of those organising the trial.)

— By putting any of his patients into the proposed trial is he depriving them of the rights or benefits to which they are entitled? This would apply if, for example, he were asked to test a new peptic ulcer treatment against a placebo when it is known that antacids (which are safe) would give the patient some benefit. In testing the new peptic ulcer treatment it must not, therefore, be compared with a placebo but with an accepted alternative active remedy. Using placebos in a trial when there is a known proved active remedy for the condition is in most cases considered unethical.

A common form of drug trial in which general practitioners are asked to participate is that carried out when a new formulation of a product is introduced—for example, the testing of a combined beta-blocker and diuretic. These trials provide an acceptable treatment and therefore do not deprive the patient of any potential benefit. The assumption is that the individual components have been tested and found satisfactory and that their safety factor is known. What the manufacturer—and future prescriber—wants to know is whether this new formulation is an improvement on the individual components.

Legal liability in a clinical trial

Legal liability has never been tested in courts, but it would seem unlikely that a general practitioner could be sued by a patient he has put into a trial, provided he complied with the guidelines above and provided he obtained the patient's informed consent. Should any mishap befall the patient—for example, an unsuspected drug reaction—it is almost certainly the manufacturer who would be responsible (and who would, anyway, have insured himself against such an event).

Much publicity has been given to a proposed EEC regulation that should a patient suffer any harm from using a prescribed medication the physician who wrote the prescription would be liable to compensate the patient. There is some argument whether, if the drug were a proprietary preparation, the manufacturer would be responsible and not the physician, but none of these issues have yet been resolved. The present position is that if a patient suffers any damage from a properly prescribed drug he would only be likely to succeed in a claim for damages if he could prove that the product was of faulty manufacture—that is, some error had crept into the manufacture of one or more batches, causing the drug to be harmful.

The appearance of an allergy to a drug is not, under present

and is an "original," the remainder may be written (or typed) by a deputy—for example, a receptionist—or be a carbon copy of the original, except in the case of a controlled drug, when the following further restrictions apply:

— A prescription for the drug and the name and address of the patient as well as the doctor's signature must be in the doctor's own handwriting and be a copy (neither typewritten nor carbon copies are permitted for controlled drugs).

— The form of the preparation—for example, tablet, capsule, linctus, injection—must be written on the prescription and, where appropriate, the strength must be given in both words and figures; to too much the total quantity of the drug to be dispensed or the number of dosage units, as the case may be.

A prescription, whether NHS or private, is valid for six months (13 weeks in the case of a controlled drug) from the date the prescription was signed. Provided that the first dispensing of a repeat prescription was within the six months, the subsequent prescriptions may be dispensed at any time, though after a very long interval the chemist may want to check with the prescriber that his intention remains.

Repeat prescriptions

The role of the repeat prescription and its importance in the doctor-patient relationship is complex.¹ Many patients are on prolonged courses of treatment, sometimes indefinitely; once started on a remedy a few continue to take the same medication for the rest of their lives although the dose may vary with time and monitoring will be necessary—for example, thyroxine.

Providing patients with repeat prescriptions—that is, arranging for a further supply of a previously supplied medication without the patient having to see the doctor—is a universal custom: it is not just a British practice. Where the practice does differ it is that under the NHS the doctor's prescription enables the patient to get a medicine either free of charge or at what is usually a very reduced rate. Thus many items which would in other countries (or by private patients in Britain) be bought over the counter without a prescription—for example, antacids—will on the NHS be obtained on prescription because it is cheaper for the patient, though for those not exempt from prescription charges—presently £1—some items may now cost less if purchased over the counter.

Often the patient on a repeat prescription has an illness which though not lethal is incurable—for example, asthma, eczema. Changing treatment of control is rarely beneficial. Once such a regimen has been found, the patient can learn to manage his own illness. Thereafter he will need repeats of his medicine and since his doctor will have to supply any necessary prescription to enable him to get them, it is clear that some way of providing repeat prescriptions will be needed. For the patient with a chronic disease the problem is not whether the medication should be continuously prescribed but at what intervals the patient should be reassessed and how he is to obtain medication in the intervening period. Difficulties may, however, arise when the patient tries—and sometimes succeeds—in getting a prescription for a new symptom or a change in an existing symptom without actually seeing the doctor. The repeat prescription is now acting as an alternative to a proper medical consultation.

There are various reasons why a patient should continue to take a prescribed medicine:

- (1) Those having long-term prophylactic medication—such as an oral contraceptive. Once established on a suitable pill further examinations are only needed at 6- to 12-month intervals.
 - (2) Those having endocrine replacement treatment—for example, hypothyroid patients and diabetic patients on insulin.
 - (3) Those having other continuous long-term drug treatment—for example, epileptic and hypertensive patients.
- Some also include in the definition drugs prescribed for patients who are having intermittent but usually long-term treatment—for example, patients with dyspepsia and those who

suffer from migraine. Some extend the definition to include drugs prescribed for acute exacerbations of a chronic illness—for example, chronic bronchitis—and even for recurrent infections. I am sometimes phoned by a new patient who asks if I would leave a prescription on the shelf for an antibiotic "because the child has another attack of tonsillitis or otitis media." The parents seem surprised—but usually pleased—when I reply that I prefer to examine the child each time before prescribing a fresh course of antibiotics however often the child has had it in the past.

There are also patients who attend a specialist clinic at hospital, at regular but often infrequent intervals, who expect the general practitioner to prescribe the hospital-initiated drug in the intervals between outpatient appointments. Many hospital doctors make the same assumption, particularly since the recent cuts in hospital expenditure. It has long been the policy of the Department of Health that the doctor who decides the treatment should write the prescription. Furthermore, the doctor who writes the prescription must remain, at least in part, responsible for any harm that might befall the patient: a lot will depend on the acceptability or otherwise of such treatment. (So far as NHS prescribing costs are concerned, it is only a limited defence for a general practitioner whose prescribing costs are being questioned to say that what he prescribed was on the advice of a specialist.) If an NHS patient sees a specialist privately he may ask his general practitioner to rewrite the specialist's prescription on an NHS form. Provided that the general practitioner referred the patient to that specialist, it would be reasonable for the patient to have his drugs on the NHS so long as the general practitioner has vetted them for cost effectiveness as well as safety. It would not be reasonable for a patient to have a more expensive preparation on an NHS prescription merely because it was recommended by a private specialist. If the general practitioner is of the opinion that there is an alternative and cheaper drug that is equally effective.

Except for controlled drugs, the prescriber may indicate on a private prescription that the drug may be repeated once, several predetermined times, or as often as wanted either at specific intervals or as and when the patient wishes. Under the existing NHS regulations, this facility is not available. The FP10 is not only a prescription, it is also a voucher against which the pharmacist will be paid by the NHS for supplying the drug. Under the NHS, if a prescription is to be repeated the doctor has three alternatives:

- (1) The patient may have to see the doctor every time he wants a further prescription. This should normally only be necessary if a potentially dangerous drug—for example, a controlled drug—is being prescribed.
- (2) The doctor may issue a patient with an additional postdated prescription form for the same item, either carbon copies (except for his signature) or additional "top" copies. For example, a patient might see a specialist privately for 28 days' supply of a drug and two further prescription forms for the same drug and for the same period but dated 29 June and 27 July respectively.

A few practices confine their repeat prescriptions to these postdated copies: no patient can have a further supply without actually seeing the doctor. These doctors claim they exercise a greater control over their treatment and avoid patients concealing the development of new symptoms.

(3) In most practices arrangements are made by the doctor for his patients to call (by telephone, letter, or in person) at the surgery and collect (or have posted or delivered to them) a further prescription when the previous supply is running out. Many practices have introduced various forms of organisation that ease the burden of this work, regulate the flow, control possible abuse, and even perform some sort of audit.

There are a number of repeat prescription cards (produced either by individual practices or by pharmaceutical companies) in which the doctor writes the name of the drug(s) that may be repeated and the dose and the quantity to be given on any one time. Often, but by no means always, the number of times the

law, grounds for using a manufacturer unless he is aware of the risk and has failed to disclose it to those who prescribed the drug. Similarly a doctor who prescribed a drug to which a patient developed an allergy could also not be held liable unless it could be shown that he had been negligent by failing to take reasonable precautions. For example, he had not asked the patient before prescribing penicillin whether he had previously had any known reaction to that antigen or whether he had been identified during the early stage and hazards may not come to light until after the product has been in general clinical use for some time—for example, thalidomide and practolol. Whether the manufacturer

Unemployment in My Practice

Waterlooville

K B THOMAS

The area my practice is in has not had an unemployment problem over a long period of time, nor has it ever been a severely affected part of the country. Indeed, only in the past few months have we reached the national average of 10% of people out of work, and it would be wrong to suggest that the practice is obviously distressed by this trouble. There are, however, unmistakable signs that things are not quite as they were. The old confidence, the feeling that everything is possible, that everything is within reach, that each of us is the focus of a benign beneficial organisation has gone. The threat of unemployment lies like a pale shadow across the practice.

One of the first things I noticed was that some patients were reluctant to come off work. This is exemplified by the case of a gentleman who is much better known to me than I would wish. He used to burst into my room without knocking, with the words, "It's gone again, Doc." I always asked, "What's gone?" although I knew perfectly well. And he would reply, "My back." Then, after a negative examination, I would mention the subject of work. He would purse his lips and inhale a volume of air audibly, while half closing his eyes. His expression was a mixture of incredulity, sadness at the thought that I should expect him to work, and pain at the idea of actually performing it. A moment later he had taken my proffered certificate and departed with agility through the door—he was never interested in treatment. He came to see me recently, not because of his back but with a cough and cold. I was surprised; he even looked ill. I examined him and gave him a prescription. Then there was a pause as I waited for him to ask for a certificate. He was looking at me with a faint wistful smile that was almost apologetic, regretful that he could not fulfil his traditional role. I felt a little sad, and perhaps this showed in my face. I think the episode is what my colleagues of the left would call non-verbal

communication. At all events we understood one another. For so long I had been able to supply his needs, and now the rules of the game had been changed.

These are bad times

Often as an undercurrent or as background one learns that the patient or his wife or her husband is unemployed. I was told by one wife how restless and irritable her husband has become since he has been at home; by a teenager that she is quarrelling with her parents because "I cannot go out now; I've no money." Some young people get into more serious financial difficulties. In one family the husband, who had been unemployed for three months, could not face the mounting bills and attempted to end his life. Yet most people I talk to are philosophical: "These are bad times," they say, "and I am not the only one."

What disturbs me most as I roam round the practice is to find, perhaps in the afternoon, a young man sitting watching television. "Hello, what are you doing at home?" I ask, and back comes the answer, "I am out of work." Thinking and feeling not as an economist but as a doctor who has been in the practice so long that I might well have known this man all his life, I find it so absurd that he is being paid for doing nothing that I can hardly believe that it is true. He would be better off being something with other people digging holes and filling them in again. He would be better off physically, physiologically, and socially—to borrow a phrase from the Royal College of General Practitioners.

Work is such an important part of our lives that if there is to be no work for some people, and I fear that full employment may never come back, then work must be created, perhaps second-class work at a lower rate of pay. To have a large section of the community doing nothing for a long time is a certain recipe for disaster, disaster for the individual and perhaps disaster for society. A patient told me that her husband, who was epileptic and had been unemployed for a long time, had attacked her. "You see," she said, "he has nothing to do all day now. He used to have to go once a week to draw his money, and this took him out of himself, but now they send it to him by post."

Waterlooville, Portsmouth, PO7 6AH
K B THOMAS, *MD, general practitioner*