PRACTICE OBSERVED

Emergencies in the Home

ENT emergencies in general practice

H DE LA HAYE DAVIES

The amount a general practitioner can do in managing ear, nose, and throat (ENT) emergencies depends on his experience and the availability of the right equipment. Ideally if he has experience in an ENT house job or as a clinical assistant then he will be expert in using the head mirror, a technique which requires some practice and which leaves his hands free to carry out various manipulations with his instruments. These manipulations are performed in dark, deep cavities or sensitive areas in patients who are usually frightened and liable to move at the slightest touch. Good illumination is therefore essential and, if one is not adept at using a head mirror, an electrically operated head lamp can provide a good source of illumination and at the same time leave both hands free. Most family doctors, however, are equipped only with the ordinary operating otoscope, which can if used properly provide quite good illumination for looking down ears and can also double up as a nasal speculum (in a child a perfect view can be obtained if the tip of the nose is just lifted and there is no necessity to insert a speculum). The light can also be used to visualise the pharynx and—with special attachments—the postnasal space and the larynx.

The experienced ENT doctor will probably also have his surgery equipped with Jobson Horne probes, wax hooks, aural forceps, nasal speculums, etc, which will enable him to carry out some of the procedures normally carried out in the ENT department. Whether two or expert, however, the golden rule is never to attempt anything unless you have not only the correct equipment but also the experience and confidence that you will succeed. A patient who is anxious about the ENT department nervous and frightened after an attempt by a well-meaning amateur is more difficult for the houseman or registrar to deal with and may even need to be given a general anaesthetic for a simple procedure which, if seen untouched on arrival at the department, could well be done without any anaesthesia at all.

Department of Otolaryngology, Northampton General Hospital
H DE LA HAYE DAVIES, BA, PhD, clinical assistant and general practitioner

If general practitioners are going to follow this course of action, however, the local hospital must in turn provide a good service to the family doctor. In addition to the routine outpatient clinics a daily treatment clinic should be held by one of the senior house officers so that not only may new cases be seen quickly but the correct daily treatment carried out. So often ENT treatment fails in practice, for example, because of failure to remove debris before antibiotic drops are inserted.

Having been a clinical assistant for over 20 years in a busy department and also having dealt with emergencies in my general practice I have the advantage of seeing both sides of the fence, and my only criticism of fellow practitioners is that they seem reluctant to use the emergency services of the department and are often incorrect in assessing what may be dealt with at home and what should go to hospital. In some areas the facilities provided by the hospital may not be as good as in my own area, and in such cases it is up to the general practitioners to put pressure on the hospital authorities to provide a proper service.

The ears

Earache is perhaps the most painful emergency to which one may be summoned, especially in a small child; most of us are used to the mother telephoning for help, usually in the small hours of the morning, with a screaming child heard clearly in the background. Most of these cases, particularly of sudden onset, are due to middle-ear disease, and a heavy dose of analgesic is the first priority. I have no hesitation in giving an adult dose of Dypirin or paracetamol to a child in such circumstances or in leaving instructions that the dose should be repeated until the pain stops. I have found Auralgan ear drops to be the best pain-relieving drops, and the child should be started on penicillin (double the normal dose for the first 24 hours) given orally as I try to avoid giving injections to children. It may be argued that if adequate analgesia has been achieved by tablets it complicates matters to fill the ear with "treacle." In practice,

10 weeks whatever treatment is used, and permanent paralysis is rare. In its early stages, however, the condition is distressing to the patient, and for this reason I think one must do something positive. My own favourite technique is to prescribe a short, sharp course of steroids starting off with a heavy dose—for example, two 5-mg prednisolone tablets four times daily, reducing over a week to 5 mg twice daily. Referral to hospital for a specialist opinion should follow so that in rare cases of non-recovery the patient feels that the family doctor has done all that he could.

Nose

EPISTAXIS

Epistaxis can be an alarming condition yet in many cases the bleeding can be arrested with simple first-aid measures if timed off properly by the patient. The bleeding area is in most cases from a prominent blood vessel in Little's area, which is at the tip of the nasal septum and can easily be compressed if pressure is exerted by the thumb and forefinger of the right hand applied firmly over the soft tip of the patient's nose. This pressure should be maintained for at least five minutes. Plugging the nose should be avoided as this opens more blood vessels and when the plug is pulled out the bleeding starts again as the dried blood clot is pulled away. If simple first-aid measures do not stop the bleeding then the patient should be dealt with in hospital, especially if the bleeding is arising from further back in the nose.

TRAUMA AND FOREIGN BODIES

Fracture of the nasal bones usually requires reduction in hospital under an anaesthetic within 10 days of the trauma, when the swelling has subsided, but if the doctor is available immediately after the trauma while the nose is still numb a firm push with the thumb on the side of the deformed bone will often result in a satisfying click being heard and felt and no further reduction will be necessary. A case of foreign body inside the nose should also be referred to hospital without interference so that the object may be removed by an experienced operator using the proper equipment and lighting.

Throat and oropharynx

TONSILLITIS

The most common emergency that one meets in general practice is tonsillitis, and in smaller children this may be because of the accompanying high temperature, make the child appear very ill (especially to the relatives) and also often results in infantile convulsions. It is my clinical impression that most cases of infantile convulsions due to hospital for investigation can be sure that an infantile convulsion is due to tonsillitis then I tend to treat the condition at home, but if I am not sure of the diagnosis then it is my general rule to admit all "first time" cases of infantile convulsions to hospital for investigation by paediatricians. Although most cases of tonsillitis respond to oral penicillin, repeated attacks lead to increasing parental anxiety (especially when both parents are out at work) and an increasing number of out-of-hours visits are requested, which are all part of the gentle pressure being exerted for the child's tonsils to be removed. This article is not the place to discuss the pros and cons of tonsillectomy, but these families may often be reassured if left with a starter dose of penicillin to be given at an early stage during an attack. If the general practitioner is clever and plays for time until the child develops his or her own immunity it may be possible to save an unnecessary tonsillectomy.

RESPIRATORY OBSTRUCTION

Perhaps the most dramatic ENT emergency and sometimes one that requires courageous action on the part of the attending doctor is obstruction due to swelling of the tissues of the throat. Doctors who qualified during the preantibiotic era, when diphtheria was common in general practice, had no hesitation in performing a tracheotomy when the signs of laryngeal or pharyngeal obstruction manifested themselves. Most of us who qualified after the war have never had to do an emergency tracheotomy under such circumstances and indeed would be reluctant to perform such a procedure until the patient was almost "in extremis." I have been told that as an emergency first-aid measure a large-bore intravenous needle or chest aspiration needle inserted into the cricothyroid membrane would certainly give a child, and possibly an adult, enough of an airway to at least reach hospital. I have never had to perform such an operation in practice but I do have a large-bore needle in my road accident bag (primarily intended for the relief of a tension pneumothorax). Swelling of the tissues of the throat commonly arise owing to allergic conditions—for example, bee sting, drug reactions—and intravenous chlorpheniramine maleate (Pirnon) may be supplemented by intravenous or intramuscular injections of 100 mg hydrocortisone and also subcutaneous injection of 1 in 1000 adrenaline (up to 0.5 ml as a single dose followed by slow injection at a rate of 0.1 ml per minute). The swelling of the tissues of the throat can also be caused by acute infections, and here is a definite indication for intramuscular injection of antibiotic, penicillin being the first choice.

EPIGLOTTITIS

Epiglottitis that occurs in young children and is sometimes mistaken for croup is a desperate emergency and requires immediate admission to hospital. The epiglottitis swells as a

result of infection due to *Haemophilus influenzae* and this results in laryngeal obstruction, which is also accompanied by laryngeal spasm. The condition is all the more hair-raising for the doctor because the very act of putting a spatula on the tongue can precipitate an acute attack. I have never seen an attack in my general practice but I have been at the hospital receiving end and witnessed more expert colleagues dealing with what is a serious life-threatening condition. The only advice I can give is that one should not forget the possibility of such a diagnosis in any obviously ill and toxic child with symptoms of croup, whose illness started off as a sore throat or upper respiratory infection and developed its present symptoms rather rapidly. One should do no more than a purely visual examination of the back of the throat, and no instruments should be introduced into the mouth. In fact some experts claim even opening the jaws wide can precipitate an attack, so it is probably justified to order the patient to hospital immediately on suspicion alone.

Do what you can but no more

In the final analysis, the successful treatment of ENT emergencies depends on good communication with your local hospital and mutual understanding by the staff of the ENT department and the family doctors of each other's problems, but most important of all I reiterate the advice given in the first part of this article that no doctor should tackle an emergency problem unless he is sure that he is competent to do the job that he has the right equipment and surroundings to do this properly, otherwise removal to hospital should be the rule rather than the exception so the patient is in the right place at the right time.

I am grateful to Mr Keith Thomas, ENT consultant, Northampton General Hospital, for advice in preparing this article and to Mrs Daphne Brennan, secretary to the department, for secretarial help.

Patient Participation

Aberdare Patients Committee

D J THOMAS

What is a patients committee?

Our patients' committee is an elected body whose sole object is to build up a close relationship between doctor and patient. Medically a close relationship already exists between general practitioners and their patients. Our committee highlights the areas where a closer relationship can exist mainly by means of lectures on primary care and films.

The first committee was set up in 1973 at the request of a general practitioner and patients at the Aberdare Health Centre. It set about making more democratic the relationship between the general practitioners and the patients. The doctors were

Aberdare Health Centre, Mid-Glamorgan

D J THOMAS, chairman

quick to respond and gave the fullest co-operation from the outset. Without that co-operation there cannot be an effective patients committee, for the theme is participation.

Who are the members?

The committee meets once a month. We are 26 members from a patient list of 10 000. In addition there are subcommittees for social activities, lectures, care of the elderly, etc, that meet more often. Committee members are elected at the annual general meeting, which is well advertised in the local press and at the health centre, and an average of 75 people attend. The members of the committee cover a broad spectrum—I am a community health council member, as are two others; there is a person from the old age pensioners group; miners; factory workers; a local editor; teachers; an accountant; county and

local councillors; a nurse; a housewife; and general practitioners. During the past year we have held 11 committee meetings, with an average attendance of 22. All sign an attendance register.

What does the committee do?

We have held 10 public lectures in the past year (over 100 in all on health education, attended by an average of 30 people, some of whom are not patients of the health centre. We have had socials and films and visited hospitals and other Health Service premises. We have entertained 14 doctors and social workers, some from abroad, at meetings. Our last AGM was filmed by the Welsh Arts Council, and a teaching film was made by two patients and a doctor which has been shown in many places. We were also filmed for Grapevine on TV.

We helped to get a psychiatric clinic at the health centre. We run a stroke club, a caring group, a well-woman clinic (committee of three women and a lady GP), and are concerned about cervical screening and screening for the elderly, perinatal mortality, mental health, and the disabled.

We are concerned about the patients who work in collieries and factories and the effect of industrial disease. We make our views known rather forcibly on such matters as ENT and general waiting list delays. We co-operate closely with the community health council and the Age Action Committee and supported a seminar on the problems of the elderly in our community. We work closely with the local housing association, who provide sheltered accommodation and special flats for people who have recovered from mental illness.

Law and the General Practitioner

Committing a patient

S E JOSSE

The rules for committing mentally ill patients to hospital are over 20 years old now and are contained in the Mental Health Act of 1959. Sections 25 to 29 of the Act are particularly important in relation to "admission for treatment."

Section 25 is otherwise known as "admission for observation" and requires medical evidence from two doctors. The period of detention is up to but must not exceed 28 days.

Section 26 is referred to as "admission for treatment" and again requires two doctors independently to provide the medical grounds for admission. The period of detention may be up to one year initially.

In emergency cases one doctor can provide the medical recommendation for admission under Section 29, but the authorised period is reduced to 72 hours from when the patient is admitted.

There are other sections concerned with the compulsory

What are the benefits of a patients committee? Members of the committee take the chair at lectures and group meetings and feel that they are helping to promote good health in the community.

The doctor is told about difficulties patients have that only the nurse or receptionist is aware of—such as patients having to wait for a long time to see the doctor; repeat prescriptions of drugs; receptionists taking decisions on matters that are outside their jurisdiction.

The psychological barrier between the doctor and the patient is broken down. The GP must retain his professional status but not allow himself to be above it all. The ideal GP is a friend to the patient.

A more enlightened patient understands the doctor's problems and can be more tolerant, and the GP has a better understanding of the patient's problems and can relate to his patients.

Is a patients committee worth while?

As a member of many volunteer groups in the community, I believe a patient participation committee is the most interesting, instructive, and satisfying of them all. You cannot possibly know what you are missing by not having one. If only all GPs could appreciate the value of a patient participation group in their practices it would go a long way to making for more efficient and satisfying practice—the patient satisfied that he is getting the best service and the GP happy knowing that there is a better understanding between them both.

detention of mentally ill patients, but these are of little or no relevance to the general practitioner.

Criteria for commitment

Certain key criteria must be fulfilled before a patient can be committed: the patient must be suffering from a mental disorder as defined in the Act; the extent of the disorder must warrant detention; the patient does not agree to be admitted to hospital voluntarily; either the patient's own health or safety must be at risk or else other people must require protection from the patient; there must be somebody to make an application for admission; and this application must be supported by either one or two registered medical practitioners.

MENTAL DISORDER

In general terms the Act (Section 4) defines mental disorder as mental illness, arrested or incomplete development of mind, psychopathic disorder, and any other disorder or disability of mind. In the next few subparagraphs of this section, however,

however, the "instant analgesia" provided by drops complements the prolonged analgesia obtained by oral medication, which should err on the side of overdosage rather than underdosage. In most cases of middle-ear infection I prefer to carry on the antibiotic for 10 days instead of five days, which is normal for other upper respiratory tract infections. I certainly continue the antibiotic until there is no increase in temperature, the tympanic membrane appears normal, and any discharge has stopped. Later the child should be followed up to exclude glue ear, which can result in deafness. Some children who are prone to recurrent ear infections may well benefit from prolonged courses of anti-histamine agents when they have a cold or appear to be deaf—for example, Actifed Elixir 5 ml twice daily. In these children adenoids should be excluded as a cause of infection and also the possibility of the ears being infected from recurrent sinus infections.

A perforation may well result from an acute middle-ear infection and this should be observed until it is healed, and if a perforation has not healed in six months the child should be referred for a specialist opinion. In an older child a repair may be considered provided the ear is not chronically discharging; otherwise the graft will break down. The important thing about chronic discharging ears is a correct assessment, and the patient should be referred to a specialist if the discharge does not dry up.

I find that mothers with children who suffer repeated upper respiratory tract infections, particularly earaches, are usually sensible and become expert in forecasting when trouble is liable to erupt, and I have no hesitation in leaving such mothers an emergency starter dose of penicillin which they can make up themselves and give after a telephone consultation with me. I know this is considered lazy doctoring by the purists, but it works, and I have a sinking suspicion that many of my colleagues believe, like me, that prophylactic penicillin and antibiotics protect many ears, save a lot of unnecessary pain, and let us sleep a little longer.

EXTERNAL OTITIS

External otitis can also be an emergency condition, especially when the inflammation results in oedematous swelling which, because of the bony cartilaginous wall of the external canal, results in the epithelium being pushed inwards, narrowing the canal and causing pain for the patient. Secondary infection may give rise to a furuncle and infection can spread back causing perichondritis especially if the furuncle is incised. A swab for bacteriological culture should be taken before treatment starts and in addition to systemic antibiotics local treatment of the painful cause of a furuncle or a grossly narrowed meatus should be started by inserting ichthammol glycerin wick. The ichthammol is bacteriostatic, the glycerin is hygroscopic, and this old-fashioned treatment not only opens the canal but is also very soothing to the patient.

Once the acute painful stage of external otitis has been dealt with it is important to treat the underlying condition. This requires regular daily toilet under direct vision and there may be necessary to refer the patient to an ENT department because it is no use putting in local antibiotic drops if the ear is filled with pus or debris. Prolonged treatment with antibiotic drops may also result in secondary fungal infection, and indeed sometimes patients can be sensitised to the antibiotics in the drops. Proper toilet and keen vigilance are the cornerstones of proper treatment. It is important to assess early on whether the external otitis is due to secondary infection from a middle-ear infection or whether it is associated with a dermatological cause.

Bell's palsy

The most common form of paralysis of the facial nerve is Bell's palsy and fortunately 85% of patients recover within

Bristol Medical Centre, London N11 2BD

S E JOSSE, FRCS, DPM, general practitioner and police surgeon

The Act goes on to define "severe subnormality," "subnormality," and "psychopathic disorder." Severe subnormality means that a patient is incapable of leading an independent life or guarding himself against serious exploitation. A subnormal patient is one whose condition is less severe but who requires treatment or care. Psychopathic disorder in a patient means that he persistently displays abnormally aggressive or seriously irresponsible behaviour and requires medical treatment.

The Act makes it clear that promiscuity or other immoral conduct on their own are not grounds for compulsory detention. Though not mentioned in the Act, drug addiction is not if causing severe physical effects is not a reason for commitment.

THE EXTENT OF THE MENTAL DISORDER

Apart from the specific definitions given above mental illness is usually regarded as meaning psychosis. The effects of drugs including alcohol or certain physical disorders—for example, hypoglycaemia—may, however, temporarily produce a clinical picture similar to a psychosis, and it may be difficult to distinguish one from the other.

CONSENT FOR ADMISSION

If the patient voluntarily agrees to enter hospital then this is all that may be required. It may be known, however, that a particular patient discharges himself from hospital after being admitted voluntarily, and if this has occurred several times, then it may be assumed that the patient has no real wish to enter hospital as a voluntary patient. Under these circumstances compulsory admission may be the only option. Clearly where a patient is incapable of giving consent due to his mental condition, then hospital detention can only legally be effected through the Act.

THE PATIENT'S HEALTH OR SAFETY, AND OTHER PERSON'S PROTECTION

An actual or implied threat of suicide, attempted suicide, or self injury (including care neglect), or some action against another are the other circumstances in which compulsory admission can be argued. These must, however, arise from a mental disorder not criminal behaviour; the patient must be "mad not bad."

THE APPLICANT

There has to be an actual applicant who has seen the patient within the previous 14 days (the threshold for Section 29) in order to start the admission procedure. This means that doctors cannot act on their own by solely providing the medical opinions, unless the patient is already in hospital voluntarily. An applicant is defined in Section 27 of the Act as either the nearest relative of the patient or a mental welfare officer appointed for the purpose by a local authority. If the admission needs to be arranged urgently then any relative will do. Section 49 of the Act carefully defines who is a relative, what the nearest of that relationship has to be, and what modifying circumstances (such as the nearest relative being abroad) may apply.

It is clear that the mental welfare officer does not have to be a party to an admission. A relative, however, may not wish to be responsible for such an act. Also most general practitioners do not carry the appropriate forms without which admission cannot be authorised. Apart from providing the documents the mental welfare officer will usually arrange the admission, and may even physically convey the patient to hospital as he is authorised to do under Section 31.

The mental welfare officer is not, however, an entirely free agent. If the nearest relative objects to the application being made, the mental welfare officer has to apply to a court for permission to continue with the procedure (Section 52). Furthermore, the mental welfare officer has to seek out and consult with the nearest relative before making an application unless this would result in undue delay or is not reasonably practicable.

At times the patient may have locked himself in his house making it impossible for the applicant or the doctor to assess him. Under these circumstances a mental welfare officer under oath to a justice of the peace can declare that he believes that the mentally disordered patient is neglected, ill treated, or unable to look after himself. The justice may then issue a warrant authorising a constable (accompanied by a mental welfare officer and a doctor, who may be the patient's general practitioner) to enter the patient's house and move him to a safe place. He can then be committed if it is thought appropriate (Section 135).

MEDICAL RECOMMENDATION

As I have said, the application for admission has to be supported by two written recommendations by registered medical practitioners under Sections 25 and 26, and by only one under Section 29 (for emergency admission). One doctor should have had previous acquaintance with the patient, and clearly this may well be the general practitioner. It is not a legal requirement, however, that either of the two doctors must have known the patient previously—even for Section 29. The second doctor providing the medical recommendation, whether he previously knows the patient or not, must have been approved by a health authority under Section 28 of the Act as being a practitioner with special experience of mental disorder. Psychiatrists normally are appointed but doctors in general practice, such as police surgeons, have also been approved under this section.

Under Sections 25 and 26 the two doctors must have personally examined the patient (as must the single doctor in Section 29) but not necessarily together. If the examinations are done separately the interval between them must not exceed seven days. Section 28 details certain relationships between the two doctors or between a doctor and the patient or applicant which prohibit either one or both doctors from providing the medical recommendations. These include: the two doctors being in partnership together or one the assistant to the other; where a doctor also acts as the applicant or is a partner of the applicant; where a doctor has certain financial interests in the maintenance of the patient; and where certain blood or marriage relationships exist with the patient or the other doctor.

Since Section 26 allows compulsory admission initially for a period up to one year, the medical grounds have to be stated in writing by both doctors in some detail. Although the two statements do not have to be identical, the diagnosis of the mental disorder (or disorders) must, in part at least, be identical. Any recommendation must not give as the sole grounds for admission the fact that the patient was suffering from a psychopathic disorder or subnormality and was not less than 21 years of age.

Disposal

Once the application for admission has been correctly completed, a patient may be taken to a hospital even against his will. A hospital must be found, however, that is willing to take the patient. In other words, "certifying" a patient lays no legal obligation on a hospital to accept him.

Again, under Sections 25, 26, or 29 no specific duty is laid on the police to aid the transfer of a violent or struggling patient

Beyond the Surgery

General practitioner on the lifeboat

O C PARRY-JONES

Little did I think some 20 years ago when I started in general practice that I should be in active service for more than a decade as a lifeboat doctor or HMA (honorary medical adviser), as we are called. Years later I asked the now retired mechanic, who was out with me, about my first trip. Most of the crew at that time were the old "Dick Evans," double Gold Medal crew of the Moelfre Lifeboat. The trip had been what has become more commonplace for me since: transferring in a near gale from the lifeboat to a Jacob's ladder on the cargo vessel that was four or five miles offshore to examine the patient before he was brought ashore. I asked Ian Owen, "What would have happened if I had fallen off that night? Would you have found me? Would you have been able to see me?" "Not at all," he said. And of course after that I went through a period of wondering whether it was a good idea for a family man to continue as a lifeboat doctor, and I almost resigned.

Since then I have become more philosophical and consider it no more than a part of my normal practice duties that extend a few miles offshore, and that is what it is. Having the opportunity of being a lifeboat doctor depends very much on your choice of location for your general practice, and living by the sea one has a special responsibility. I have been very fortunate in always having a crew in whose hands I felt completely safe at all times and a coxswain on whom I could rely, so that when he said "jump," I jumped. Normally the HMA is required only in fairly good weather, and a force seven or eight gale would be about the worst, with a bit of luck.

As chairman of the lifeboat station committee I am involved in local lifeboat politics and running the station. I teach the crew first aid, examine new crew members, and re-examine the present crew for three-monthly regular intervals. A medical newsletter is circulated by the Royal National Lifeboat Institution (RNLI), edited by Dr Geoffrey Hale, and there are regular meetings at the BASICS (British Association for Immediate Care Schemes) annual conference at which the chairman of the Medical and Survival Committee, Professor Sir George Smart, is present. The 200 HMAs provide an immediate care scheme as part of BASICS.

Merchant ships and swimmers

Our lifeboat station is near the shipping lanes to and from Liverpool, and Moelfre Bay is one of the best places to shelter from a gale on the west coast. Calls come at any time of the year, but there is a difference between those in summer and winter. In winter the medical calls are specifically to merchant

Benlloch, Anglesey, Gwynedd
O C PARRY-JONES, Llanfyllter, IMA, general practitioner

The Moelfre lifeboat, RNLI Robert Clarke.

ships in the bay, while in summer most work is in rescuing small boats and swimmers in distress, either using the big boat or more often the small inflatable inshore lifeboat with an outboard motor to rescue children who have floated out on mattresses, for instance. These are the rescues that bring in the victims of drowning or apparent drowning. Tragically, these are often holiday-makers or day-trippers. I like to make a link with the relations of the victims, possible for counselling.

The medical calls can be as varied as those in ordinary general practice. Often the crisis is that the ship is bound for foreign waters and that any time wasted ashore or anchored in the bay is time lost by the committee, therefore the sick person has to be got ashore as quickly as possible. The person may be suffering from a coronary with chest pains, abdominal pain precipitated by the celebrations of the last night ashore, an accident on a tanker or fishing vessel, and even a gynaecological condition such as an abortion in the wife of a crew member. In these cases the lifeboat may act just like an ambulance for what is called a medical evacuation, but often the person has to be examined while aboard and the condition stabilised before taking him ashore. It is therefore wise for the lifeboat HMA to have a fair idea of what type of cases he might be called to treat and to carry a special bag at all times with fresh supplies of equipment such as pain killers, intravenous fluids, drugs and antibiotics, antiseptics, not forgetting insulin, glucagon, and an antiarrhythmic.

Our boat is among the few in the fleet that carry Entonox (nitrous oxide), and I think that I was the first to use this in the inshore lifeboat years ago on a schoolboy who fractured a tibia in a cliff fall. A modern lifeboat, unless it is one of the large ones, is just not equipped to treat sick persons on board. Our particular lifeboat, the Rother, is 37 ft (11.4 m) long and has accommodation if necessary for 70 survivors, but in the forward cabin where I usually work there is room at most for

to hospital, but usually the police are helpful as are ambulance crews.

A doctor may consider giving some tranquillising drug in these circumstances, and if the patient consents to the treatment no difficulty arises. From the very nature of the problem, however, the patient is usually unable or unwilling to consent. Might a doctor lay himself open to a legal action by forcibly treating the patient against his will by, say, giving him an injection of chlorpromazine? If the patient was being admitted under Section 26 the words "for treatment" are actually stated in the section and so no difficulty should arise. Sections 25 and 29 do not contain such a phrase, however, and some argue that there is no specific legal authority for the doctor to treat patients under these sections against their will. All that is allowed is the authority to transfer the patients to hospital and observe them. Treatment may then only be given compulsorily if the section is converted into a Section 26.

Parliament presumably intended, however, that having authorised a patient's transfer to hospital against his will if necessary, the doctor should be able to ensure that this happens peacefully and should be able to give suitable drugs. Similarly permission to use drugs to quieten a restless patient so as to prevent personal injury is perhaps also implied in Section 25.

Clinical Curio: diligence rewarded

The temptation to stop thinking about a diagnostic problem is sometimes irresistible, especially when a press of patients is outside the consulting room door. Even in the more leisurely atmosphere of the ward a spate of negative investigations may lure the clinician into a forest of false assumptions. At least one of our patients owes his survival to an assiduous medical registrar who refused to give up.

The patient, a bachelor lecturing in metallurgy, was then aged 51 and had suffered intermittent fever for three years or more. He was cachectic, normally demented, and had increasing bowel symptoms and a severe peripheral neuropathy. What misled his own general practitioner and the first referral specialist was his history. At the age of 26 he had undergone an operation for sarcoma of the tonsil. He subsequently had a succession of sore throats and respiratory illnesses that seemed to culminate in the febrile episodes for which he was eventually referred to hospital. Apart from an erythrocyte sedimentation rate of 13 mm in the first hour, the results of all tests were negative. His liver and spleen were palpable, but a liver biopsy showed no abnormality. The patient remained in hospital for five weeks and was thoroughly investigated (sinus). "The diagnostic favourites were some form of reticulosis, brucellosis, or perhaps sarcoidosis," wrote the registrar at the first hospital. No confirmatory evidence was forthcoming.

In March 1966 the patient was referred to a teaching hospital under one of London's most scintillating diagnosticians. Gall stones were found, and a lymphangiogram showed an enlarged group of para-aortic glands and possibly glands in the groin. A biopsy of one of these showed no abnormality. Lymphosarcoma or other tumour was suspected, especially as the sections from the original tonsil operation were proved to be reticulum-cell sarcoma. Laparotomy was advised, but the patient was unwilling, a viewpoint with which the first consultant concurred. Six months later, despite a general remission, the patient was suffering odd aches in the stomach and bouts of vomiting. A barium-meal examination suggested partial obstruction of the small bowel. The report referred to an abnormal mucosal pattern. The patient now agreed to laparotomy, which confirmed the enlarged para-aortic and mesenteric lymph-glands, but no reticulosis was found. Over the next few months his abdominal symptoms continued, there was a pronounced deterioration in his mental state, and he was asked to give up his job. In May I received a call from his mother, with whom he lived, saying that he was worried about his behaviour. I was met at the door by the patient, who took me into a side room to tell me in a confidential whisper that his mother had gone out of her mind. I listened sympathetically,

These views have never been challenged in the courts but the specific absence of such authorisation in the Act is worrying.

Section 136 or a mad MP

Occasionally a general practitioner may be contacted for information by the police when a constable has removed one of his patients to a place of safety (which may be a police station) under Section 136. The patient must have been in a public place or a place to which the public have access (not in his own home in other words) for this action to take place, and the period of detention can be up to 72 hours.

For those general practitioners who have patients who are Members of Parliament it is interesting to note that the MPs' authority admirably provides the special commission mentioned in the Act (Section 137). It is complicated by the fact that on detention the MP has to be examined by, in addition to the usual two doctors, other specially appointed medical practitioners who have to report back to the Speaker. In due course his seat may be declared vacant. Such provisions do not exist for members of the House of Lords, who, of course, have no seats to vacate.

half-persuaded, and then interviewed his mother, who told me that her son had tried to drive the car out of the garage without first opening the door. The patient was admitted to a mental hospital. After discharge his physical condition deteriorated considerably with considerable sensory loss, imbalance, and pronounced ataxic gait, and he was presumed to be nearing end of his reticulosis. He was, however, mentally alert enough to ask if he could have yet another specialist opinion. I agreed with reluctance, fearing to waste further time and resources. After an initial outpatient interview he was put onto prednisone and admitted to another teaching hospital, where extensive investigations were again carried out. The erythrocyte sedimentation rate was still raised, but little else was found except coarse folds in the small-bowel mucosa and the old gall stones. An attempt at jejunal biopsy failed. He was discharged with a label of malabsorption syndrome (causae). Vitamin treatment was added to the steroids.

By good grace the medical registrar had not given up and decided to repeat his attempt at jejunal biopsy, and this was successful. The report mentioned large numbers of phagocytes with foamy granular cytoplasm containing para-aminoazobenzene-positive material in the submucosa, strongly suggestive of Whipple's disease. The saga was not yet over; the house officer in the follow-up clinic had the temerity to write, "This will not affect our management of the case." The registrar meanwhile had been reading up on the disease and it seemed that the condition was a strange infectious disease of the bacteria in the gut, and "cures" of up to eight years had been recorded after prolonged treatment with tetracycline. Within a few weeks of starting tetracycline the patient was much better, although it was many months before his strength had substantially returned, the symptoms of peripheral neuropathy being the slowest to recover. After 18 months he had resumed a reasonably normal life and consults only for relatively trivial conditions unrelated to the Whipple's disease. A fortnight as the registrar was writing the Whipple's disease I came upon a letter from a tired locum specialist who wrote in 1967, "I had great difficulty tracing a useful history from the patient, who seemed to be in a convalescent stage of the disease." No comment.—MYRICK EMMYS-ROBERTS, general practitioner, Walton on Thames, Surrey.

We shall be pleased to consider for publication other interesting clinical observations in general practice.—Ed, BMJ.

only two stretcher cases in very cramped conditions. The stretcher is invariably the most versatile of all—the Neil Robertson model. We can sometimes sling another one or two on the deck. In fact last year we treated a cardiac arrest on deck on top of the cabin for the three-hour journey from the Liverpool-to-Dublin car ferry back to the slipway. He was one of the rare cases that was a fake. He was a real case of Münchhausen syndrome, having simulated cardiac arrest on the car ferry, fooling three medical members of the intensive care team of Jervis Street Hospital, Dublin. He also took me in. I took an electrocardiogram in the ambulance by the boat house but it was not until the next morning after he had been in hospital several hours that he was shown to be a malingerer. That is a very unusual story, however. Most of our cases are quite genuine even if they do sometimes walk ashore undetected.

Sometimes Anglesey Radio, the General Post Office radio station, will ask me to speak to the skipper of a merchant ship maybe 100 miles away who is approaching our area asking for medical advice. This past autumn I treated the captain of a small Navy minesweeper who was brought ashore by his own boat to my local beach 200 yards from my surgery. The Post Office radio station is apparently directed to ask for medical advice from the nearest hospital. This sometimes causes difficulties, and we have to attend to the patient at the hospital asking the skipper to bring the patient ashore so that he can be seen in outpatients next morning. When a vessel approaches our patch, Anglesey Radio has agreed to ask me for advice. Being the doctor most rewarding work and the most in evidence there may not be the slightest bit of difference from one's council estate patients ringing up during surgery time for advice, for this is the responsibility one has in general practice—to provide primary care.

Rescues

Rescues are entirely in the charge of the area coastguard rescue headquarters, in our case Holyhead. They may ask for the assistance of a helicopter from the RAF Search and Rescue Squadron 22 at Valley. Once the coastguard think that it is necessary to launch the lifeboat, he or she (we have lady coastguards) will inform the honorary secretary of the lifeboat station who will authorise launching. I will have been given the relevant information already and possibly discussed the patient with the ship's captain. If I am in the surgery it will take me only a few minutes to go down to the lifeboat station; otherwise I can be informed by radio in my car. It will then take only a

few minutes to slip on my Javelin boots and an extra jumper and, putting my flasher on top of my car and sometimes using my siren, judiciously, to approach the boat house. I have my personal RNLI protective clothing and air jacket hanging in the boat house and wear crash-helmet-type headgear. I always receive every courtesy from the crew, and the head launcher or one of the shore crew, who are such a very essential part of any lifeboat station, take charge of my ignition key until the boat is retrieved.

If the weather is easier it can be difficult, and until our new breaker is built we have to shelter in the Menai Straits and are off station for some hours.

The coastguard will have also informed the ambulance station to stand by and an ambulance will be waiting for us when we return to the slipway. I always find the launch down the slipway particularly exciting, and I am quite a steep one of the steepest in the country. I think everybody who does emergency work secretly enjoys that little extra perk of the thrill of the job—in my case to be involved in an exhilarating launch. Once at sea everyone knows his place and I just keep out of everybody's way. I think in this respect that there is some similarity between the lifeboat HMA and the ship's surgeon in the Royal Navy. It has been said, "Like the engineer, the surgeon had a long battle before his status as a naval officer was recognised."

The view of the sea surgeon, which—unchanging as the sea—still holds today, is a blend of acceptance and rejection, of a claim to treat the methods while, conceding the surgeon's services. Dismissed as a man taking no part in the life on the ship, he is accorded the guarded welcome of the stranger within the gates. Once one gets to know the crew, however, and has the good nature to haul on a line here and there and keep out of the way during launching and retrieval, one is treated with the greatest of good nature and one has great affection for one's boat.

RNLI

The RNLI, as everybody knows, is a voluntary organisation that depends on charity. It is also a Christian organisation and new lifeboats are dedicated and blessed by the archbishop and representatives of other denominations and faiths. This gives the crew and their families great comfort. Last year in Moelfre it was our 150th anniversary, and we were honoured by the presence of His Grace the Archbishop of Wales at the ceremony.

"In the end we lived and slept in the big family bed in the room above the surgery, in the front of the house, which did not come to bed. He would lie for a while at the end of the bed with fluffy clothes and then go out again on his call. In his car he had Winchester's mistal and mist expect and big bottles of aspirin to give out in the street to groups of colliers staggering home from work with their arms round each other to support themselves. He told me that of six miners coming home in tux, two would probably die within 24 hours of the epidemic pneumonia. Our family were kept alive on Welsh broth (cawl) by William Henry's sister, Miss Davies.

The old whelwright and his wife both died of influenza within hours of each other, and I was well enough to go to our bedroom window to see their coffins being lifted onto wooden biers to be carried by relays of colliers to the local cemetery. We all lived through the epidemic, but when we looked through the fence into the yard next door we realised with sadness that our manly was also condemned to remain for ever in this silent graveyard of immobile whelwrights.

AUSTIN WILSON, general practitioner, Aberystwyth.

We shall be pleased to consider for publication other interesting stories in general practice.—Ed, BMJ.