Hospital Topics

Present use of five-day wards

R DAVIES, K S CLIFF, W E WATERS

Abstract

A survey of five-day wards serving medical and surgical specialties indicated that 30 such units were operating in England in 1979. Six mixed surgical specialty, eight mixed specialty (medicine and surgery), four medical investigation, and 12 single specialty wards had opened since 1969. There was a varied, and often low, use of these wards. Nurse-staffing levels and allocation of operating theatre time also varied considerably. In busy wards nurse-staffing levels were high and access to operating theatre time reasonable. Medical investigation wards showed some of the highest levels of occupancy and throughput. The potential savings and service improvements that can result from five-day wards have yet to be realised on a wide scale.

Introduction

"Acute" five-day wards exist to examine and treat those patients whose stay is to be short and can be planned. This includes patients undergoing waiting-list surgery and some medical investigations.¹⁻⁴ The less intensive environment of the five-day ward is an attractive feature both in terms of nursing care required and the routine nature of the work, interrupted by few emergency admissions. In addition, the lack of weekend work is important in view of the reduction over the past decade of hours worked by nurses, which should lead to easier recruitment, particularly of married and part-time staff.

It may be argued that five-day wards are an uneconomic use of ward space and beds since, with closures at weekends and holidays, they may effectively be open for only half the year. But weekends and holidays are the most expensive periods as staff must be paid for overtime and even in traditional wards, bed occupancy is low.

Background and method

In 1974 a largely unpublished survey indicated that there were 23 five-day wards including wards for longer-stay cases.⁵ We excluded the latter type of ward. The interest in five-day wards was enhanced in 1975 by the publication of the DHSS circular *Management Arrangements for Waiting Lists*,⁶ which commended establishing five-day wards to reduce waiting lists. Central Government made

money available to health authorities to establish such wards. They were again cited as a main priority for the acute sector in *The Way Forward*.⁷ In 1975-6 the University of Southampton's department of community medicine carried out studies to examine the data sources of suitable surgical and investigatory procedures and to identify the organisational pattern, based on computer models, for admissions to five-day wards.^{8,9} In 1979, as part of a larger study funded by the South-west Hampshire Health District, a survey questionnaire was sent to the district administrators of districts identified as having five-day wards. Much of the data obtained related to 1978.

Results

Initially 32 health districts were identified as having five-day wards, but four failed to respond and two indicated that their five-day wards had closed. The results are based on the analysis of the completed questionnaire received in respect of 30 five-day wards in the 26 health districts.

Table I identifies the work in four types of ward and the reasons given for their establishment; nearly half had been opened as facilities additional to existing resources, most being new and purpose built. All the wards in the survey had opened since 1969.

TABLE I—Origin of five-day wards by type

Type of ward	Origin				Total
	Additional facility	Replaced conventional ward	Re-opening of closed ward	Part of a reorganisation (excludes other 3 reasons)	
Mixed specialty medical and surgical	6	—	1	1	8
Mixed specialty surgical	3	2	1	—	6
Medical investigation	2	1	1	—	4
Single specialty	2	5	1	4	12
Total	13	8	4	5	30

COMPOSITION AND SIZE OF WARDS

Mixed specialty medical and surgical wards—In these eight wards the number of beds varied from 12 to 32 (mean 20). All contained general medical and general surgical beds, and six also had orthopaedic beds; other specialties included dental and gynaecological surgery and cardiothoracic investigations.

Mixed specialty surgical wards—In these six wards (general surgery, orthopaedics, and sometimes dental and gynaecology) the number of beds ranged from 18 to 36 (mean 23).

Medical investigation wards—In these wards the number of beds ranged from eight to 29.

Single specialty wards—There were 12 single specialty wards, of which four were for gynaecological surgery. Of the remainder, two were devoted to ENT surgery for children, and the others to orthopaedic, dental, ophthalmic, and thoracic surgery, cardiothoracic investigations, oncology, and radiotherapy. The number of beds available varied from four to 21 (mean 12).

Community Medicine, South Academic Block, Southampton General Hospital, Southampton SO9 4XY

R DAVIES, MSC, research fellow (present address: Faculty of Mathematical Studies, University of Southampton, Southampton SO9 5NH)

K S CLIFF, FFCM, DPH, senior lecturer

W E WATERS, FFCM, DIH, professor

Half of the wards had fewer than 20 beds. Ward organisation varied considerably; some were independent units, others part of a large unit. Among those providing facilities for surgical inpatients, 11 out of the 26 had to share not only the theatre but also the lists with other general wards in the hospital. The single specialty surgical wards, even taking into account their small size on average, seemed to fare worst in respect of allocation of theatre sessions.

HOW RESOURCES ARE USED

Examination of the occupancy and throughput was based on the number of weeks each ward was available in the year immediately before the survey. Percentage occupancy is a misleading indicator of how resources are used, particularly where periods of stay are short. Throughput per bed is more commonly used, but this may have disadvantages with a large number of day cases. We have combined the two measures for the purposes of comparison (table II). Table III, which summarises throughput and occupancy by type of ward, must be interpreted with some caution because of the small numbers in each of the different types of wards. The results suggest that the

TABLE II—Classification of throughput and percentage occupancy. (The groups are exclusive)

	Admissions and day cases (per bed per week)		Average percentage occupancy (over nights open)	
High	Both > 1.50	and	> 70	
High-medium	Both > 1.30	and	> 60	
Medium	Both > 1.00	and	> 40	
Low	Either < 1.00	or	< 40	

TABLE III—Throughput and occupancy (see table II) of the wards by type

Type of ward	Throughput and occupancy			
	High	High-medium	Medium	Low
Mixed specialty medical and surgical	1	2	3	2
Mixed specialty surgical	—	—	3	3
Medical investigation	2	1	1	—
Single specialty	—	1	7	4

single specialty wards were not particularly efficient: this may be because of their relative shortage of theatre sessions. The mixed surgical wards were in the medium and low categories owing mainly to poor occupancies rather than low throughput. Occupancy and throughput were highest in the medical investigation and mixed medical and surgical wards.

Very different levels of nurse staffing between the wards were found, the mean number of nurses per bed ranging between 0.21 and 0.72. The busiest wards, however, had high staffing levels, and the one mixed specialty ward in this category had its own operating theatre.

Discussion

Plainly, the introduction and maintenance of five-day wards would be more readily justified by both high occupancy and high throughput. The results of our study suggested that in units with surgery these were more likely to be achieved where there was ready access to theatre sessions and high supporting nurse-staffing levels. Occupancy and throughput rates in units with low staffing levels and poor access to theatre sessions were poor. Districts with such wards gave various reasons for this, but it appeared that wards distant from the acute work in the district general hospital were particularly vulnerable to both low occupancy and low morale of staff. Consultants need to show interest and perhaps even enthusiasm for a ward to operate efficiently.

There seem to be two important and distinct theoretical factors in favour of five-day wards. The first is as an additional resource to reduce waiting lists for certain specific procedures. The second is economic, especially if weekends and holidays pose staffing problems, as five-day wards should reduce the cost or difficulty of running traditional wards. The results of

the present survey suggests that both these potential advantages are still rarely realised. This is surprising in view of the concern expressed in recent years about waiting lists and the limitations on expenditure. Even when five-day wards are achieved many are small, and their throughput and occupancy, especially in single specialty wards, leaves much room for improvement, although a few are very efficient. In general, the theoretical advantages of five-day wards do not yet seem to have become reality.

We thank the regional, area, and district officers for their co-operation in the study, Dr J G Dawe, Mr B Sartain, and Mr P A Shaw of the Southampton and South-west Hampshire Health District for their help, and Mr V Andrews of the community medicine department for data collection.

This study was supported by a grant from the Southampton and South-west Hampshire Health District.

References

- Longston D, Young B. The Manchester Royal Infirmary programmed investigation unit. *Br Med J* 1973;iv:528-31.
- Ingram RM, Traynar PM. Five-and-a-half-day ophthalmic ward. *Br Med J* 1976;ji:445-6.
- Sachdev Y, Gomez-pan A, Fletcher PM, Hall R. Programmed investigation unit. *Br Med J* 1976;jii:91-3.
- Burgess C, Chant ADB, Beschi J. Elective surgery and a programmed investigation unit in a five-day ward. *Hospital and Health Services Review* 1978;427-8.
- Anonymous. Five-day wards in 1974. *Br Med J* 1974;jii:71-2.
- Department of Health and Social Security. *Reduction in waiting times for in-patient admissions—management arrangements for waiting lists*. London: HMSO, 1975. (Circular HSC (IS) 181.)
- Department of Health and Social Security. *Priorities in health and social services. The way forward*. London: HMSO, 1977.
- Cliff KS. The use of hospital activity analysis data in assessing operative procedures suitable for five-day care. *Health and Social Services Journal* 1979; April: A29-32.
- Cliff KS. A computer model for a five-day surgical ward. *Public Health* 1979;93:4-10.

(Accepted 15 May 1981)

Is there any damage to children from clothes or bedding that have been stored with naphthalene moth repellent?

Naphthalene (tar camphor) is a major constituent of coal tar and is widely used industrially in manufacturing other chemicals including dyes, synthetic resins, and solvents. Its use as a moth repellent and insecticide decreased after the introduction of certain more effective chlorinated compounds for this purpose. The pattern of use is now changing again, however, because at very high doses some of the chlorinated compounds increase the risk of cancer in laboratory animals. It is to some extent absorbed through the human skin, and exceptionally heavy exposure by this or other routes—for instance, ingestion—has been associated with nausea, vomiting, headache, haemolytic anaemia, haematuria, rashes, cataracts, hepatic necrosis, convulsions, and coma in man. At one time naphthalene was used clinically as an anthelmintic, and there is a case report of a 6-year-old child who died after ingesting 2 g naphthalene over two days. The dose levels associated with signs of systemic toxicity are several orders of magnitude higher than those that might be absorbed from clothes stored with naphthalene moth repellent, and, although some individuals appear to be sensitive to relatively slight skin contact with naphthalene,¹ the consequential rashes usually disappear quickly after exposure ceases. A recent report of four cases of laryngeal cancer among a group of 15 cleaners using naphthalene,² although uninterpretable on its own because details of the smoking habits and other chemicals to which the group were exposed are not available, clearly indicates a need for further studies. There appears to have been no adequate test of naphthalene, using modern methods, for carcinogenicity in animals, although, from a consideration of its chemical formula, there are no grounds for expecting it to be dangerous in this regard.

¹ White RP. *The dermatogoses or occupational affections of the skin, giving descriptions of the trade processes, the responsible agents and their actions*. London: H K Lewis & Co, 1934.

² Wolf, O. Larynxkarzinome bei Naphthalinreinigern. *Z Gesamte Hyg* 1978;24: 737-8.