treatment the facial paraesthesia had ceased. After a further two days sensation was normal in the right arm and the sensory level on the trunk had descended to T6 on the right and T8 on the left. Grip in the right hand had improved (grade 3-4) and he was again able to move his right leg at the hip and knee (grade 3) and ankle (grade 2). After 10 days' treatment he was able to walk with a single stick. Five weeks after the start of the treatment he was able to walk unaided, having only slight proximal weakness in the right leg. Power was normal in his right hand, though dexterity was impaired. Sensation was normal, with the exception of vibration sensation, which was impaired below the sternum. The plantar response was extensor on the right, and reflexes were brisk in the right leg.

Comment

Herpes-zoster myelitis is rare, and in their review of the published reports Rose *et al*¹ found only seven reported cases and added a case of their own. At necropsy their case showed extensive haemorrhagic necrosis of the spinal cord. The neurological symptoms and signs developed four days to six weeks after the herpetic rash, and the myelitis progressed over hours, days, or weeks. Four of the patients had died while the others recovered with variable neurological deficits. Although an autoimmune response had been proposed as the process causing central nervous system damage after herpes-zoster infections,^{1,2} Hogan and Krigman³ subsequently showed direct invasion of the spinal cord by herpes-varicella virus in a case of ascending myelitis.

In our patient symptoms and signs of myelitis progressed inexorably for four weeks until treatment with vidarabine was begun. Vidarabine is an antiviral agent which has proved valuable in the treatment of herpes-simplex encephalitis and in herpes-zoster infections in immunocompromised patients.⁴ Its use has not been reported in herpes-zoster myelitis. Within days of starting the treatment our patient began to improve, and within five weeks he could walk unaided. Although herpes-zoster myelitis may improve spontaneously,¹ we think that the improvement was so dramatic after the start of treatment that it was not a mere coincidence. We therefore suggest that a parenteral antiviral agent should be given as soon as zoster myelitis is diagnosed.

¹ Rose FC, Brett EM, Burston J. Zoster encephalomyelitis. *Arch Neurol* 1964;**11**:155-72.

² Appelbaum E, Kreps SI, Sunshine A. Herpes zoster encephalitis. *Am J Med* 1962;**32**:25-31.

³ Hogan EL, Krigman MR. Herpes zoster myelitis: evidence for viral invasion of spinal cord. *Arch Neurol* 1973;**29**:309-13.

⁴ Whitley R, Alford C, Hess F, Buchanan R. Vidarabine: a preliminary review of its pharmacological properties and therapeutic use. *Drugs* 1980;**20**:267-82.

(Accepted 2 June 1981)

Department of Neurology, General Hospital, Nottingham NG1 6HA
R N CORSTON, MA, MRCP, senior registrar
S LOGSDAIL, MB, MRCP, senior house officer
R B GODWIN-AUSTEN, MD, FRCP, consultant neurologist

Gastric mucus secretion in ranitidine-treated patients

Ranitidine is a new H₂-receptor antagonist which at a dosage of 200 mg daily is reportedly as effective as cimetidine in the short-term treatment of duodenal ulcer.¹ Despite the remarkable antiulcer properties of cimetidine, this drug alters the composition of gastric mucus,²⁻³ which apparently influences the rate of ulcer recurrence when treatment is stopped. The quality of gastric-soluble mucus appears to affect duodenal as well as gastric mucosal defences. Maintenance treatment with carbenoxolone after healing of duodenal ulcers by cimetidine is associated both with return to normal of the gastric mucus and with a comparatively low relapse rate once treatment is stopped. In contrast patients healed and maintained with cimetidine whose gastric mucus is of persistently poor quality show a significantly higher rate of ulcer recurrences.³

Acute intravenous infusion of ranitidine failed to affect pentagastrin-stimulated output of N-acetyl-neuraminic acid, suggesting a lack of effect on mucus secretion.⁴ Nevertheless, the influence of prolonged ranitidine administration on gastric mucus has not been investigated.

Indeed, the effect of a single dose of the drug cannot be taken to predict the behaviour of mucus in ranitidine-treated patients. While gastric mucus secretion is unaffected by a single intravenous dose of cimetidine,⁴ appreciable alterations in the mucus properties are detectable after prolonged oral administration.²⁻³ We have studied the behaviour of gastric mucus in patients given ranitidine for one month.

Patients, methods, and results

Twelve outpatients aged 26-73 years with endoscopically proved duodenal ulcer were given ranitidine 150 mg twice daily for four weeks. Treatment was then stopped and all patients subjected to endoscopy. Neutral and acid mucoproteins in fasting gastric juice were measured before and after treatment and a "mucoprotective index" calculated as: $(B/(A+B)) \times 100$, where A=acid mucoprotein complex (g/l) and B=neutral glycoprotein complex (g/l), the latter containing the bulk (75.2%) of carbohydrates.^{2,3}

After the four weeks of ranitidine endoscopy showed complete healing of the ulcers in 11 of the 12 patients. No significant changes in mucoprotective indices were found after ranitidine (mean values before and after treatment 69.9 ± SEM 1.23 and 69.8 ± 0.9 respectively; $p > 0.05$ (Student's *t* test for paired data)).

Comment

Our results show that, unlike cimetidine, ranitidine does not influence gastric mucus secretion. This suggests that the cimetidine-induced alterations in gastric mucus are not related to the H₂-receptor blockage but are due to the drug itself. Differences between the two drugs have been reported. Aminopyrine metabolism, for example, is altered by cimetidine but is unaffected by ranitidine.⁵ Indeed, ranitidine appears to be structurally different from cimetidine, being a substituted furan derivative with no imidazole ring; and this may account for different behaviour of the two drugs in many respects. Whether the lack of damaging effects on gastric mucus by ranitidine will result in a lower rate of ulcer recurrence in healed patients remains to be determined.

Ranitidine was kindly supplied by Glaxo SpA, Italy.

¹ Langman MJS, Henry DA, Bell GD, Burnham WR, Ogilvy A. Cimetidine and ranitidine in duodenal ulcer. *Br Med J* 1980;**281**:473-4.

² Guslandi M, Testoni PA, Fesce E, Ballarin E, Tittobello A. Behaviour of gastric mucin during pirenzepine treatment: a double-blind controlled study versus cimetidine. *Current Therapeutic Research* 1980;**27**:714-8.

³ Guslandi M, Cambielli M, Tittobello A. Carbenoxolone maintenance in cimetidine-healed patients. *Scand J Gastroenterol* 1980;**15**:369-71.

⁴ Domschke W, Lux G, Domschke S. Furan H₂-antagonist ranitidine inhibits pentagastrin-stimulated gastric secretion stronger than cimetidine. *Gastroenterology* 1980;**79**:1267-71.

⁵ Henry DA, MacDonald IA, Kitchingman G, Bell GD, Langman MJS. Cimetidine and ranitidine: comparison of effects on hepatic drug metabolism. *Br Med J* 1980;**281**:775-7.

(Accepted 13 May 1981)

Gastroenterology Unit, 3rd Medical Clinic, University of Milan, Milan 20100, Italy

M GUSLANDI, MD, research assistant
E BALLARIN, MD, postgraduate fellow
A TITTOBELLO, MD, senior registrar

Primary systemic amyloidosis presenting as extreme hyperlipidaemia with tendon xanthomas

Hyperlipidaemia is not a recognised complication of amyloidosis in the absence of nephrotic syndrome. We describe a patient with primary systemic amyloidosis in whom tendon xanthomas associated with extreme rise in plasma remnant and low-density lipoproteins were presenting features and preceded detectable renal disease.