ABC of 1 to 7

J A FIXSEN
H B VALMAN

LIMP IN CHILDREN

The child with a limp may have a minor injury which will recover spontaneously in a few hours or a condition which may affect him for life if treatment is delayed. A diagnosis must be made, or if this is not possible the child should be observed in hospital until he has fully recovered. A full history and careful examination of the whole child is essential. A radiograph of the affected area is taken, but pain in the thigh referred from the hip can lead to the wrong area being examined. Parent's observations may sometimes sound odd but should not be dismissed. Abnormal physical signs are usually present in any child with a limp which has a serious cause. Limp may be due to pain, leg inequality, neuromuscular dysfunction, or, rarely, psychological disturbance. These groups often overlap.

Pain

The commonest cause of limp is pain. Injury may produce muscle strain or fractures but non-accidental injury or foreign body should always be considered. Any pain which does not settle within a few days or recurs should be investigated. Perthes's disease, slipped upper femoral epiphysis, and benign tumours such as osteoid osteoma often produce intermittent pain, which can easily be dismissed as a recurrent muscle strain.

Transient synovitis (observation hip, coxalgia fugax) is a common condition of unknown cause. The child suddenly starts limping and there is spasm of the hip muscles but no other abnormal signs. He may have a mild fever and raised erythrocyte sedimentation rate. Transient synovitis usually settles with a short period of bed rest. Traction is helpful to relieve muscle spasm and pain. Most children have no further problems, but a few may have recurrent attacks and subsequently develop changes of Perthes's disease or juvenile arthritis. It is important to eliminate more serious conditions such as septic arthritis or osteomyelitis, Perthes's disease, juvenile chronic arthritis, slipped upper femoral epiphysis (which is outside the age group of this article), and tuberculosis.

Septic arthritis—The child with septic arthritis is usually seriously ill but beware of the child suffering from immune deficiency, immune suppression, or who has been given a short course of antibiotics which mask the signs and symptoms. Perthes's disease and slipped upper femoral epiphysis should be seen on a radiograph. It is important to take a frog lateral view as well as an anteroposterior view of the hips to diagnose and evaluate these two conditions. Juvenile chronic arthritis and tuberculosis are likely to have a longer time course and also show radiographic changes.

Perthes's disease—Treatment for Perthes's disease is confusing both to the general practitioner and the orthopaedic surgeon. There is evidence which suggests that Perthes's disease is due to temporary ischaemia of the femoral head. Treatment cannot prevent the disease but is aimed at improving the outcome. Many patients need no treatment after the acute episode of pain and spasm. Regular radiographic monitoring of the hip is necessary to ensure that the femoral head remains within the acetabulum. If the femoral head begins to move out of the acetabulum abduction splints, plasters, or surgery may be needed to achieve containment of the head.

Osteochondroses—Osteochondritis of the tarsal navicular may cause pain over the medial arch of the foot. A radiograph will confirm the diagnosis. Most cases settle with reduction of activity. Sometimes a short period in a below-knee plaster is necessary. Osteochondritis affecting the heel or the tibial tubercle and osteochondritis dissecans of the knee rarely occur in this age group.

Verrucas, foreign bodies, and minor fractures may also cause pain in the foot. Persistent swelling, particularly around the ankle, suggests juvenile chronic arthritis, which commonly presents in the foot in children.

In this age group pain around the knee is uncommon although it becomes increasingly common in early adolescence. Causes of pain in the knee include bone and joint sepsis, chronic inflammatory conditions, and mechanical derangement such as a discoid meniscus.

Leg inequality

A short leg must be distinguished from apparent shortening due to pelvic obliquity, scoliosis, or joint deformity. Congenital defects such as short femur or tibia are usually obvious. Despite early screening programmes congenital dislocation of the hip must always be considered. Typically the leg is short and externally rotated, and abduction in flexion is limited. The child usually walks with the leg straight on the dislocated side and the knee flexed on the normal side. The doctor may consider that the flexed knee is abnormal and fail to recognise that the opposite hip is abnormal. Hemiplegic cerebral palsy, spinal dysraphism, and poliomyelitis cause poor growth of the affected leg and can be detected by a neurological examination. The whole spine should be examined for the warning signs of dysraphism: a naevus, hairy patch, pit, or lipoma. Epiphyseal injury from trauma or sepsis can cause shortening and deformity.

Neuromuscular and hysterical disorders

Complex disturbances of gait, such as incoordination or weakness due to cerebral palsy, muscular dystrophy, or spinal or cerebral tumour, may be described by the parents as a limp. A careful history and examination will usually suggest the diagnosis.

If a limp has an emotional origin diagnosis may be difficult. Usually the pattern of gait is bizarre and the physical signs inconsistent. It is essential to eliminate any possible organic cause. Often it is necessary to admit the patient to hospital for a few days' observation. A bone scan may help to eliminate the possibility of an abnormality of bone which has not been detected by radiographs.

Mr J A Fixsen, MCHIR, FRCS, is consultant orthopaedic surgeon, Hospital for Sick Children, Great Ormond Street, and St Bartholomew's Hospital, London, and Dr H B Valman, MD, FRCP, is consultant paediatrician, Northwick Park Hospital and Clinical Research Centre, Harrow.