PRACTICE OBSERVED

Trainees' Corner: Managing Chronic Disease

Diagnosing arthritis

RAY MILLION, TERRENCE REILLY

This article is based on an audiovisual presentation made for vocational trainees in general practice by the MSD Foundation. Further information about the video-cassette programme on which this series is based is available from the MSD Foundation, Turretstock House, Turretstock Square, London WC1.

Although this article deals with specific arthritides, we want to make some remarks about rheumatic disorders in general. When presenting evidence to the Royal Commission on the National Health Service the British League Against Rheumatism wrote:

"Rheumatic disorders are responsible for a large part of the general practitioner's work load, but all too often the quality of primary care for patients with these conditions leaves much to be desired. Doctors with special expertise, including some engaged in primary care, appear to be able to respond effectively to greater demands without distortion of their work burden. There is an urgent need for more relevant education for those intending a career in general practice as well as for provision of appropriate supporting facilities. A great deal of improvement could be accomplished without the need for additional resources."

During a year something like 150 patients for every 1000 on his list will consult a general practitioner with a rheumatic problem, and over the age of 65 this proportion may be doubled.

Eccles, Manchester M30 9EQ

RAYMOND MILLION, MR, FRCP, general practitioner and clinical assistant, Rheumatic Diseases Centre, Hope Hospital, Salford

Weiss National School of Medicine, Cardiff

TERRENCE REILLY, MR, MRCP, senior lecturer in general practice

Objectives

This article aims at stimulating trainees to consider the objectives of managing the patient with arthritis in general practice and in particular:

- to consider the methods available for diagnosis;
- to emphasize the need to detect early signs of the disease which might be aborted or have its subsequent course modified;
- to recognise and manage wide ranges of common conditions which need not require the more costly interventions of the hospital services;
- to gain insight into the factors requiring continuing care in chronic conditions.

In Britain more than 8m people consult their general practitioner each year with a rheumatic problem, and such problems account for the loss of some 40m working days each year. What impact can general practice make on this problem? The authors of the Brindle Lodge report wrote in 1974:

"Primary care in the community is the first line of contact for those with locomotor diseases. It was felt that with appropriate support and education and ready access to expert advice when difficult problems are encountered, general practitioners could deal with a large proportion of the management of these problems without the need for referral to hospital. Difficulties in diagnosis would, of course, still be likely to call for a specialist opinion in a number of instances."

Arthritis in its many forms is thus one of the commonest problems in general practice. Surveys have shown that musculoskeletal disorders account for between 20 and 25% of general practice consultations. They pose problems of diagnosis, investigation, and management. This article is in two parts: in

sanitation. He was a good parent and a concerned family man with regular but poorly paid employment. He was not a frequent attender at the surgery, but he had come because he was becoming increasingly worried about his painful knee. The doctor discovered that he had first noticed the pain when playing with his children. There were no medical records available, and he did not give a history of related or associated disorders. During the consultation the doctor elicited that there was no family history of hereditary disease and that his children, aged 5 to 14, were all at school and were well.

On examination the right knee was warmer than the left, and it had lost its normal dimples, suggesting an effusion. On moving the knee there was crepitus in mid-flexion and pain at full flexion but not at full extension. The patellar tap was positive on the right but not on the left. The right quadriceps measured 38 cm round 12 cm above the top of the patella as opposed to 40 cm on the left; this difference was probably important as Mahmet was right-handed. Further examination showed that there was no finger clubbing, no cervical lymphadenopathy, the trachea was central, and the apex beat not displaced.

So what was the diagnosis? This was a big problem and difficult to manage in general practice. From the history we had a man, a first generation immigrant, who had gone back to Bangladesh a year ago for a visit, who was now in Britain working, and who had had a minor injury to one joint. The trauma might be dismissed by the doctor as a trivial event and the patient might be advised to go away, do something simple, take some rest and analgesic, and come back next week if it was still bothersome. And this could be a disaster.

He may have had the first evidence of an atypical form of an inflammatory polyarthritis, and one might be tempted to sit back and wait and see what happened. Conversely, you might think that he had got into an acute form, but the joint had been uncomfortable for a few weeks, which is not typical of gout. A septic arthritis was a possibility, and one might think it could have cost him his leg or his life—a dramatic statement but quite true, and this was one of the most important points. Because of the findings elicited in the history, tuberculosis also had to be considered.

With this differential diagnostic dilemma this was the point at which the general practitioner should cease to work alone in the case. He should telephone a rheumatologist or, if not available, a consultant physician, and possibly ask him to come out and discuss the case. Specialised investigation was of paramount importance, and before the day ended that man should have been in hospital.

Mary Evans

The last patient was Mary Evans. She was also complaining of knee trouble but in both of her knees. She was 56 years old, overweight, and had had the pain in her knees for several weeks. The pain was bad enough to make her limp but was relieved slightly when she took the weight off her feet and rested. She also had pain in all of her right leg, down the side of the ankle, up to the hip, and into the groin. She had no other complaints and her previous medical history was unremarkable. To her doctor's frustration she had recently gained a little weight.

On examination her hands were normal and she had full painless extension and flexion in both of her knees, although there was some crepitus in the right knee. There was a fixed flexion deformity in the right hip and some loss of flexion in the left hip. How can you distinguish hip from knee involvement, and what is the cause of these symptoms?

This was a woman in late middle age whose sole problem was pain of fairly recent origin in the knees. The doctor found normal peripheral joints and elicited a positive Thomas test—the fixed flexion deformity in the right hip. She also showed that the knees, which the patient had said were being painful, were in fact mobile, stable, and had a little crepitus. An alternative technique for differentiating the source of pain between hip and

knee is to sit a patient on the side of a couch with the legs dangling free and the popliteal fossa cushioned by the side of the couch and then ask her to extend the knee fully, and if this is painful it is a good sign that the source of pain is the hip and not the knee. It was important to bear in mind that she had a limp and that the examiner had elicited some short shortening in the right leg. Such patients walk in a dot-and-carry-one fashion and often therefore complain of low backache, which may be a misleading symptom if leg measurement is not precisely assessed.

This particular woman had normal hands, but commonly an examination of the peripheral joints in the upper limbs is valuable because middle-aged and elderly patients have signs of generalised osteoarthritis in the form of Bouchard's nodes in the proximal interphalangeal joints or Heberden's nodes in the distal interphalangeal joints. In this instance there was no morning stiffness, and no synovial thickening, swelling, or tenderness was elicited anywhere. It was fairly clear that the diagnosis was osteoarthritis in the right hip. Steps would have to be taken to show that this was the case and that there was no other overriding cause for her symptoms, but that would be straightforward.

Diagnoses confirmed

We have now considered four patients with joint disorders: Brian; Doris; Mahmet; and Mary. We have studied their case histories, and we have made presumptive diagnoses of ankylosing spondylitis, rheumatoid arthritis, tuberculous arthritis, and osteoarthritis. What tests were necessary to confirm or reject each of these diagnoses? And which of these should be handled in general practice?

Brian Hall's presumptive diagnosis was ankylosing spondylitis. All test results were within normal limits, though the erythrocyte sedimentation rate (ESR) was at the highest limit of normal at 20 mm in the first hour. Urine analysis showed normal glucose and protein. X-ray examination of the dorsolumbar spine and sacroiliac joints were requested, and the first showed the changes typical of early ankylosing spondylitis. Some patients may show the later stages characterised by ossification of the spinal ligaments plus erosion and fusion of the apophyseal joints, the so called bamboo spine. In Brian's case the sacroiliac joint surfaces were eroded with sclerosis in subchondral bone. Brian had ankylosing spondylitis.

Doris Taylor was found to have a haemoglobin of 9.8 g/dl and an ESR of 75 mm in the first hour. Sheep cell agglutination and latex fixation tests were positive. Urine analysis showed normal protein, red cells, and the throat swabs were sterile. The diagnosis of rheumatoid arthritis was probable. Mahmet Kashmi was admitted to hospital for investigation and treatment. His haemoglobin at 14 g/dl was normal, but he had an ESR of 66 mm in the first hour and a leucocytosis of $15 \times 10^9/l$. Aspiration of the joint and synovial biopsy specimen confirmed the diagnosis. He had a tuberculous knee joint. Mary Evans had a normal ESR as expected. The diagnosis was made on the x-ray examination of the pelvis. She had severe osteoarthritis of both hips.

So our four diagnoses are confirmed. Now come the problems of management, which will be discussed in the next article.

We are grateful to Dr Tom Smith and Dr Sally Hull, who helped with the preparation of this programme.

References

- Summary of the evidence, Section 6, relating to rheumatic disorders, presented to the Royal Commission on the National Health Service, British League Against Rheumatism, 1977.
- Options in the delivery of medical care. The Brindle Lodge Report, 1974, para 2.

the first we consider four typical patients, all with joint disease, who might present themselves at any surgery. You are asked to put yourselves in the position of their general practitioner. By the end of the first article their diagnosis should be clear. The second article deals with management, which of course encompasses much more than prescription writing.

The four pseudonymous patients who might present themselves in your surgery are: Brian Hall, a 30-year-old man with backache; Doris Taylor, a woman nearing 50 with pains in her joints; Mahmet Kashmi, a 40-year-old immigrant from Bangladesh with a painful knee; and Mary Evans, a woman in her late 50s whose knees are bothering her. At first sight they have little in common, yet they all have disorders of their joints, and a systematic approach to their problems will help to ensure speedy and accurate diagnosis and the selection of the most appropriate management.

Brian Hall

Brian had already labelled his own illness as sciatica, as his uncle had had a similar complaint some years ago. He came saying that the pain in his back had become bad again. His medical records showed that he rarely attended the doctor, and his only consultations had been for well-defined incidents. He dated his present symptoms from a fall two years previously. He was stiff in the morning, but felt better in the evenings. Exercise hurt, but later it relieved the stiffness.

Examination showed no scoliosis but some lumbar lordosis, and Brian indicated that he felt pain in his buttocks and between the shoulders. When he touched his toes the doctor's fingers, placed on his spinous processes, did not separate at all, indicating that he had a very stiff spine. When asked to slide his hand down his right leg and then his left leg, he could manage only about 10 cm movement on each side, showing that the limitation of movement was considerable but equal. When asked to lean back as far as possible Brian could not move his back but did all his leaning from his hips. Testing neck movements showed loss of lateral mobility. Pressure on the sacroiliac joints gave Brian pain, particularly on the left side. Finally, measuring Brian's chest in inspiration and expiration at the nipple line showed that there was only 2 cm of expansion.

So what was the next step and what were the possible diagnoses? Several features in the history were very important: one was that he thought he had sciatica; another was that he had morning stiffness; and a third was that he had pain both between the shoulder blades and in his buttocks. Fourthly, his stiffness was relieved by exercise. The examination brought out evidence of limitation of movement in the cervical and thoracic spine, and of normal hips (the pain in the buttocks was a confusing sign because it could arise in the back and it could arise in the hip joints); the straight-leg raising test indicated that probably there was no most compression in the lumbar spine. This was important because he gave a history of having had a fall and he may have had a mechanical disorder of the back. The history suggested that this was not because of the morning stiffness and because of the vague distribution of pain—where it mechanical it would probably have been more localised. This man showed all the signs of an inflammatory disorder of the spine of some severity.

It is very important to understand that not every patient who has a disorder like this will show such obvious signs. In particular, the lateral flexion of the spine where the fingertips failed to reach the popliteal crease by several centimetres will not always be present. In an early case it may be possible to show that the symptoms are all the same, but the man may reach his popliteal crease and touch his toes with little difficulty. But the fact that he had difficulty with the pelvis fixed in rotating his dorsal and lumbar spine was a pointer to an inflammatory disorder in the spine.

So we have built up a picture of a man quite severely affected by a disorder that looked inflammatory in its history and in its clinical presentation; a mechanical disorder of the lumbar spine

was very unlikely to show limitation of rotation and limitation of lateral flexion but only of forward flexion and extension. This was a man with a spondyloarthropathy of the ankylosing spondylitis type, and we would want to do further investigations to make a more accurate diagnosis.

Doris Taylor

Doris was 48, and her record folder was remarkably thin. She had had few clinical problems, but she was very upset: in recent months she had developed pain and swelling of small joints all over her body. She had morning stiffness for two hours and felt feverish and vaguely unwell. Obviously, the first diagnosis that came to mind was rheumatoid arthritis—but how to confirm it?

Examining Doris's hands showed typical spindling and some palmar erythema but no vasculitis of the fingertips. There were no subcutaneous nodules to be felt. Squeezing the proximal interphalangeal joints was painful, but extending the fingers Doris described as uncomfortable rather than painful. The distal interphalangeal joints were normal. Extension of the wrist joint was painful on both sides. The doctor felt for nodules or trigger fingers by placing her hand on the flexor tendons and then flexing and extending the fingers—she could not feel anything. Doris had difficulty putting her hands behind her neck and her back. Trying to abduct Doris's right arm with her scapula fixed, the doctor showed a considerable restriction of glenohumeral movement, but when she took her hand off the scapula further elevation of the arm was possible. There was no limitation of movement in the neck, and Doris reported that movement was not painful but was noisy. She had no pain on opening her mouth wide. Further examination showed tenderness in the joints of the forefeet.

Doris again

With Doris we can piece together a picture that is fairly conclusive. In the last few months she has had some pain with a lot of stiffness in the morning, and this had affected her mainly in the joints of the limbs. She had felt ill; she had not been able to get about and fulfil all her responsibilities; and she had been disturbed for all those reasons. The examination showed that she had pain, tenderness, and limitation of motion in a lot of joints: in the small joints of the hands—the distal interphalangeal joints—and the wrists, the shoulders, and the forefeet. The conclusion we reach is that the inflammation is in soft tissue, in synovial tissue. The examiner has tried to elicit tenderness in the temporomandibular joints; people sometimes spontaneously complain of difficulty in chewing their food, and, of course, that is a synovial joint like the others. So we have a picture again of an inflammatory arthritis, but this time affecting the peripheral joints. It affects them symmetrically, is of recent onset, and omits the distal interphalangeal joints—this is a diagnostic pointer. We can conclude that she has an inflammatory polyarthritis of the rheumatoid type.

The general practitioner must decide whether this is rheumatoid arthritis or whether it is one of a variety of disorders that mimic it: psoriatic arthropathy; sarcoidosis; systemic lupus erythematosus; or polyarteritis nodosa—rarer syndromes. A general practitioner would not be expected to deal with. Now complex investigations are needed, but the general practitioner will be able to organise these.

Mahmet Kashmi

The next patient was 40-year-old Mahmet Kashmi. Mahmet had come to Britain several years previously from Bangladesh and had returned only once, last year. He now lived with his wife and children in an overcrowded terraced house with poor

Organising a Practice

A general practice expands

P G KAY

There has been a tendency for group practices to get larger in recent years. This paper describes how one large practice is coping with further expansion. Large practices have their own administrative problems, and perhaps it is time that we took stock to identify these problems and discuss solutions.

Setting

The practice is based in Witney, an old market town 12 miles west of Oxford. Two-thirds of the patients live in the town, and the remaining third in villages, mainly within five miles of the town centre but up to eight miles in one direction. The main employers in the town are a car components factory, the ancient blanket-making industry, and a food processing factory. Many small industries have grown up recently, and the area acts as a dormitory for Oxford. There is a full spread of patients throughout the social classes, with classes 3 and 4 predominating.

Since 1948 the family practitioner services in the town have been provided by one practice, which had grown to six partners by the late 1950s. On the periphery of the practice there is territorial overlap with neighbouring practices. A branch surgery, owned by the practice, is in a village six miles from the town centre. There has been no cottage hospital in the area, so we looked to Oxford for all hospital services, except that orthopaedic and ophthalmology clinics were held on a regular basis in a church hall.

The 1960s

By the late 1950s the rented practice premises had become too small. After extensive negotiations the Nuffield Provincial Hospitals Trust agreed to finance a health centre to bring together the three strands of the health service—hospital, local authority, and general practice—under one roof. At the end of 1965 the partners moved into the Nuffield Health Centre, which provided them with a suite of six consulting rooms, an examination room, a treatment room, reception area, and secretary's office. There were rooms for health visitors and district nurses, who had previously worked from their homes. There was a physiotherapy department under a physiotherapist, a small x-ray department, a small x-ray department, capable of taking chest and "cold" orthopaedic x-rays, and a suite of consulting rooms for visiting specialists from Oxford who held outpatient clinics in the evenings. The district nurses, district nurses, and health visitors increased rapidly and an effective primary care team was built up, while the links between the team and the hospital services steadily improved.

Witney Health Centre, Witney, Oxon
P G KAY, MR, MRCP, general practitioner

At this time the population was expanding fast—about 500 extra patients a year were being added to the practice list—so that by 1968 there were an average of 3000 patients per partner. We felt we were a large enough group, but we also realised that this number of patients was too large for good practice so we decided to take on two new partners in the next three years. We were able to take over some rooms in the Nuffield Health Centre as consulting rooms.

The early 1970s

We became a training practice, which added another "pair of hands," but we soon realised that the advantages were balanced by the time taken up in teaching.

At the beginning of the decade the local planners predicted that there would be 31 000 people in our practice area by 1981—a further 10 000—and we knew that there was no room for further expansion in the health centre. In the early 1970s the Oxford Regional Hospital Board planned to develop its community hospital programme and wanted to site a new community hospital in Witney. In 1972 it decided to do this and include a new health centre as part of the building. It was agreed in principle that half of our practice would move into the new centre. A planning team included two of the partners.

In the Nuffield Health Centre there is only one examination room, the treatment room is some distance from the consulting rooms, and the reception area is too small. We used our influence to ensure that the reception area was much larger and that the treatment room was close to the consulting rooms. The health centre design guide at the time insisted on one examination room for each consulting room and, though we thought that these would be too many, they were provided. The project was ready to go to tender by the end of 1973, when, because of the sudden cut-backs in public expenditure, the project was shelved.

The mid-1970s

The Oxford Regional Hospital Board was replaced by the Oxford Regional Health Authority, and the Oxfordshire Area Health Authority (Teaching) (OAHAT) came into being. Of the original planning team only the two GPs and one nursing officer remained in the area. When limited funds became available it was decided to build the health centre as a separate block. Work started in 1976 and was completed by the end of 1977. It provided eight consulting rooms and examination rooms, a treatment room, a large reception area, a health education room, and offices for the district nurses and health visitors, with a medical loan store and chiropody room.

At completion, the OAHAT had no money to allocate for the costs, and the predicted population expansion had not occurred. So the practice was several thousand patients smaller