may not prove to be cause and effect. The fetal abnormalities caused by thalidomide and the oculomucocutaneous syndrome caused by practolol were brought to doctors' attention by letters to the *Lancet*¹ and the *BMJ*⁴ respectively, and in each case the association was quickly confirmed by further reports.

In contrast, associations may prove false alarms which cast an unwarranted blight on a useful drug. One such was the report associating skeletal malformations with Debendox,⁵ which has not been confirmed by later large-scale studies⁶ but played a part in the loss of public and medical confidence in the use of the drug by pregnant women.

The *BMJ*'s current policy aims at steering a path between the extremes of crying wolf too often and insisting on near-certain evidence. In general, we are likely to accept a report of a drug side effect if it describes more than one case and if the evidence points clearly to a single drug. Something more than simple coincidence in time is usually required—rechallenge (with the patient's informed consent) or immunological investigations may tip the balance of probabilities. Two other vital sources of information should be checked by authors before submitting a report: Has the manufacturer been told of any similar episodes? And have any reports been made to the Committee on Safety of Medicines?

¹ McBride WG. Thalidomide and congenital abnormalities. *Lancet* 1961; ii:1358.

² The Sundry Times Insight Team. *Suffer the children: the story of thalidomide*. London: Andre Deutsch, 1979.

³ Griffin JP, Diggle GE. A survey of products licensed in the United Kingdom from 1971 to 1981. *Br J Clin Pharmacol* 1981;12:453-64.

⁴ Wright P. Untoward effects associated with practolol administration: oculomucocutaneous syndrome. *Br Med J* 1975;ii:595-8.

⁵ Donnai D, Harris R. Unusual fetal malformations after antiemetics in early pregnancy. *Br Med J* 1978;ii:691-2.

⁶ Harron DWG, Griffiths K, Shanks RG. Debendox and congenital malformations in Northern Ireland. *Br Med J* 1980;281:1379-81.

Episiotomy

Childbirth in Britain has never been safer than it is today, though we are not yet among the European leaders in the league table of lowest maternal and perinatal death rates. Understandably, pregnant women, their husbands, and doctors are all paying more attention to subtler aspects of care, especially those concerned with making the experience of pregnancy and childbirth more comfortable and pleasurable. With increasing insistence individual women, and sometimes well-organised groups, are asking whether some procedure is manifestly to the advantage of mother and baby or amounts to unnecessary interference by doctors. Among the targets have been the steadily rising incidence of induction of labour (the subject of considerable debate a few years ago¹⁻⁵), forceps delivery, and caesarean section; and the whole matter of intervention in obstetric practice was recently the subject of special study at a scientific meeting organised by the Royal College of Obstetricians and Gynaecologists.⁶

The spotlight of public concern has now moved on to episiotomy. The National Childbirth Trust has recently published a collection of essays⁷ on the physical and emotional aspects of episiotomy with contributions from obstetricians and midwives, concluding with Sheila Kitzinger's assessment of its effects on postnatal sexual adjustment. On page 243 Reading *et al* report their account of patients' attitudes towards the pain and discomfort that may occur after episiotomy. All these studies show how many questions remain unanswered.

Here is a surgical procedure widely used by doctors and midwives, yet we have few objective data to support claims that perineal incision performed correctly at the appropriate moment eases delivery, protects the head of a small baby from trauma, is more easily repaired than a ragged tear and will heal more quickly and effectively, is less liable to infection than a bruised and torn perineum, and reduces the risk of later complications such as dyspareunia and prolapse. Reading *et al* were largely concerned with the discomfort associated with episiotomy, and the findings are disturbing: many women had severe pain at the time and afterwards, and nine out of 10 who subsequently experienced dyspareunia attributed this directly to the episiotomy scar.

In some aspects, the study of Reading *et al* confirms work⁸ that the *BMJ* discussed in 1973,⁹ and the questions remain the same. Unfortunately we still do not know whether the pain associated with episiotomy is greater or less than that associated with a perineum that has been allowed to become bruised and torn. On first principles most—though not all¹⁰—obstetricians would argue that a clean surgical incision in the perineum, correctly timed and repaired, is more likely than a ragged, bruised tear to heal by first intention and cause less trouble at the time and later. But it would be helpful to have firm evidence to support or refute this belief. And as women themselves become better informed and more articulate they are sure to have strong views on this important subject. It would, however, be a pity if clinical practice were changed on insufficient evidence because of a patient-led protest. The answers should come from clinical research.

J K RUSSELL

Professor of Obstetrics and Gynaecology,
Princess Mary Maternity Hospital,
Newcastle upon Tyne NE2 3BD

¹ Anonymous. A time to be born. *Lancet* 1974;ii:1183-4.

² O'Herlihy C, McDonald D. Planned delivery. *Lancet* 1975;ii:126.

³ Bonnar J. Selective induction of labour. *Br Med J* 1976;ii:651-2.

⁴ Chalmers I, Zlosnik JE, Johns KA, Campbell H. Obstetric practice and outcome of pregnancy in Cardiff residents 1965-73. *Br Med J* 1976;ii:735-8.

⁵ Leeson J, Smith A. Induction of labour and perinatal mortality. *Br Med J* 1977;ii:707.

⁶ Beard R, Paintin DB, eds. *Outcomes of obstetric intervention in Britain. Proceedings of a scientific meeting of the Royal College of Obstetricians and Gynaecologists, May 1980*. London: Royal College of Obstetricians and Gynaecologists, 1980.

⁷ Kitzinger S, ed. *Episiotomy: physical and emotional aspects*. London: National Childbirth Trust 1981.

⁸ Baker S. *A survey into post-natal perineal discomfort*. London: S Maw and Sons, with the Royal College of Midwives, 1973.

⁹ Anonymous. Pain after birth. *Br Med J* 1973;iv:565.

¹⁰ Barker GH. The unkindest cut of all. *World Medicine* 1981;16:40-1.

Children's accidents

1980 was designated Year of the Child and 1981 International Year of Disabled People, and all too often the two concepts overlap. Increasingly, accidents have become a main cause of mortality and morbidity in children of all ages. Reduction in mortality from infectious disease and improvements in chemotherapy and surgery have reduced infant and childhood mortality from other causes, with a consequent increase in the proportion attributable to accidents.

Concern about children's accidents has been given a focus with the formation of the Child Accident Prevention Committee, which is supported by Government and voluntary funding. In conjunction with the BBC, the Health Education

Council, and the Scottish Health and Education Group the committee has brought children's accidents into the public eye through the television series *Play it Safe*. The series is accompanied by a booklet,¹ edited by Dr Hugh Jackson, and contains a foreword by the presenter, Jimmy Savile, urging that "accidents hurt children, often seriously . . . So do yourself and your kids a favour. Play it safe."

Such an appeal from a celebrity relies on people's emotions to change their attitudes, but the programme also uses facts convincingly—and for the apostles of prevention the end justifies any means.

What are the problems that the Child Accident Prevention Committee is trying to tackle? The *Play it Safe* booklet sets out some bleak statistics. Four children each day die as the result of an accident; one in six of the children in a paediatric ward has accidental injuries.

These figures raise other questions, starting with the fundamental "What are the distribution and determinants of this disease in the childhood population?" Accidents can be investigated by epidemiological techniques in just the same way as other disease. Four questions then have to be answered: Who is affected? Where does the event occur? When does it happen? and What is the cause?

Data from the Office of Population Censuses and Surveys² show that the proportional mortality from accidents is 3% in children under 1 year and around 26% in 1-4-year-olds. From 5 to 14 years the proportional mortality in boys is 40-45% and in girls around 28%. Analysis of data from the same source on age-specific mortality rates from road-traffic accidents shows that the peak occurs between the ages of 5 and 9 years and that boys have twice as many deaths as girls. This should lead the epidemiologist to examine in greater depth the type of road user and the mortality experience by the different groups—for example, the rates among pedestrians and cyclists. Next, we need to know where the road users' accidents occurred.

Caution is needed, however, in interpreting these findings. The analysis should recognise the problem of deciding why there are these differences both between sexes and, for example, between road-user groups. Are these real differences or are they due to increased exposure to risk? Are boys really different from girls in their concept of danger, or are they more likely to be exposed more frequently to risk because of their particular life style?

Consideration of where accidents occur to children is equally important. For example, in children's accidents on farms one important danger is slurry pits, which may look just like a rather boggy part of a field. Studies of road-traffic accidents to child pedestrians have shown that the walk or ride home from school presents the major dangers, and specifically where children have to cross at a T junction. Moreover, the relative distribution of different types of accidents plays an important part in determining where efforts should be concentrated on prevention. Should our efforts be put into home accidents, road accidents, school accidents, or farm accidents? Analysis of where an accident occurs gives an indication of the relative risks, home accidents being by far the greatest problem in the 0-4-year age group, whereas road-traffic accidents play more significance for the age group 5-14 years.

Finally, when do accidents occur? Hancock's study³ of accidental poisonings in children found that 15% occurred between 0800 and 0900 and 27% between 1700 and 2000. Both periods are times of maximum family busyness, when the mother in particular may have to turn her attention to a range of activities and family members.

Answers to questions about who, when, and where may lead

to a greater understanding of why accidents occur. Sabey and Taylor's investigation⁴ of road-traffic accidents identified three factors: the road user, the vehicle, and the road environment. These provide useful analogies for all accidents. We need to ask, firstly, was the user experienced enough; did he or she understand the potential dangers? Secondly, was the object that caused the injury as safe as was possible? and, thirdly, was the environment as safe as possible?

The three factors may contribute individually, or in association with one or both of the others, to differing degrees in different accidents. Our understanding of the interaction, however, is the key to prevention.⁵

The question still sometimes asked, however, is, "Is it all worth while?" What effect can the medical profession have? Doctors can try to educate the public and give them the facts; that is what *Play it Safe* is trying to do. Next comes legislation, where the medical profession can have a powerful influence on Government thinking. Without medical commitment, or at least commitment by individuals and groups within the profession, there would be no Child Accident Prevention Committee. Without basic epidemiological studies into the effectiveness of seat belts,⁶ and the ineffectiveness of educational programmes,⁷ the British Medical Association and other professional bodies could not have pressed for and seen legislation introduced. Perhaps the profession needs, therefore, to reconsider where it is going and give greater thought to the possibilities of prevention rather than treatment of children's accidents; or will doctors be satisfied to carry on as usual and *Play it Safe*?

K S CLIFF

Senior Lecturer in Community Medicine,
University of Southampton,
Southampton SO9 4XY

¹ Jackson RH, ed. *Play it Safe*. London: Health Education Council/Scottish Health and Education Group, 1981.

² Office of Population Censuses and Surveys. *Mortality statistics: cause 1979*. London: HMSO, 1981. (Series DH2 6.)

³ Hancock BW. Accidental poisoning in children. *Br J Clin Pract* 1973;**27**: 77-80.

⁴ Sabey BE, Taylor H. *The known risks we run: the highway*. Crowthorne: Department of Transport, 1980. (TRRL report SR 567.)

⁵ Jackson RH. *Children, the environment and accidents*. Tunbridge Wells: Pitman Medical, 1977.

⁶ McDermott F. Control of road trauma epidemic in Australia. *Ann R Coll Surg Engl* 1978;**60**:437-50.

⁷ Wessex Positive Health Team. Promoting the use of seat belts. *Br Med J* 1980;**281**:1477-8.

Effect of transplantation on non-renal effects of renal failure

The patient who receives a kidney transplant is usually rapidly restored to wellbeing thanks to the return of normal renal function, which causes many of the secondary effects of chronic renal insufficiency to disappear. For example, when there are adequate iron stores anaemia resolves in six to 12 weeks and occasionally erythrocytosis occurs, and with the restoration of normal haemoglobin concentrations angina may also disappear.

About half of all transplanted patients, however, have appreciable hypertension; in most this predates the transplant and has not been improved by it. But the incidence of hyper-