PRACTICE OBSERVED

Organising a Practice

Organising preventive medicine

D J PEREIRA GRAY

An important job for the general practitioner is providing practical preventive advice. This is true in both industrialised and less developed societies and is now recognised nationally and internationally.¹ The approach to preventive medicine may be classified according to the "model for decision making."² What are we trying to do?—aims; How can we do it?—methods; How well have we done it?—assessment.

Aims

Medical activities may be arranged in a hierarchy according to when the doctor acts (table 1). Most medical work has traditionally taken place at levels (4) to (8). General practitioners are slowly learning to perform at all nine levels at once and to pay more attention to (1) and (2).

The main block to organising preventive care lies in the doctor's mind: once there is a will it is not difficult to find a way. Doctors need not feel swamped by preventive medicine. There

are relatively few interventions that have been validated and are effective, and most of these are quick and simple to do. The only facts about my patients that I try to record in writing are given in table 11 (these are my personal aims).

TABLE 11—Facts recorded about patients

Children	Adults
Head circumference at birth	Smoking habits
Abduction of hips	± cigarette smoking
Presence of testicles in scrotum (boys)	Cervical smears—sexually active within previous five years
Absence of 3rd molar or squarer	Blood pressure—adults 10-64 recorded within five years
Weight	Rubella status (women)
Absence of talipes	Contraception (all offered)
Immunisation status	(a) accepted—pill
Measles	—intruterine contraceptive device
Diphtheria	—sterilisation
Polio	
Polio	
Polio (girls after age 10 years)	

TABLE 1—Medical activities

- (1) Prevention of disease
- (2) Presymptomatic detection of disease
- (3) Early diagnosis of disease
- (4) Diagnosis of established disease
- (5) Management of disease
- (6) Management of the complications of disease
- (7) Rehabilitation after active treatment has been completed
- (8) Terminal care
- (9) Counselling the bereaved

Department of General Practice, Postgraduate Medical School, Bristol.
D J PEREIRA GRAY, MB, FRGP, general practitioner and senior lecturer in charge

How to do it (methods)

Once a general practitioner has decided to try seriously to provide practical preventive care for his or her patients the next step is to find a way of doing it. There are several approaches, but the following six are particularly useful.

710

WORK AS A TEAM

Practical preventive care for over 2000 patients cannot be done by the doctor alone, but requires a team approach from the doctor, health visitor, practice sister, and secretary. Health visitors already have a long and distinguished history in promoting preventive care in the community. Teams need time for discussion, planning, and analysing. They work best with a room for each member, and if there is no practice sister working in a treatment room a comprehensive preventive programme is very hard to sustain. Each team member needs to know the role of the others, and all should meet together periodically to review the results for their immunisation and attendance rates.

REGARD THE LIST OF PATIENTS AS A POPULATION

General practitioners have for decades been content to react to the problems of patients and to think of only one patient at a time. By regarding patients as a population at risk for whom they are responsible it becomes possible for family doctors to practise miniprevention. General practitioner trainers have a special responsibility to show trainees this concept. A simple way of doing this is by using an age-sex register. This is now essential for informing the doctor, for instance, exactly how many children or how many women aged from 15 to 44 years he is responsible for.

In 1977 an editorial in the *Journal of the Royal College of General Practitioners* commented: "A practice which either does not have or does not use an age-sex register is indicating that it is not particularly interested in preventive medicine or the surveillance of age-sex groups among its patients."³ Much of this information is easily transferred to computers, and the RCGP⁴ predicted that computers will be in common use in general practice by 1985.

TAKE THE INITIATIVE

Patients in the future may come knocking on the surgery door asking for preventive care, and they should be encouraged to do so. Meanwhile the responsibility to take the initiative lies with the primary health care team. This can be done in two ways. The first is to offer preventive care to groups of patients selected by age and sex. The simplest practical application is to send for 10-year-old girls for their 10th birthday. A reasoned letter can be sent from the doctor with a rubella leaflet (available free from the Health Education Council).

Secondly, doctors and nurses may use ordinary consultations to promote preventive care as well as special sessions or clinics, such as ante-natal clinics, which have long been established, and newer clinics, such as a children's clinic or a well-woman clinic.

The real challenge, however, is to integrate preventive care with therapeutic care in the same place, with the same person, at the same time. This new responsibility for general practitioners is well encapsulated by Stott and Davis⁵ in their theoretical model of the consultation in general practice.

They introduced this as a method of teaching, but it is even more important as an aide-memoire for the practical general practitioner.

Since on average 64% of all patients consult a general practitioner in Britain each year and as many as 95% within five years,⁶ the possibility of reaching virtually the whole population through case finding already exists.

USE PREVENTIVE MEDICINE RECORD CARDS

The absence of special record systems may have been the biggest practical block to the mass application of preventive

The exceptional potential in each primary care consultation—an aide-memoire

A	B
Management of presenting problems	Modification of help-seeking behaviour
C	D
Management of continuing problems	Opportunistic health promotion

care in general practice. A system is needed which reminds the doctor quickly during ordinary consultations of the preventive health status of the patient. One practical way is to use separate coloured summary cards. Brief information like "smokes 20/day" or "sterilised" may be recorded. In the future, separate preventive health cards may come into general use. Meanwhile coloured summary sheets are available to general practitioners free of charge on application to family practitioner committees.

A system is also needed for recording serial changes over time—for example, a child's weight in relation to percentiles. Special record cards for children have been available for years. Scaled down copies of percentile charts may be used, or the child record card, which fits the National Health Service and is available from the RCGP Birmingham Research Unit.

ENCOURAGE PERSONAL LISTS

Personal lists are not a requirement for promoting preventive medicine, but are emerging as an aid. Compliance is improved by personal lists,⁷ and this is likely to be even more important for problems of excessive smoking, eating, and drinking. If a general practitioner has an individual personal list and an age-sex register it is much easier for him to take an interest in his own immunisation rates and to take a personal pride in returns that improve over the years.

EFFICIENT BUSINESS MANAGEMENT

General practitioners are managers. They own premises, employ staff, and are directly responsible for expenditure. If practices are to take on substantial new management roles then the principals, who are responsible, must be clear about the additional costs in staff, stationery, and postage, and how these are to be recovered. Taking the initiative in practising preventive medicine, especially for age groups of patients, is expensive.

Jones et al⁸ have recently examined the financial incentives available to general practitioners for preventive medicine—for example, the immunisation fee of £2.50 (higher rate). They conclude: "We believe the treatment-room should pay its way."⁸ "An efficient nursing team should generate enough item-of-service fees to cover at the 30% of their salaries which is not directly reimbursed by the NHS."

Assessment

Thorough case assessment or audit. The modern general practitioner accepts the professional responsibility of knowing what is going on in his own practice⁹⁻¹¹—that is, measuring or counting achievements.

What percentage of children are immunised at a defined age? What percentage of girls have been immunised by the practice against rubella by their 12th birthday?

What percentage of adult patients have their smoking habits recorded in writing?
How many of the smokers have been given advice and a leaflet?¹²
What percentage of boys under 10 years have the presence of both testicles recorded in writing?
What percentage of women aged 15 to 44 have been (a) offered, (b) accepted contraceptive advice?
What percentage of women aged 15 to 44 have had a cervical smear and when?
What percentage of adults aged 35 to 64 have had their blood pressure recorded¹³ within the previous three years?
Each general practitioner can create his or her own list.

I would like to acknowledge my partners, Dr Ann Buxton, Dr Russell Steele, and Dr Margaret Mitchell, with whom these policies have been worked out in the practice. Dr Mollie Donohoe, Dr Roger Peppatt, and my wife have made valuable suggestions.

Practice Research

Variations of plasma potassium concentrations during long-term treatment of hypertension with diuretics without potassium supplements

FRANCIS F SANDOR, PETER T PICKENS, JOYCE CRALLAN

All the diuretics commonly used to treat patients with hypertension, such as the thiazides and related drugs including chlorothalidone, frusemide, and bumetanide, increase potassium excretion. Fearing that hypokalaemia may occur with or without the loss of total body potassium, many doctors prescribe potassium supplements routinely to patients with hypertension. MIMS lists eight potassium supplements in addition to the preparations listed in the *British National Formulary* and nine combinations of a thiazide or similar drug and a potassium supplement. Long-term studies in hospital have shown that appreciable potassium depletion occurring with hypokalaemia is uncommon,¹⁻³ and the danger of taking potassium supplements has been pointed out.⁴⁻¹¹ No comparable study has been done in general practice, where most hypertensive patients are managed in Britain. These include most of the patients with mild to moderate hypertension who take small doses of drugs, lead a normal working life, and take their normal diet, which in Britain contains far more than the minimum requirement of potassium. On the other hand, a wide variety of diuretics and other antihypertensive drugs is prescribed, compliance is probably not as

good as in hospital, and laboratory tests are probably performed less frequently and less uniformly than for patients serviced in hospital. Thus if hypokalaemia or potassium deficiency occurs it may go undetected for a long time. We therefore decided to study the incidence and severity of hypokalaemia and potassium depletion in a large group of hypertensive patients managed in general practice to assess whether potassium supplementation was necessary.

Methods

This study was planned with the collaboration of all general practitioners in the Hartlepool area, which has a stable population. Patients were eligible for entry if they had been treated or were newly diagnosed, but the following were excluded because they were judged to be susceptible to potassium depletion:

- (1) Patients aged over 65, who often take insufficient food.¹² Patients were admitted to the study only if a qualitative assessment of their diet showed that it was adequate.
- (2) Patients who had oedema, whether cardiac, hepatic, or renal.
- (3) Patients who were receiving cardiac glycosides.

Patients with signs or a history of myocardial infarction or cerebrovascular accident were entered, provided that their general condition was good and there had been no cardiac failure, as were patients with renal failure: hypokalaemia may be a risk when such patients receive potassium supplements.¹³ One patient who was known to have lymphatic leukaemia was entered.

One hundred and sixty-nine patients were entered. Of these, 158 were followed up at least to the first assessment: 89 were men aged 22-76 (mean 54), weighing 51-101 kg (mean 78 kg); 69 were women

Hartlepool Postgraduate Medical Centre, Hartlepool, Cleveland TS24 8AQ

FRANCIS F SANDOR, FRCP, consultant physician, General Hospital, Hartlepool
PETER T PICKENS, FRCP, consultant physician, General Hospital, Hartlepool
JOYCE CRALLAN, AMLT, senior scientific officer

712

aged 28-70 (mean 56), weighing 38-94 kg (mean 65 kg). Doctors entered their patients to the trial directly from general practice or through referral to the outpatient clinic of Hartlepool General Hospital, where they were assessed by FIT. The choice of antihypertensive drugs was left to the general practitioner. The only condition of entry into the trial was that the patient received a potassium-sparing diuretic that was not combined with any potassium-sparing diuretic. For later analysis patients were divided into four groups according to treatment before entering the trial:

Group	Diuretic treatment
A (n=87)	Diuretic without potassium supplements*
B (n=14)	Diuretic without potassium supplements*
C (n=31)	Diuretic with potassium supplements
D (n=66)	No treatment

*With or without other antihypertensive drug

After entering the trial all patients took diuretics without potassium supplements. General practitioners were free to withdraw diuretic drugs or add potassium supplements, or both, if hypokalaemia or other side effects of diuretic treatment supervened, but were asked to keep the patients on diuretics throughout the trial if they were well tolerated.

At entry and at three-monthly intervals the following were measured: height; weight; concentrations of plasma potassium, sodium, chloride, bicarbonate, urea, and creatinine by standard Technicon Autoanalyser techniques; whole blood potassium concentrations by lysate of blood by saponin and measurement of the lysate on a Technicon Autoanalyser; and haematocrit value by Coulter 1 counter. Some patients volunteered for a study of whole-body potassium by shadow-shield whole-body scintillation counter.

To avoid the fluctuations of plasma potassium concentrations with meals and diurnal variations all patients reported at 9 am after a 12-hour fast. All blood samples were taken by two people, who avoided contracting patients' muscles and applying tourniquets. Plasma was immediately separated by centrifugation and transferred within minutes to the Technicon SMA Autoanalyser. For analysis, hypokalaemia was subdivided into grade 1: 3.0-3.3 mmol(mEq)/l (<3.4 mmol(mEq)/l) and grade 2: 2.6-2.9 mmol(mEq)/l (<3.0 mmol(mEq)/l). Most other authors have chosen 3.5 mmol(mEq)/l as the lower limit of normal.¹⁴⁻¹⁶ A few 2.4-2.7-3.0 to indicate definite hypokalaemia.¹⁴⁻¹⁶ Groups A and D were compared by using Student's *t* test, and changes in each group with time were evaluated with paired *t* tests.

Results

INITIAL PLASMA POTASSIUM CONCENTRATIONS

There was a wide scatter of plasma potassium concentrations in all groups, but the mean concentration was highest in group C (previous treatment with antihypertensive drugs excluding diuretics) and lowest in group B (previous treatment with diuretics without potassium supplements) (table 1).

PLASMA POTASSIUM CONCENTRATIONS DURING FOLLOW-UP

Table 11 shows the results for the 158 patients at three months and for fewer patients at one and two years. Plasma potassium concentrations over the first three months in groups A, B, and C over the first three months and in group D over the first three months, and at one year and two years differences between the groups had disappeared. Previous treatment influenced the results over about three months only, as all four groups were combined for subsequent analyses (table 11). After the fall in the mean plasma potassium concentrations in the first three months there was no further mean fall. The proportion of patients with mild hypokalaemia increased over the first three months but thereafter remained almost constant. The small number with more severe hypokalaemia (grade 2) was almost constant throughout the study. Paired samples analysis in table IV confirm the same trends. Plasma potassium concentrations fell slightly in the first three months, then stabilised or rose slightly over the next 21 months.

In individuals plasma potassium concentrations fluctuated throughout the study despite eliminating variations in meals and diurnal variations. In several patients plasma potassium estimation was repeated over periods of four to 42 days, and the differences between paired samples varied from +0.7 to -0.7 mmol(mEq)/l. The plasma

TABLE 1—Mean and range of initial plasma potassium concentrations (mmol (mEq)/l) in the four groups

	A	B	C	D
No. of patients	47	14	31	66
Mean	3.8	3.7	3.9	3.8
Range	3.1-4.5	2.6-4.3	3.2-5.0	2.4-5.4
No. of patients with hypokalaemia	2 (4.3%)	2 (14.3%)	2 (6.5%)	2 (3.0%)
Grade 1—mild	2 (4.3%)	2 (14.3%)	2 (6.5%)	2 (3.0%)
Grade 2—moderate	0	0	0	0

Group C vs group A: *p* < 0.02.
Group C vs group B: *p* < 0.001.

TABLE 11—Mean and range of plasma potassium concentrations (mmol(mEq)/l) after three months, one year, and two years in groups A, B, C, and D

Assessment	A	B	C	D
Initial	47	14	31	66
3 months	3.8	3.7	3.9	3.8
1 year	3.8	3.7	3.9	3.8
2 years	3.8	3.7	3.9	3.8
No. of patients	34	10	28	57
Mean	3.6	3.3	3.7	3.7
Range	2.9-4.7	2.4-4.1	3.0-5.1	2.7-4.8
1 year	32	9	23	39
Mean	3.7	3.5	3.7	3.6
Range	2.9-4.5	2.4-4.4	3.1-4.5	2.7-4.4
2 years	13	4	11	13
Mean	3.7	3.6	3.8	3.6
Range	3.2-4.4	3.0-4.0	2.9-4.3	2.9-4.5

TABLE III—Mean and range of plasma potassium concentrations (mmol(mEq)/l) for all the patients initially and after three months, one year, and two years

	Initial assessment	After 3 months	After 1 year	After 2 years
No. of patients	158	124	100	41
Mean	3.8	3.7	3.7	3.6
Range	2.6-5.0	2.7-5.3	2.7-5.4	2.4-5.4
No. of patients with hypokalaemia	9 (5.7%)	21 (17%)	18 (18%)	12 (29%)
Grade 1—mild	9 (5.7%)	21 (17%)	18 (18%)	12 (29%)
Grade 2—moderate	0	0	0	0

TABLE IV—Mean and range of plasma potassium concentrations (mmol(mEq)/l) between paired samples in the four groups over two years

Assessment	A	B	C	D
Initial to 3 months	3.8	3.7	3.9	3.8
Mean	3.8	3.7	3.9	3.8
Range	-1.2 to +1.3	-0.4 to +0.6	-1.1 to +1.3	-1.1 to +1.3
Initial to 1 year	3.8	3.7	3.9	3.8
Mean	3.8	3.7	3.9	3.8
Range	-0.4 to +1.1	-0.6 to +0.8	-0.8 to +1.3	-0.4 to +1.1
Initial to 2 years	3.8	3.7	3.9	3.8
Mean	3.8	3.7	3.9	3.8
Range	-0.5 to +1.1	-0.2 to +0.6	-0.6 to +1.7	-0.4 to +1.1
2 months to 1 year	3.8	3.7	3.9	3.8
Mean	3.8	3.7	3.9	3.8
Range	-0.5 to +1.1	-0.2 to +0.6	-0.6 to +1.7	-0.4 to +1.1
2 months to 2 years	3.8	3.7	3.9	3.8
Mean	3.8	3.7	3.9	3.8
Range	-0.5 to +1.1	-0.2 to +0.6	-0.6 to +1.7	-0.4 to +1.1
1 year to 2 years	3.8	3.7	3.9	3.8
Mean	3.8	3.7	3.9	3.8
Range	-0.5 to +1.1	-0.2 to +0.6	-0.6 to +1.7	-0.4 to +1.1

*Significant difference of at least *p* < 0.05.
In the paired samples analysis of plasma potassium concentrations (groups A and B) the difference between the mean potassium concentrations at three months, confirming the rise of plasma potassium concentration over the first three months, was significant only in the first three months, indicating that the rise was sustained. In the patients who started diuretics only at the beginning of the trial (groups C and D) the difference remained significant at one year.