Hypophosphataemia after parathyroidectomy in chronic renal failure

K FARRINGTON, Z VARGHESE, R A BAILLOD, O N FERNANDO, J F MOORHEAD

Abstract

Out of 24 patients receiving haemodialysis who were subjected to parathyroidectomy, 13 developed hypophosphataemia; this persisted for 3-52 weeks (mean 10.6 weeks). Before operation these 13 patients had had significantly higher plasma alkaline phosphatase activities ($p < 0.01$) and significantly higher values in iliac crest bone biopsy samples for active resorption surface and active formation surface ($p < 0.05$ in each case) than the group who remained normophosphataemic. Significantly more of the patients who remained normophosphataemic had shown periarticular calcification in preoperative skeletal radiographs ($p < 0.001$).

Hypophosphataemia may result from reduced mobilisation of phosphate from bone or its increased accretion into bone, and resorption of phosphate from periarticular mineral deposits may protect against development.

Introduction

Serum phosphate concentrations often fall after parathyroidectomy in patients with end-stage renal failure,¹⁻⁶ but frank hypophosphataemia is rarely reported.^{7,8} Hypophosphataemia in patients receiving haemodialysis may contribute to osteomalacia,⁹ and the importance of identifying hypophosphataemia in these patients after parathyroidectomy has been emphasised.⁸ We observed hypophosphataemia after parathyroidectomy in 13 out of 24 patients receiving haemodialysis. We therefore examined clinical, biochemical, radiological, and histological variables to try to identify factors contributing to hypophosphataemia.

Patients and methods

The study group comprised all 24 patients receiving haemodialysis (13 male, 11 female) who underwent parathyroidectomy at this hospital between 1973 and 1980. Their ages ranged from 9 to 55 years (mean $35 \pm \text{SD } 13$ years), and the duration of dialysis was 16-180 months (mean 71 ± 43 months). All patients had had treatment with phosphate-binding antacids and 16 with vitamin D compounds. In most cases these had been discontinued many months before parathyroidectomy because of hypercalcaemia. Four patients had also been given courses of human calcitonin.¹⁰ The main indication for parathyroidectomy was severe bone disease associated with hypercalcaemia.

Plasma alkaline phosphatase activity and calcium, phosphate, and hydroxyproline concentrations were measured in all patients as described.¹¹ Immunoreactive parathyroid hormone was measured in 21 patients as described.^{10,12} Preoperative skeletal surveys were performed in all patients and the radiographs examined for subperiosteal erosions, osteosclerosis, vascular calcification, and periarticular calci-

fication. Iliac crest bone biopsy specimens were obtained preoperatively in 17 patients and processed and analysed as described.¹³ Variables measured were total bone volume (% of total tissue), osteoid volume (% of total bone), active resorption surface (% of total bone surface), and active formation surface (% of total bone surface). Active resorption surface is the proportion of the total bone surface covered by active osteoclasts, and active formation surface is the length of osteoid surface covered by active osteoblasts expressed as a percentage of total bone surface.

Table I shows the types of parathyroidectomies performed.

TABLE I—Types of parathyroidectomy performed in 24 patients receiving haemodialysis

	Subtotal (3 1/4)	Total				Total + auto-transplantation
		5/5	4/4	3/3	2/2	
No of patients	1	1	9	1	1	11

All the glands removed showed histological evidence of parathyroid hyperplasia. Plasma calcium and phosphate concentrations were checked at least daily for 10 days after operation, and thereafter estimations were determined by clinical need. A group of patients developing hypophosphataemia was identified and compared with the group who remained normophosphataemic with respect to clinical, biochemical, radiological, and histological variables. Oral calcium supplements (calcium carbonate) and vitamin D analogues (dihydroxycholesterol or 1α -hydroxycholecalciferol) were given in the post-operative period to maintain plasma calcium values. Oral phosphate-binding antacids were discontinued at the time of operation. Patients who developed hypophosphataemia were given phosphate supplements during dialysis by means of phosphate infusions into the dialysis fluid.

Statistical methods used were Wilcoxon's rank sum test and the χ^2 test with Yates's correction.

Results

Thirteen patients developed hypophosphataemia after parathyroidectomy, and the condition persisted for 3-52 weeks (mean $10.6 \pm \text{SD } 12.4$ weeks). The figure shows the mean serial estimations of plasma calcium and phosphate concentrations in the patients who developed hypophosphataemia and those who remained normophosphataemic. In both groups there was an immediate fall in plasma calcium and phosphate values after parathyroidectomy. The hypocalcaemia was more profound in the group who developed hypophosphataemia, and in this group the duration of hypophosphataemia was far greater than the duration of hypocalcaemia.

The two groups showed no significant differences in age, sex, duration of dialysis, or number who had received previous treatment with vitamin D compounds (table II). Nor was there any appreciable difference between the groups in the dialysis schedule after parathyroidectomy or type of operation performed. Though preoperative plasma calcium, phosphate, hydroxyproline, and immunoreactive parathyroid hormone concentrations were higher in the hypophosphataemic group, the differences were not statistically significant. Plasma alkaline phosphatase activity was significantly greater ($p < 0.01$) in the hypophosphataemic group (table II).

Table III shows the radiological findings in the two groups before operation. There was a higher incidence of subperiosteal erosions, osteosclerosis, and vascular calcification in the hypophosphataemic group, but the differences were not statistically significant. The incidence of periarticular calcification, however, was significantly higher

Department of Nephrology and Transplantation, Royal Free Hospital, London NW3 2QG

K FARRINGTON, BSC, MRCP, senior registrar
Z VARGHESE, MSc, PhD, principal biochemist
R A BAILLOD, MB, BCH, first assistant
O N FERNANDO, FRCS, consultant surgeon
J F MOORHEAD, FRCP, director of department

TABLE II—Preoperative clinical and biochemical details of patients receiving haemodialysis who were subjected to parathyroidectomy (values are means $\pm$ SD)

	No of patients	Sex		Age (years)	Duration of dialysis (months)	No with past vitamin D treatment	Calcium (mmol/l)	Phosphate (mmol/l)	Alkaline phosphatase (KAU/l)	Hydroxyproline (μ mol/l)	Immunoreactive parathyroid hormone (ng/l)
		M	F								
Patients developing hypophosphataemia	13	6	7	34 $\pm$ 13	72 $\pm$ 39	10	2.82 $\pm$ 0.16	2.16 $\pm$ 0.48	520 $\pm$ 310*	165.5 $\pm$ 120.5	3600 $\pm$ 2227
Patients remaining normophosphataemic	11	7	4	35 $\pm$ 13	69 $\pm$ 46	6	2.78 $\pm$ 0.22	1.87 $\pm$ 0.25	170 $\pm$ 100	93.1 $\pm$ 70.9	2922 $\pm$ 1374
Normal range $\pm$ 1SD							2.43 $\pm$ 0.10	1.16 $\pm$ 0.13	60 $\pm$ 20	12.2 $\pm$ 4.6	541 $\pm$ 204

Conversion: SI to traditional units—Calcium: 1 mmol/l $\approx$ 4.0 mg/100 ml. Phosphate: 1 mmol/l $\approx$ 3.1 mg/100 ml. Hydroxyproline: 1 μ mol/l $\approx$ 13.1 μ g/100 ml.

* $p < 0.01$.

Serial mean postoperative plasma calcium and phosphate concentrations in patients who developed hypophosphataemia (■) and those who remained normophosphataemic (○). Bars indicate SEM. Shaded area represents normal range.

Conversion: SI to traditional units—Phosphate: 1 mmol/l $\approx$ 3.1 mg/100 ml. Calcium: 1 mmol/l $\approx$ 4.0 mg/100 ml.

TABLE III—Preoperative radiological findings in patients receiving haemodialysis who were subjected to parathyroidectomy

	Patients developing hypophosphataemia (n = 13)	Patients remaining normophosphataemic (n = 11)
Subperiosteal erosions	13	8
Osteosclerosis	11	5
Vascular calcification	6	3
Periarticular calcification	0	6*
Looser's zones	1	0

* $p < 0.001$.

TABLE IV—Preoperative bone histomorphometric findings in patients receiving haemodialysis who were subjected to parathyroidectomy (values are means $\pm$ SEM)

	No of biopsies	Total bone volume (% total tissue)	Osteoid volume (% total bone)	Active resorption surface (% total bone surface)	Active formation surface (% total bone surface)
Patients developing hypophosphataemia	9	41.9 $\pm$ 3.9	13.5 $\pm$ 1.5	11.2* $\pm$ 2.0	10.0* $\pm$ 3.0
Patients remaining normophosphataemic	8	34.6 $\pm$ 3.4	10.7 $\pm$ 2.3	6.8 $\pm$ 0.9	2.7 $\pm$ 1.2
Normal range		18-25	< 2.8	0.1-0.8	2-8

* $p < 0.05$.

in the normophosphataemic group ($\chi^2 = 12.5$; $p < 0.001$). There were also differences between the groups in the bone biopsy findings (table IV). Total bone volume and osteoid volume were higher in the hypophosphataemic group, though the differences were not significant. Values for active resorption surface and active formation surface were significantly higher in the hypophosphataemic group ($p < 0.05$ in each case).

Discussion

Though a fall in the plasma phosphate concentration is common after parathyroidectomy in patients with end-stage renal failure,¹⁻⁶ there are few reports of frank hypophosphataemia.⁷⁻⁸ Prolonged hypophosphataemia after parathyroidectomy was noted in three patients,⁸ though two of these were taking anti-convulsant drugs; these compounds interfere with vitamin D metabolism,¹⁴ so that a true reflection of the effect of parathyroidectomy on phosphate metabolism in chronic renal failure could not be obtained. We report the development of hypophosphataemia immediately after parathyroidectomy in 13 out of 24 patients receiving haemodialysis. The mean duration of the hypophosphataemia was 10.6 weeks, though in one patient hypophosphataemia persisted for 52 weeks.

There are several possible causes for the decrease in plasma phosphate in these patients. Johnson *et al*³ thought that it was due to a reduction in the mobilisation of phosphate from bone under the influence of reduced immunoreactive parathyroid hormone concentrations. Alternatively increased accretion of phosphate into healing bone may occur. In patients developing hypophosphataemia while receiving dialysis other factors may be important, including phosphate-binding antacids, low dietary intake of phosphate,⁹ repeated dialysis with phosphate-free dialysate,⁹ and reduced absorption of phosphate due to oral calcium supplements.¹

We found that patients developing hypophosphataemia after parathyroidectomy had significantly greater preoperative plasma alkaline phosphatase activities, and their plasma phosphate and hydroxyproline concentrations were also substantially higher, but not significantly so (table II). Hypophosphataemic patients also had greater histological indices of bone turnover, having significantly higher values for active resorption surfaces and active formation surfaces than the normophosphataemic group (table IV). A higher proportion of hypophosphataemic patients also had radiological erosions and osteosclerosis (table III). These data indicate that hypophosphataemia tends to occur in patients with more severe hyperparathyroid bone disease and

hence that the major cause of hypophosphataemia is likely to be either reduced mobilisation of phosphate from bone or its increased accretion into bone.

Other factors are probably less important. Since phosphate binders were routinely stopped before operation these cannot have contributed to the postoperative reduction in plasma phosphate concentrations, and the temporal relation of the fall in these values with parathyroidectomy tends to exclude deficiencies in dietary phosphate as a possible cause. Phosphate depletion from excessive dialysis was also unlikely to account for the hypophosphataemia, since the dialysis schedules of the hypophosphataemic and normophosphataemic groups were not appreciably different. Patients who developed hypophosphataemia, however, were also those who had the most profound hypocalcaemia (figure), who would therefore tend to require higher amounts of oral calcium supplements; hence intestinal malabsorption of phosphate due to oral calcium supplements might be a contributory factor.

A significantly high proportion of patients who remained normophosphataemic after operation had periarticular calcification in their preoperative skeletal radiographs (table III). In such patients serum calcium concentrations tend to remain near normal after parathyroidectomy,¹⁵ probably owing to auto-transfusion of calcium from these deposits.⁵ A similar mechanism may be responsible for the maintenance of normophosphataemia in these patients.

Phosphate is considered to be an important modulator of bone mineralisation,¹⁶ and a significant inverse correlation exists between plasma phosphate values and histological indices of osteomalacia in patients with chronic renal failure.¹⁷ Severe osteomalacia has also been described in patients with hypophosphataemia receiving haemodialysis.⁹ Hypophosphataemia after parathyroidectomy may lead to delayed healing of bone lesions,⁵ and a vitamin-D-resistant osteomalacia has been reported after parathyroidectomy¹⁸ to which hypophosphataemia may have contributed. We found that hypophosphataemia may be predicted from biochemical, radiological, and histological criteria in patients who have severe bone disease. Careful postoperative monitoring of plasma phosphate as well as plasma calcium concentrations is necessary in these patients so that appropriate treatment may be instituted. In many of our patients adding phosphate to the dialysis fluid did not prevent persistent hypophosphataemia, and oral phosphate supplements may be necessary in these patients.

References

- 1 Stanbury SW, Lumb SA. Metabolic studies of renal osteodystrophy. *Medicine* 1962;41:1-34.
- 2 Gill G, Pallotta J, Kashgarian M, Kessner D, Epstein FH. Physiological studies in renal osteodystrophy treated by subtotal parathyroidectomy. *Am J Med* 1969;46:930-40.
- 3 Johnson JW, Wachman A, Katz AI, et al. Calcium metabolism after subtotal parathyroidectomy in chronic renal failure. *Trans Am Soc Artif Intern Organs* 1969;15:333-40.
- 4 Massry SG, Coburn JW, Popovtzer MM, Shinaberger JH, Maxwell MH, Kleeman CR. Secondary hyperparathyroidism in chronic renal failure. *Arch Intern Med* 1969;124:431-41.
- 5 David DS, ed. Mineral and bone homeostasis in renal failure. Pathophysiology and management. In: *Calcium metabolism in renal failure and nephrolithiasis*. New York: John Wiley and Sons, 1977:1-76.
- 6 Varghese Z. PhD thesis. Biochemical studies of metabolic bone disease in uraemia with special reference to dialysis and transplantation. University of London, 1979:463.
- 7 Berens JJ, Potter DJ, Nickas GM. Parathyroidectomy. Treatment for secondary hypoparathyroidism and azotemic osteodystrophy. *Arch Surg* 1970;100:338-41.
- 8 Lefavour GS, Brensilver JM, Pierce JC, Cortell S. Persistent hypophosphataemia following parathyroidectomy in end stage renal disease: report of three patients. *Clin Nephrol* 1980;13:40-3.
- 9 Ahmed KY, Varghese Z, Wills MR, et al. Persistent hypophosphataemia and osteomalacia in dialysis patients not on oral phosphate binders: response to dihydroxycholesterol therapy. *Lancet* 1976;ii:439-42.
- 10 Farrington K, Varghese Z, Moorhead JF. Human calcitonin in the treatment of renal osteodystrophy. *J Lab Clin Med* 1980;96:399-406.
- 11 Varghese Z, Moorhead JF, Tatler GLV, Baillod RA, Wills MR. Plasma hydroxyproline in renal osteodystrophy. *Proc Eur Dial Transplant Assoc* 1973;10:187-96.

- 12 Varghese Z. PhD thesis. Biochemical studies of metabolic bone disease in uraemia with special reference to dialysis and transplantation. University of London, 1979:230-2.
- 13 Meinhard E, Wadbrook DG, Ring C, et al. Computer card morphometry in uremic bone disease. In: Norman AW, Schaefer K, Grigoleit HG, von Herrath D, Ritz E, eds. *Vitamin D and problems related to uremic bone disease*. New York: Walter De Gruyter Inc, 1975:547-52.
- 14 Hahn TJ, Herdin BA, Scharp CR, Haddad JG. Effect of chronic anticonvulsant therapy on serum 25-hydroxycalciferol levels in adults. *N Engl J Med* 1972;287:900-4.
- 15 Massry SG, Coburn JW. Divalent ion metabolism and renal osteodystrophy. In: Massry SG, Sellers AL, eds. *Clinical aspects of uremia and dialysis*. Springfield, Illinois: Charles C Thomas, 1976:304-87.
- 16 Rasmussen H, Bordier P. *Physiological and cellular basis of metabolic bone disease*. Baltimore: Williams and Wilkins, 1974:347-9.
- 17 Kanis JA, Henderson RG, Heynen G, et al. Vitamin D, osteomalacia, and chronic renal failure. In: Norman AW, Schaefer K, von Herrath D, et al, eds. *Vitamin D, biochemical, chemical and clinical aspects related to calcium metabolism*. Berlin, New York: Walter De Gruyter, 1977:671-3.
- 18 Coburn JW, Brickman M, Sherrard DJ, et al. Clinical efficiency of 1,25-dihydroxy vitamin D₃ in renal osteodystrophy. In: Norman AW, Schaefer K, von Herrath D, et al, eds. *Vitamin D, biochemical, chemical and clinical aspects related to calcium metabolism*. Berlin, New York: Walter de Gruyter, 1977:657-66.

(Accepted 11 December 1981)

ONE HUNDRED YEARS AGO The following account of the Montefiore medal and prizes, which have been given for the first time at the late session of the Army Medical School, at Netley, was read by Professor Longmore, at the closing meeting on February 6th. In the spring of the year 1881, Mr Nathaniel Montefiore, FRCS, having observed that no prize was given at the Army Medical School in the department of military surgery, although prizes were awarded in the sections of hygiene and military medicine, and being desirous of stimulating practical study and improvement in this branch of professional duties of military surgeons, offered to fund a sum of two thousand pounds for the purpose of establishing a prize to be competed for in the military surgery class at the school. This generous offer was submitted through the Director-General of the Army Medical Department for the consideration of the Right Honourable the Secretary of State for War; and the Secretary of State was pleased to accept the offer, and expressed to Mr Montefiore, by letter, his appreciation of the patriotic motives which had prompted the gift, and his gratification in the acceptance of it. Mr Montefiore left the arrangements for the disposal of the fund to the governing body of the army medical school. After due consideration, it was determined to appropriate the proceeds of the fund to two prizes: the first to consist of a bronze medal and sum of money; the second, of a selection of books relating to matters connected with military surgery. The subjects to be engraved on the proposed medal were arranged and designed at Netley. One face of the medal is intended to illustrate a military surgeon's duty in attending to wounded men in the field, and to represent some of the principal articles of modern field-hospital equipment; the other face bears the arms and name of the founder of the prize, and the date of its foundation. As soon as the drawings for the medal were completed, the work was submitted to a leading English medallist for execution; but the designs, as proposed, being declared unsuited for medallic effect, it was then offered to an eminent French medallist, M Dubois, of Paris, who both expressed his approval of the plan, and at once offered to undertake the execution of it. The plaster cast of the model of the principal face of the medal, triple the size of the intended medal itself, has been recently received from M Dubois, and has been regarded, by all who have seen it, as most artistic and satisfactory. In order that no delay might occur in giving the prize, Mr Montefiore has, with still further liberality, undertaken to defray the cost of the medal. It would have taken more than a year and a half to pay for it, if its cost had to be met by the interest accruing from the sum originally funded by Mr Montefiore. This has enabled the prizes for military surgery to be given during the present session. The first prize this session consists of the medal and twenty guineas; the second, of four books. As the medal is not yet completed, it will be forwarded to Mr James Barry Gibbons, of Her Majesty's Indian service, who has gained it, as soon as it is received from Paris, and the successful competitor's name has been engraved on it. (*British Medical Journal*, 1882.)