of the alimentary tract was intermittent or more localised than we explored, similar to the localised tender segments in the oesophagus described by Edwards.²⁰ Alternatively, pain may arise in other areas such as the pancreas or biliary tract or in some cases from a non-alimentary source.

Referral of pain to extra-abdominal sites has now been shown from the small as well as the large bowel. A wider appreciation of the protean presentation of functional abdominal pain and the existence of potential tender areas throughout the whole gut should expedite the recognition of non-organic abdominal pain and prevent unnecessary investigations and even exploratory laparotomy.

KJM is in receipt of a Medical Research Council training fellowship.

References

- Isselbacher KJ. Indigestion. In: Isselbacher KJ, Adams RD, Braunwald EB, Petersdorf RG, Wilson JD, eds. *Harrison's principles of internal medicine*. 9th ed. Maidenhead: McGraw-Hill, 1980:195.
- Hawkins CF. The alimentary system. In: Mann WN, ed. *Conybeare's textbook of medicine*. 16th ed. Edinburgh: Churchill Livingstone, 1977:285.
- Way LW. Abdominal pain and the acute abdomen. In: Sleisenger MH, Fordtran JS, eds. *Gastrointestinal disease*. 2nd ed. Eastbourne: W B Saunders, 1978:398-404.
- Rains AJH, Ritchie HD, eds. The intestines. In: *Bailey and Love's short practice of surgery*. 18th ed. London: Lewis, 1981:990.
- Schwartz SI, Shires GT, Spencer FC, Storer EH, eds. Manifestations of gastrointestinal disease. In: *Principles of surgery*. 3rd ed. Maidenhead: McGraw-Hill, 1979:1043.
- Jones CM. *Digestive tract pain*. New York: Macmillan, 1938:8.
- Bentley FH, Smithwick RH. Visceral pain produced by balloon distension of the jejunum. *Lancet* 1940;ii:389.
- Ray BS, Neill CL. Abdominal visceral sensation in man. *Ann Surg* 1947;126:709-24.
- Chapman WP, Jones CN. Variations in cutaneous and visceral pain sensitivity in normal subjects. *J Clin Invest* 1944;23:81-91.
- Waller SL, Misiewicz JJ. Prognosis in the irritable-bowel syndrome. *Lancet* 1969;ii:753-6.
- Manning AP, Thompson WG, Heaton KW, Morris AF. Towards positive diagnosis of the irritable bowel. *Br Med J* 1978;iii:653-4.
- Swarbrick ET, Hegarty JE, Bar L, Williams CB, Dawson AM. Site of pain from the irritable bowel. *Lancet* 1980;ii:443-6.
- Connell AM, Hilton C, Irvine G, Lennard-Jones JE, Misiewicz JJ. Variation of bowel habit in two population samples. *Br Med J* 1965;ii:1095-9.
- Hurst AF. On the sensibility of the alimentary canal in health and disease. *Lancet* 1911;ii:1051, 1119, 1187.
- Lennander KG. Über die Sensibilität der Bauchhöhle und über lokale und allgemeine Anästhesie bei bruchund Bauchoperationen. *Zentralbl Chir* 1901;28:209.
- Ryle JA. Visceral pain and referred pain. *Lancet* 1926;ii:895.
- Payne WW, Poulton EP. Visceral pain in the upper alimentary tract. *Q J Med* 1923;17:53.
- Payne WW, Poulton EP. Experiments on visceral sensation. *J Physiol* 1927;63:217.
- Morley J. Visceral pain. *Br Med J* 1937;ii:1270.
- Alvarez WC. Abdominal pain: the sensitive regions of the abdomen and ways in which they may be stimulated to produce pain. *JAMA* 1934;102:1351.
- Dworkin HJ, Biel FJ, Machella TE. Supradiaphragmatic reference of pain from the colon. *Gastroenterology* 1952;22:222-9.
- Thompson WG, Heaton KW. Functional bowel disorders in apparently healthy people. *Gastroenterology* 1980;79:283-8.
- Holdstock DJ, Misiewicz JJ, Waller SL. Observations on the mechanism of abdominal pain. *Gut* 1969;10:19-31.
- Connell AM, Jones FA, Rowlands EN. Motility of the pelvic colon. IV. Abdominal pain associated with colonic hypermotility after meals. *Gut* 1965;6:105-12.
- Horowitz L, Farrar JT. Intraluminal small intestinal pressures in normal patients and in patients with functional gastrointestinal disorders. *Gastroenterology* 1962;42:455-64.
- Kalser MH, Zion DE, Bockus HL. Functional diarrhoea: an analysis of the clinical and roentgen manifestations. *Gastroenterology* 1956;31:629-46.
- Thompson DG, Laidlow JM, Wingate DL. Abnormal small-bowel motility demonstrated by radiotelemetry in a patient with irritable colon. *Lancet* 1979;ii:1321-3.
- Sekar ASC, Thompson WG, Turner LJ, Barr R. Duodeno-gastric reflux and gastritis in non-ulcer dyspepsia. *Clinical Research* 1977;24:667A.
- Edwards DAW. History and symptoms of oesophageal disease. Diseases of the oesophagus. In: Vantrappen G, Hellemans J, eds. *Handbuch der inneren Medizin*. Berlin, Heidelberg, New York: Springer-Verlag, 1974:112.

(Accepted 23 April 1982)

SHORT REPORTS

"Scalded mouth" caused by angiotensin-converting-enzyme inhibitors

Captopril, an angiotensin-converting-enzyme inhibitor, is used to treat resistant hypertension.¹ Side effects include rash, proteinuria, leukopenia, ageusia, angio-oedema, and aphthous and tongue ulcers; its sulphhydryl moiety may be an aetiological factor.¹ Enalapril, a new converting-enzyme inhibitor, does not contain a sulphhydryl group.² We describe three patients who reported a scalded sensation of the oral mucosa during treatment with captopril or enalapril.

Case reports

Case 1—A 53-year-old hypertensive woman receiving hydrochlorothiazide 50 mg daily was prescribed captopril 12.5 mg thrice daily in addition; she took no other medication. After two weeks the dosage of captopril was increased to 25 mg thrice daily. Six days later she complained of a burning sensation on the upper surface of her tongue, comparable to having been "scalded by a hot liquid," though she had no recollection of such an occurrence. Examination of her tongue and mouth was unremarkable. She reported no alteration in taste. With her consent the regimen was continued, the symptom persisted for five days and then dissipated despite two additional weeks of captopril treatment, without subsequent recurrence.

Case 2—During a research study a 54-year-old man with essential hypertension received enalapril 20 mg twice daily. Seven days later he said that his tongue and mouth felt as though they had been "scalded by coffee or pizza," though he could not recall such an occurrence. Examination of the mouth showed only periodontal disease and questionable hyperaemia; neurological evaluation including taste perception was normal. He took no

other medications. He continued treatment with enalapril for four additional days without resolution. The symptom abated, however, when enalapril was stopped, and he was treated with hydrochlorothiazide 50 mg/day. Two weeks later enalapril was restarted in a dose of 20 mg daily, the diuretic being continued. On the twelfth day of treatment he again complained of the "scalded-mouth" sensation, which was similar to but milder than the first episode. This continued for the next two days but abated when enalapril was stopped at the end of the trial. Similar symptomatology did not recur.

Case 3—A 64-year-old hypertensive woman taking propranolol 160 mg and hydrochlorothiazide 50 mg daily was prescribed captopril 25 mg thrice daily in addition. Other medications included nitroglycerine and isosorbide dinitrate. After two weeks the captopril dosage was increased to 50 mg thrice daily. Nine weeks later she complained of a burning sensation in her throat, palate, and tongue. Examination of the mouth was unremarkable. After two additional weeks of treatment with captopril the symptoms persisted and mild injection at the margins of the soft palate was noted; there was no alteration in taste. Because of continued discomfort captopril was stopped with a distinct improvement during the next week; slight discomfort, however, localised to the tip of her tongue, persisted.

Comment

Although disorders of taste as well as aphthous and tongue ulcers have been reported during captopril treatment,³⁻⁵ the symptom described by our patients has not, to our knowledge, been previously associated with use of either captopril or enalapril. The mechanism of this side effect is unclear. Neither fever nor eosinophilia accompanied the patients' complaints. No concomitant laboratory abnormalities were noted, and all the patients had normal renal function. Since the side effect was noted with both captopril and enalapril the aetiology of this scalded-mouth reaction appears to be independent of the sulphhydryl moiety of captopril. The hyperaesthesia of the oral mucosa abated during continued treatment with captopril in case 1, recurred

on rechallenge with enalapril in case 2, and was troublesome enough to require captopril treatment to be stopped in case 3. Doctors treating patients with converting-enzyme inhibitors should be aware of the potential occurrence of this side effect.

Requests for reprints should be addressed to: Dr Peter H Vlasses, Clinical Pharmacology (M-504), Thomas Jefferson University Hospital, 11th and Walnut Streets, Philadelphia, PA 19107, USA.

- ¹ Ferguson RK, Vlasses PH. Clinical pharmacology and therapeutic applications of the new oral angiotensin converting enzyme inhibitor, captopril. *Am Heart J* 1981;101:650-6.
- ² Gavras H, Biollaz J, Waeber B, Brunner HR, Gavras I, Davies RO. Anti-hypertensive effect of the new oral angiotensin converting enzyme inhibitor "MK-421". *Lancet* 1981;ii:543-7.
- ³ Vlasses PH, Ferguson RK. Temporary ageusia related to captopril. *Lancet* 1979;ii:526.
- ⁴ Seedat YK. Aphthous ulcers of mouth from captopril. *Lancet* 1979;iii:1297-8.
- ⁵ Nicholls MG, Maslowski AH, Ikram H, Espiner EA. Ulceration of the tongue: a complication of captopril therapy. *Ann Intern Med* 1981;94:659.

(Accepted 5 February 1982)

Division of Clinical Pharmacology, Jefferson Medical College, Thomas Jefferson University, Philadelphia, Pennsylvania, USA

PETER H VLASSES, PHARM D, associate director and clinical associate professor of medicine

HESCHI H ROTMENSCH, MD, assistant director and clinical assistant professor of medicine

ROGER K FERGUSON, MD, director and professor of medicine and pharmacology

Department of Pharmacy, Thomas Jefferson University Hospital, Philadelphia

STEVEN L SHEAFFER, PHARM D, assistant director

Prenatal serological diagnosis of intrauterine cytomegalovirus infection

Cytomegalovirus commonly causes intrauterine infection. Finding a primary or reactivated cytomegalovirus infection in the mother by isolating the virus from the throat, urine, cervix, or blood, or by maternal serology, does not necessarily indicate that the fetus will be infected. Fetal infection appears to occur in no more than 25-50% of even primary maternal infections. The presence of intrauterine infection has been established by culture of cytomegalovirus from amniotic fluid,¹ but such isolation may be time consuming. We report a case that shows that the availability of a sensitive radioimmunoassay for the detection of cytomegalovirus-specific IgM² and the ability to obtain fetal cord serum³ mean that intrauterine serological diagnosis is possible.

Case report

A 28-year-old rhesus-positive woman without atypical antibodies was referred at 25 weeks' gestation after ultrasonography showed a hydropic fetus. She had felt well throughout the pregnancy, denied taking any medication, and reported no relevant personal or family history. At 16 weeks' gestation an ultrasound scan had confirmed her dates and shown no major structural abnormalities. Examination was unremarkable, with fundal height being consistent with dates. A further ultrasound scan confirmed a single fetus with gross ascites, but no other abnormalities were observed. Fetoscopy was carried out and 8 ml pure fetal blood and 260 ml ascitic fluid collected. Fetal serum showed appreciable hypoalbuminaemia of 6 g/l (normal $21 \pm SD$ 2 g/l (unpublished)). A fetal blood film showed severe erythroblastosis.

A follow-up ultrasound scan showed re-collection of the ascites. Intra-vascular fetal albumin infusion was planned but precluded by spontaneous premature rupture of the membranes at 27 weeks' gestation. A conservative policy was adopted but labour ensued and a 1480 g severely depressed male infant delivered. Resuscitative attempts were stopped after 20 minutes. Initial serology for cytomegalovirus on maternal sera showed an antibody titre of 32 when a complement fixation test was performed on sera collected at 21 weeks' and 25 weeks' gestation. No cytomegalovirus-specific IgM was

detected in the fetal serum using indirect immunofluorescence (Electro-nucleonics Laboratories Inc, Maryland, USA).

Necropsy showed non-icteric hydrops fetalis (figure). Macroscopically the only abnormal findings were a large liver with a finely nodular and mottled surface and massive ascites causing gross elevation of the diaphragm with consequent pulmonary hypoplasia. Microscopy showed cytomegaly and intranuclear "owl's eye" inclusions in many organs, particularly the kidney (figure). The appearance was typical of cytomegalic inclusion disease. The liver showed extreme active erythropoiesis. Cytomegalovirus was isolated from urine obtained at necropsy.

Male fetus of 1480 g showing gross hydrops fetalis. Inset: kidney showing cytomegalic cells with inclusion bodies. Haematoxylin and eosin. $\times 2500$ (original magnification).

These findings prompted a re-examination of stored maternal and fetal cord sera. The total IgM content of the fetal serum was 0.11 g/l. This is within the normal range for a fetus of this gestation as stated by Alford *et al.*⁴ Radioimmunoassay for cytomegalovirus IgM showed that the fetal serum had a titre of 4000. This is within the range seen in neonatal or cord serum from confirmed cases of symptomatic congenital cytomegalovirus.⁵ Examination of the maternal serum for cytomegalovirus-specific IgM by radioimmunoassay showed a titre of 800 in the serum taken at 21 weeks' gestation, but such IgM was absent at 25 weeks. These results were compatible with a primary infection occurring early in the pregnancy.

Comment

Cytomegalovirus is a recognised cause of hydrops fetalis, and intra-uterine cytomegalovirus infection was confirmed in this case by histology and isolation of the virus. Initial serology for cytomegalovirus specific IgM on fetal cord serum obtained at 25 weeks' gestation was negative using an indirect fluorescence assay. When a sensitive radioimmunoassay was used, cytomegalovirus-specific IgM was detected with ease, showing the increased sensitivity of this procedure for diagnosing congenital cytomegalovirus.⁶ We believe this to be the first case in which virus-specific IgM has been shown in fetal serum obtained in utero.

This work was supported by the National Fund for Research into Crippling Diseases.

Requests for reprints should be addressed to CHR.

- ¹ Davis LE, Tweek GV, Stewart JA, *et al.* Cytomegalovirus mononucleosis in a first trimester pregnant female with transmission to the fetus. *Pediatrics* 1971;48:200-6.