PRACTICE OBSERVED

Practice Research

Review of maternity patients suitable for home delivery

E A DIXON

The number of home deliveries of babies has been decreasing and now only 1.4% of deliveries take place at home. Opponents of home delivery believe that the safest place for the patient to be delivered is in hospital where expert facilities and attention are immediately available for the mother and her newborn baby. Those who favour home delivery believe that the hospital environment is unfamiliar and unfriendly. They also think that the routine monitoring of the fetal heart rate, artificial rupture of membranes, oxytocin stimulation, and the widespread use of episiotomy and other procedures disrupt the family atmosphere and also cause lack of bonding between the newborn and the rest of the family.^{1,2}

There are many arguments for and against home delivery.^{3,4} and the perinatal mortality rates have been compared. Hudson,⁵ James,⁶ and Lewis⁷ et al⁸ have also discussed patients who were initially booked for delivery at home or in general practitioner obstetric units (GP units) and had to be transferred to a consultant unit in hospital.

I decided to review the complications of a low-risk group of patients who were considered to be potentially suitable for home confinement when they booked: 1015 consecutive patients who were delivered at the maternity unit of this hospital between August and November 1978.

Maternity Unit, Farnley District General Hospital, Liverpool L9 7AL
E A DIXON, MRCOG, registrar in obstetrics (present address: Victoria Jubilee Hospital, Kingston, Jamaica, W.I.)

Patients

With the assistance of the Community Midwifery Service a list was compiled of the categories of patients who would not be suitable for home delivery:

- nullipara;
- gravidity more than four live births;
- history of threatened abortion in the present pregnancy;
- history of infertile or recurrent miscarriages;
- history of small-for-date babies;
- last live birth more than four years earlier;
- last pregnancy ending in perinatal death, premature labour, or spontaneous abortion after 16 weeks' gestation;
- previous postpartum haemorrhage or retained placenta;
- previous pregnancy affected by rhesus sensitisation;
- medical disorders before or during the pregnancy: hypertension, cardiac disease, diabetes mellitus, etc;
- previous operations on the genital tract: caesarean section, repair of third-degree tear, myomectomy, pelvic floor repair;
- age over 35 years;
- previous difficult forceps delivery if a normal delivery had not occurred since then.

It was decided that the height of the patient would be a factor in deciding whether home delivery would be acceptable nor would her social class. The patient's home conditions would, however, be inspected before the booking for home confinement was confirmed. There was no information on the social class of the patients who were reviewed. None of the patients under review had epidural analgesia during labour, and those who were booked for 12 or 48 hour discharge were supervised in labour by the staff of the Community Midwifery Service.

Results

During the antenatal period 41 of the 278 patients (14.7%) had complications that made them unsuitable for home confinement (table 1). Before 38 weeks' gestation a patient was considered to be hypertensive if the diastolic blood pressure was above 90 mm Hg on at least two occasions more than four hours apart over 48 hours while resting in hospital. There were six patients in this category and three

continuous monitoring of fetal heart rate. Although routine monitoring of the fetal heart rate has been criticised,⁹ Dewhurst¹⁰ stated that it is better than the Fernald's stethoscope, while Donald¹¹ stressed that anomalies in the fetal heart rate may be missed without continuous monitoring. By continuously monitoring fetal heart rate the diagnosis of fetal distress would not be made only on the basis of meconium-stained liquor as is done when the patient is at home¹² or in GP units.¹³ In some cases the birth of an asphyxiated baby can be predicted and appropriate steps taken to adequately resuscitate the baby at birth. In this review 13 of 237 babies (5.5%) needed intubation for longer than two minutes. In only three cases was asphyxiation predicted, which confirms the report that a normal delivery does not preclude an infant being asphyxiated at birth.¹⁴

Of the complications in labour, 14 (42.7%) occurred in the third stage. Placental retention or postpartum haemorrhage, or both, may occur without any warning and, like asphyxiation of the baby, may carry serious consequences if not treated immediately.

Those who favour home delivery have recognised that there are dangers accompanying childbirth and they are willing to take the risks, especially when there is a good flying squad service available. James⁶ has stated that transfer to hospital late in pregnancy or in labour is undesirable from the mother's point of view for psychological reasons, while Adamson and Gare⁴ report that the delay caused by the transfer may result in a poor outcome for the infant and that even a five-minute delay may be fatal. Hudson⁵ showed in his review that the perinatal mortality rate rose for patients who were transferred in labour and in his experience the delay was commonly more than one hour.

I agree with Adamson and Gare⁴ that the first priority must be a live and healthy mother, the second a live and healthy baby, and the third a psychologically rewarding experience for the parents and baby. This review showed that there are unexpected complications in labour in even a low-risk group of patients, confirming that pregnancy, labour, and puerperium can only be judged as normal after it has taken place.

Conclusions

From a total of 1015 patients, 278 (27.4%) were considered to be suitable for home delivery at booking. Of these, 41 (14.7%) had complications in the antenatal period which made them unsuitable candidates for home confinement. Of the remaining

MRC trials in general practice

Many general practitioners will know of the Medical Research Council's trial of treatment for mild hypertension. This trial is still in progress and most of the 176 group practices that have been brought together in this research are still participating. We hope to extend this valuable framework and carry out studies of other problems suitable for investigation in general practice. Firstly, there will be studies, such as the hypertension trial, that are particularly suitable for investigation in general practice, for both scientific and organisational reasons. We shall shortly be starting a study of the use and possible adverse consequences of hormone treatment given for the relief of menopausal symptoms; a large scale trial of treatment of hypertension in elderly people; we hope, also, to start later this year, and other studies of observational or clinical trials are being discussed. The second type of study envisaged will be concerned with the conduct and organisation of medical care in general practice. Any practice in the group is, of course, free to participate only in those studies that the partners feel are of interest.

237 patients, 33 (13.9%) had complications during labour or up to the fifth postpartum day, while 16 babies (6.8%) needed specialised paediatric care at or after birth. This review shows that 31.3% of low-risk patients who were considered to be suitable for home confinement at booking had complications that could make home delivery unsafe for the patient or her newborn infant, or both.

I would like to thank Mr S J Bay, Mr J K P Perera, and Mr P G Rae for their kind permission to review these patients who were under their care, and Mr Perera and the Community Midwifery Service for their assistance.

References

- 1 Royal College of Obstetricians and Gynaecologists and the Royal College of General Practitioners. *Report on training for obstetrics and gynaecology by joint practice posts of the RCOG and RCGP*. London: Royal College of Obstetricians and Gynaecologists and Royal College of General Practitioners, 1981.
- 2 Craft I. The future of obstetrics and gynaecology in the National Health Service. *Specialist futures*. London: Oxford University Press, 1978.
- 3 Anonymous. Home or hospital confinement? *Br Med J* 1977;1:845-8.
- 4 Klaus MH, Kennell JH. *Maternal-infant bonding*. St Louis: C V Mosby, 1976.
- 5 Stewart D, Stewart L, eds. *Twenty-first century obstetrics*. New: Chapel Hill NC: NAPSAC, 1977.
- 6 Adamson GD, Gare DJ. Home or hospital birth? *JAMA* 1980;243:1732-6.
- 7 Barry CN. Home versus hospital confinement. *J R Coll Gen Pract* 1980; 30:102-7.
- 8 Hudson CK. Domiciliary obstetrics in a group practice. *Practitioner* 1968; 201:81-2.
- 9 James DK. Patients transferred in labour from general practitioner maternity units. *J R Coll Gen Pract* 1977;27:414-8.
- 10 Lewis BV, Tipton RH, Sloper IM. Changing pattern in a general practitioner obstetric unit. *Br Med J* 1978;1:884-5.
- 11 Cox CA, Fox JS, Zinkin PM, Matthews AEB. Critical appraisal of domiciliary obstetric and neonatal practice. *Br Med J* 1978;1:84-6.
- 12 Mehl LE, Peterson GH, Whitt M, et al. Outcomes of elective home births: a series of 1,146 cases. *J Reprod Med* 1977;18:281-90.
- 13 Pritchard J, MacDonald PC. *Williams' obstetrics*. New York: Appleton-Century-Crofts, 1980.
- 14 Queney W. *Home delivery and birth*. New York: Appleton-Century-Crofts, 1980.
- 15 Friedman EA. *Labour: clinical evaluation and management*. New York: Appleton-Century-Crofts, 1978.
- 16 Donald I. *Practical obstetric problems*. 5th ed. London: Lloyd-Luke, 1978.
- 17 Dewhurst PR. *Practical obstetrics and gynaecology for postgraduate*. 3rd ed. Oxford: Blackwell Scientific Publications, 1981.
- 18 Standing Maternity and Midwifery Advisory Committee. *Report on domiciliary midwifery and maternity bed needs*. London: HMSO, 1970.

(Accepted 14 April 1982)

had proteinuria. Antihypertensive treatment was started in all, and labour was induced at 38 weeks' gestation in five cases. The sixth patient had labour induced at 37 weeks' gestation because of increasing proteinuria. Six patients had a diastolic blood pressure above 90 mm Hg after the 38th week of pregnancy. Two had proteinuria. All were admitted and labour was induced after the blood pressure reading was verified.

TABLE 1—Complications in the antenatal period (n=41)

	No	%
Hypertension + proteinuria	12	29.3
Macrosomia	5	12.2
Postmaturity	5	12.2
Multiparity	3	7.4
Antenatal haemorrhage	3	7.4
Premature rupture of membranes	2	4.9
Others	9	21.9
Rhesus sensitisation		
Intrauterine growth retardation		
Intact placental death		
Chlorydramen		
Cardiovascular accident		
Chlorydramen		
Temporal epilepsy		
Appendicitis		

The patients who were considered to be postmature were at least 10 days past term. Term was considered to be 40 weeks' gestation as calculated by menstrual dates and checked by sonar scans or x-ray films. The three patients who were excluded during the antenatal period because of antenatal haemorrhage did not have placenta praevia, but they continued to bleed intermittently throughout the pregnancy and labour was induced before term.

Spontaneous rupture of membranes occurred in two patients between the 32nd and 33rd weeks of pregnancy. In one patient spontaneous labour followed shortly afterwards, while the other was delivered 48 hours after the membranes had ruptured. Both deliveries were assisted by forceps and the babies were transferred to the special care baby unit.

Rhesus antibodies rose in a rhesus-negative patient who had no antibodies present when she booked. Labour was induced before term and the baby, who was not jaundiced at birth, was transferred to the special care baby unit. The patient who had a suspected subchorionic haemorrhage during the second trimester had a forceps delivery, and the patient who developed deep vein thrombosis was treated with anticoagulant drugs before and after delivery. Temporal epilepsy was noted for the first time in one patient at 36 weeks' gestation. The cause of intrauterine fetal death occurred at 32 weeks' gestation. The cause of death was not known.

Of the 237 patients who had no complications in the antenatal period, 33 (13.9%) had complications in labour or in the first five postpartum days (table 1). Ten patients were delivered by forceps,

TABLE 2—Complications in labour and early puerperium (n=33)

	No	%
Forceps delivery	10	30.3
Retained placenta	5	15.2
Postpartum haemorrhage	5	15.2
Retained placenta	5	15.2
Concurrent infection	3	9.1
Hypertension + proteinuria	3	9.1
Postnatal labour	2	6.1
Ptychosis	1	3.0

five because fetal distress was diagnosed. The diagnosis of fetal distress was made by noting type II (late) or variable decelerations of the fetal heart rate using either the external ultrasonic Doppler or fetal scalp electrode tracings. Two of the babies, which were born alive, needed intubation at birth and two others were born with nuchal entanglement of the umbilical cord.

The other five forceps deliveries were performed because the second stage of labour was considered to be prolonged and the maternal effort was poor. In all cases the second stage lasted longer than 50 minutes and in two cases it was more than 60 minutes. There was deep transverse arrest in one case.

Three patients had a caesarean section as an emergency procedure. One had prolapse of the umbilical cord after spontaneous rupture of membranes and the others had failed to progress after 12 hours in the first stage of labour. Two patients went into spontaneous labour before 38 weeks' gestation. The babies who were born to these patients developed respiratory distress syndrome and were transferred to the special care baby unit. Nine patients had retained placentas, which were removed manually under general anaesthesia.

Four patients had postpartum haemorrhage and all needed blood transfusion. A fifth patient received a transfusion on the second postpartum day because her haemoglobin concentration was 8.4 g/dl. Three patients were hypertensive, noted for the first time in two in labour. Both had diastolic blood pressures greater than 100 mm Hg, and one had proteinuria when a catheter specimen of urine was tested. The third patient became hypertensive immediately after delivery. There was no postpartum haemorrhage in any of these patients.

Sixteen babies (6.8%) needed specialised paediatric care. Thirteen of them had to be intubated at or soon after birth. Intubation was performed if the diastolic blood pressure was above 90 mm Hg on at least two occasions more than four hours apart over 48 hours while resting in hospital. There were six patients in this category and three

Two babies who were born at 38 weeks' gestation developed the respiratory distress syndrome and were transferred to the special care baby unit. One further baby was jaundiced because of blood group ABO incompatibility.

No perinatal deaths were recorded for the patients who were still suitable for home delivery after the antenatal period.

Discussion

Of the patients who were thought to be suitable at booking for home delivery, 31.3% had complications in the antenatal period, during labour, or in the first five days of the puerperium, or their baby needed specialised paediatric care during the first few hours of life. In the antenatal period 14.7% of the patients were found to be unsuitable for home delivery. Hudson⁵ rebooked 11.2% of multiparae women before labour, and Cox et al⁸ transferred 11.1% for hospital care. Lewis et al⁸ found an increasing number of patients who were being transferred from GP units to consultant units and said that this was due to a greater awareness among general practitioners of antenatal problems.

Of the patients who were suitable for home delivery after the antenatal period, 30 (12.7%) had complications in labour. Hudson⁵ and James⁶ reported that 4.1% and 7.3% of multiparae were transferred in labour from home and GP units respectively, while Mehl et al⁸ have shown that at least 10% of women who are expected to deliver at home will require transfer to hospital for an unexpected complication in labour.

The first stage of labour was prolonged in two cases—more than 12 hours—in my review. James⁶ stated that patients were transferred to hospital after being 36 hours in the first stage. Although there is no agreement on when labour becomes prolonged in the first stage, more than 18 hours in multiparae must be considered outside normal limits. In this series assistance was not given before 50 minutes in the second stage of labour; James⁶ however, transferred multiparae to a consultant unit after 45 minutes. Like prolonged labour in the first stage, there is no agreement about when the second stage becomes prolonged. The median duration of the second stage is 20 minutes,^{15,16} and the 95th percentile value is 1.1 hours.¹⁷ Donald¹¹ stated that after 20 minutes the patient should be reviewed so that an early decision may be made about giving assistance. Since there are so many opinions on when labour becomes prolonged, the condition of the patient or the fetus, or both, must therefore be the deciding factor about giving assistance in both the first and second stages of labour.

Forceps were used in five cases of fetal distress diagnosed by

A Family Practice

Six generations in Wiltshire

DICK MAURICE

It was in 1792 that my great-great-grandfather, Thelwall Maurice, MD, settled in Marlborough. A graduate of St Thomas's Hospital, he was 25 when he joined Dr Pinckney, who had himself been in practice in the town since 1769. Thelwall's descendants in the practice are still present. My father, Dr Dick Maurice—Marlborough "Fitz," as he is known, is of course, free to participate only in those studies that the partners feel are of interest.

From the start the practice seems to have flourished. On 26 October 1795 an advertisement appeared in the *Salisbury & Winchester Journal*: "Wanted—an APPRENTICE to a SURGEON and APOTHECARY in full practice—apply to Messrs Pinckney and Maurice—Marlborough-Fitz." We claim to be the oldest group practice in the country.

Centre of a widely scattered rural community, the distances that had to be covered on horseback or by carriage were far greater than—and indeed for ensuing generations—than they are today with the motor car. Reference to the practice day-books for 1794-5 and 1799-1800, which are preserved, illustrate this. Journeys might set out by the Kennet valley—a flag or article of clothing in the window of a cottage would be a signal for the doctor to call—then to Avebury, and on through the villages to Wroughton and Swindon. Half way on this journey fresh horses would be met, brought by the direct route over the downs.

Sundays do not seem to have been a rest day; what would a doctor today think of having to make up 12 repeat prescriptions as appears under Die Solis 2 Nov 1794? But as today, Monday seems to have been a bad day; opened at random there are 26 entries under Die Lunae 8 Decemb 1794, only 14 for Die Martis 9 Decemb.

Thelwall seems to have kept himself up to date. In 1824 Longmans published the fourth edition of *A Treatise on Dislocations and on Fractures of the Joints* by "Sir Astley Cooper, Bart., FRS, Surgeon to the King etc etc." Many personal comments are included. Case XV appears on pages 54 and 55. Headed "Marlborough Feb 12-1823," it tells how George Davis, aged 35, fell and dislocated his hip. The reduction is vividly described: 30 ounces of blood were removed, the patient was put on his back, and it was David, the third son, who led St Thomas's with FRCs, who joined his father in the family practice. Thelwall died in 1830 and

David, aged 28, married the same year. He acquired a good property on the south side and toward the western end of the High Street; his father had practised from the other end of the street on the north side, and his house Llanor House. After the ancestral home Llanor Ucha in Denbighshire; this house was to remain in the family until 1955 and an extension to the house is the practice surgery to this day: initially the surgery was a small lean-to adjoining his property.

The decade starting in 1840 was an important one for the practice. The railway from London to Bath and Bristol had just been completed and passed through Swindon, which was to grow as a railway town. Marlborough's famous Castle Inn, an important staging post for the old coaches, was acquired for the foundation of a "School for the sons of Clergy etc etc." David applied for the post of medical officer for the new school. His printed application, dated 9 March 1843, with 17 testimonials from prominent medical men (two were Fellows of the Royal Society and another, David's own brother-in-law, was to become one) together with four from local persons of importance, is still preserved; so also are the original letters from which the printed application was made. He got the job and held it for three years until the increasing size of the school made a full-time medical officer essential. It was not until 1971 that the care of the school reverted to the practice on the retirement of its fourth and last full-time medical officer.

David's work load must have been prodigious. The day-book for 1845 reveals that Mrs Letitia Codrington of Wroughton House (close to Swindon—a ride there and back by the most direct route, the track over the downs, of at least 16 miles), was visited every day including Sundays from 27 September to 7 November. During this time she received 53 medicines, tablets (less usual), or lotions. She had a further 12 visits during the following three weeks. The nature of her illness is not recorded, but Mrs Codrington's coachman was also being treated, though less frequently, over the same period. During all these time journeys of equal length were being made in the opposite direction. It is satisfactory to record that Mrs Codrington lived on until 1858.

The railway and the foundation of St Mary's hospital, so close to Paddington station, made it convenient for David's two doctor sons to do their training at the new hospital—a tradition that has continued ever since. The elder son moved to Reading and his uncle and became Hon Physician to the Royal Berks. James, the younger son, armed with FRCs like his father, joined the practice in the early 1860s. James took over the family house, which was considerably extended, and incorporated a surgery while his father built a large house, Mantion Grange, just to the west of the town beyond the college. No doubt young Jimmy had to do those tiresome night calls! The surgery had a waiting room, dispensary, and a large examination and consulting room; surgery could be carried out here but was more often done in the patient's own home. More important patients would be seen in a consulting room in the house proper. The surgery backed on to a courtyard and coach houses and stables, to which there was and is access through an arch.

Marlborough, Wiltshire
DICK MAURICE, retired general practitioner

BRITISH MEDICAL JOURNAL VOLUME 284 12 JUNE 1982

Lloran House, 1905. Dr James standing beside his carriage, Oliver on the pavement with his children, Thelwall in front porch.

1757

BRITISH MEDICAL JOURNAL VOLUME 284 12 JUNE 1982

Awareness of alcohol problems

One difficulty of dealing with alcohol problems is recognising the problem. To help increase one's awareness of patients with alcohol problems it is useful to have a check-list or a register of patients at risk (table I). The physical complaints of heavy

TABLE I—Alcohol problem check-list

Accidents	Occupations
Work, home, road	Catering trade
	Pollitons
	Seamen
Alcoholic symptoms	Physical symptoms
Smelling of alcohol at consultation	Gastrointestinal upset—pain, vomiting, diarrhoea
Morning shakes	Obesity, especially in men
Memory losses	
Withdrawal fit	
Known alcoholic	
Blood tests	Psychological symptoms
Raised mean cell volume, particularly above 98	Anxiety
Reduced platelet count, particularly above 111	Attempted suicide
	Thyroid disease
	Sexual problems
Family	Social
Psychological problems in spouse	Financial offences
Psychological problems in children	Financial problems
Referring of wife or child	Work problems

drinkers are often vague: loss of appetite, tummy upsets, morning shakes, headache, nausea, larynx, and accidents. Symptoms of psychological difficulties include unhappiness, erratic moods, sexual failure, family conflicts, and confusion. Social inadequacy is shown by requests for a sickness certificate, underachievement, problems at work, shortage of money, and trouble with the law. The patient must be asked how much he drinks, when, how often, and whether his drinking has ever caused problems. This needs to be done with sympathy and without embarrassment. When the subject is broached most patients appreciate a frank discussion of the role alcohol plays in their lives.

Records are the GP's paper instruments for detecting patients with a drinking problem, and accurate records are important, especially in a group practice where the patient may consult with a different partner. Any suspicion should be recorded in the notes, and the alcohol problem entered on a chronic morbidity register.

Safe level of drinking

It is surprising how many people will adjust their drinking habits when it is pointed out to them the difficulties that drinking is causing. This is particularly so of those who have developed a medical problem that is instantly made worse by drinking and so reminds them constantly of the need to change their drinking habits. It is necessary to tell people what is a safe level of drinking. The Royal College of Psychiatrists recommended that the equivalent of four pints of beer for men and three pints for women was the uppermost acceptable daily intake. But this was designed to stop an individual developing dependence or cirrhosis of the liver, and damage may result from drinking much smaller amounts. A more acceptable level may be the equivalent of two or three pints of beer two or three times a week.

GP's advice

Individuals must be made aware of the health hazards of drinking since heavy drinkers are less aware of the health consequences of drinking than light drinkers (table II). One should also point out that alcohol is an addictive drug and that anyone could, in a given combination of circumstances, become alcohol-dependent (table III). Attention needs to be paid to the

TABLE II—Medical conditions related to alcohol

Accidents
One in three drivers killed on the roads have blood alcohol concentrations over the legal limit. Many accidents at work and in the home are alcohol related.
Brain damage
One in 10 patients in alcohol units have an organic brain syndrome. Shrinkage of the brain can be shown in over half the remaining 90%, many of whom have cognitive impairment on psychological testing.
Cancer
A high alcohol intake increases the incidence of cancer of the mouth, pharynx, and larynx fourfold, oesophagus threefold, and primary hepatocellular carcinoma twofold.
Circulatory system
Moderate consumption of alcohol may decrease the risk of coronary heart disease, but alcohol use increases the risk of raised blood pressure and stroke.
Gastrointestinal
Heartburn, gastritis, and impaired intestinal absorption are common effects of alcohol.
Liver disease
Mortality rates for cirrhosis are 10 times higher in alcohol-dependent. Women develop cirrhosis at lower consumption levels than men. Genetic influences probably determine hepatic susceptibility, as not all excessive drinkers develop cirrhosis.
Nutrition
Many moderate drinkers are obese, whereas many alcoholics show evidence of malnutrition.
Pancreatitis
About 25% of all cases of acute pancreatitis are related to alcohol use.
Peripheral neuritis
Develops in 10% of dependent individuals.
Pregnancy
Heavy alcohol intake is associated with the fetal alcohol syndrome. Moderate intake is associated with an increased risk of spontaneous abortion, congenital malformation, and small-for-date babies.
Psychological
Alcohol use leads to increased rates of depression and sexual difficulty. Suicide rates in men alcoholics are up to 80 times higher than in non-alcoholics.

TABLE III—Symptoms of alcohol dependence: typical order of occurrence

Completely unable to keep to a drink limit
Needing more drink than companions
Difficulty getting going drink
Spending more time drinking
Missing meals drinking
Blackout, memory loss
Drinking up interests because drinking interferes
Continuing to drink despite
Changing to some drinking habit on a work day as on a day off
Feeling ill when not drinking
Change to drinking some amount whatever mood
Feeling ill when not drinking
Trembling after drinking day before
Times when can't think of anything but getting a drink
Morning sickness or vomiting
Sweating profusely at night
Shaking
Morning drinking
Waking up shaking or frightened
Hallucinations

benefits of reducing drinking rather than to the harm of continuing excessive consumption. One obvious benefit is financial. Others include better judgment and performance, safer driving, and weight loss for obese patients.

Although many people succeed in cutting down their drinking without any help, others need help and the doctor should be willing to help. The advice should be offered not only when requested but also when the opportunity presents in any consultation, especially if it can be related to the presenting medical problem. Literature may reinforce the advice and inform the patient. The Health Education Council has produced two leaflets: *Good Health?* and *Facts about Alcohol*. Further suggested reading is listed at the end of this article.

Goal of treatment

The patient has acquired a drinking habit that is damaging his personality, his family, social life, and health. Habits are hard

1758

BRITISH MEDICAL JOURNAL VOLUME 284 12 JUNE 1982

1866 was an important year for the practice. On 17 January a public meeting was held in the Town Hall and Saverne Hospital was founded. D P Maurice and J B Maurice were both appointed to the staff, together with two other medical men. The hospital now comprises some 70 or more beds and has been continuously staffed by the family ever since. This too has led to the tradition, still maintained, of having at least one fully trained surgeon in the partnership.

Four of James's 10 sons followed a medical career. Although all four helped out at times, Oliver, the third son, was the first to settle into the practice, joining about 1895. Walter, the fifth son and my father, did eight years as a naval surgeon and then in 1906 joined his father and brother. Tragedy struck in 1912, not only did James die but a few months later Oliver developed pneumonia and also died.

Two years later came the first world war. Doctors with surgical skill were hard to come by in those desperate years. On one occasion Walter had to travel to Monmouth to remove an acute appendix. During this time there was another doctor practising in the town, but the opposition was friendly and after the war the two practices merged though separate surgeries were maintained until 1955. In 1934 my cousin Jim, son of Walter's eldest brother who had devoted his life to the Army medical service, joined the partnership. Jim saw service in the second world war (he was on the RAF reserve), but meantime my elder

brother had joined the partnership in 1938. I joined the family team in 1947 to succeed my father, Walter, and when the only non-Maurice member of the four-man partnership retired in 1949 he was succeeded by his son, a surgeon like his father. The family had been addressed by this Christian name (it is hoped with the prefix doctor) since David's time, now they were all so addressed.

At this time the old surgery was much as it had been for nearly a hundred years but during the last 30 years it has been steadily extended through the old coach houses, later garages, to form the modern building of today. There are now six suites of consulting rooms, each with separate examination room (even the trainee gets his own), record room, waiting rooms, dispensary, nurse's room, and staff room. A year ago my son joined my nephew and the rest of the five-man partnership on my retirement.

Some local families have survived our medical care for nearly 200 years. The son and heir of five generations of one of these has been brought into the world by five generations of my family. When my nephew joined five years ago his very first patient in his first surgery said, "Your great-grandfather brought me into the world, your grandfather looked after me in my boyhood and early manhood, your father has looked after me in my middle age and old age, I want you to see me into my grave": a wish that was duly fulfilled.

Practising Prevention

Alcohol

PETER ANDERSON

Size of the problem

During the past 20 years alcohol consumption in Britain has roughly doubled. In 1980 the British population spent £10 200m—7.5% of consumer outlay—on alcohol, drinking the equivalent of nine pints of beer a week for each individual over the age of 16. Associated with the increase in consumption there has been an increase in the many forms of individual and social damage related to alcohol abuse. The Office of Health Economics estimate that 150 000 people, or 0.4% of the total adult population, are dependent on alcohol; 700 000 (2%) have problems related to alcohol; and 3 000 000 (8%) are heavy drinkers, showing detectable biochemical abnormalities. The number of premature deaths due to alcohol in Britain is probably about 5000 to 10 000 a year.

Role of the general practitioner

GPs have special opportunities for helping patients with alcohol problems because they are ideally placed to recognise the problem early and intervene. In terms of prevention we should

concentrate on the heavy drinkers rather than the alcohol dependents. Most people who are dependent on alcohol only present for help around their mid-forties, after 10 to 20 years of heavy drinking during which time help might have prevented the onset of serious difficulties. A GP with 1800 adult patients will have 120 heavy drinkers, 36 with problems related to alcohol, and six alcohol addicts. Because heavy drinkers have more health problems than most people it is likely that they will consult their GP more often.

Effectiveness of treatment

GPs are frequently pessimistic about the outcome of efforts to help people who are alcohol-dependent and problem drinkers. Much of the available data about treatment, however, relates only to samples of severely dependent individuals. Results might be more encouraging in the earlier stages of the cycle, since heavy drinkers are more likely to be able to cut down their drinking than those who are alcohol-dependent. Edwards's well-known study showed that 50 alcohol-dependent men who were treated intensively with all the facilities of an alcohol unit did no better after one year than a control group given only firm advice. Thus a GP can do just as much to help patients with alcohol problems in a short time with simple resources. Although many people with alcohol problems help themselves, some whose problems remit without specialist intervention attribute the change in their drinking habits to advice from their GP.

1760

BRITISH MEDICAL JOURNAL VOLUME 284 12 JUNE 1982

to change and the patient may be ambivalent about changing his drinking pattern. This ambivalence may be met by asking the patient to draw up a balance sheet of the good and bad consequences of drinking than light drinkers (table II). One should also point out that alcohol is an addictive drug and that anyone could, in a given combination of circumstances, become alcohol-dependent (table III). Attention needs to be paid to the

Changing lifestyle

For many heavy drinkers, drinking has become their predominant interest, and to achieve their desired goal they may have to make major changes in their way of life. The patient will need help to look at impediments to change and alternatives to drinking. Impediments may be a job where drink is readily available, family stress with which the drinker cannot cope without alcohol, or the occurrence of withdrawal symptoms when he tries to stop drinking. The patient should look out for situations and feelings that trigger off drinking and work out new ways of coping with them. To help change their lifestyles patients may be asked to think of activities that they enjoy that do not involve drinking. Alternatives may become clearer if specific attention is paid to past events that triggered drinking—for example, the prematch drink may be avoided by meeting at the football ground.

Family

The family will need help in supporting the problem drinker. The family may feel confused, bitter, and devalued and will welcome the chance of being understood and participating in the process of recovery.

Review

Whatever the agreed goals it is essential to review the patient's progress and to offer continuing help and support and advice on managing difficulties. A diary in which the patient makes a note of any drink consumed, the time, the quantity, and the occasion is useful for self-audit. Supportive laboratory tests (mean cell volume, blood alcohol, and γ -glutamyl transpeptidase) are useful to monitor progress, and the results and their implications may be discussed with the patient.

Relapse

Most patients will drink again whatever the original goal of treatment, but this should not be regarded as a loss of all that has been achieved. It should be viewed as an opportunity for the patient to learn more about himself and the problem. Once the relapse has been openly discussed the patient can recognise strategies for preventing a further recurrence.

Drugs

Drugs have very little place in the long-term management of patients with alcohol problems. Drugs may be needed to help withdrawal symptoms for someone who is physically dependent on alcohol. Chlormethazolin in an initial dose of 5 mg four times a day, and then reducing over one week and then stopping, are suitable alternatives. Alcohol sensitising drugs, such as

disulfiram or citrated calcium carbimide, have little use, though some patients find them helpful.

Referral

Most heavy drinkers and patients with alcohol problems can be helped by their GP. There are some whose care the GP may wish to share. Alcoholics Anonymous provides a very supportive self-help group. Referral to an alcoholism treatment unit may be needed as much for support of the GP as for the patient. When a patient is referred it is important to maintain a relationship with the patient and to give a further appointment after the initial referral to discuss what took place.

Summary

The role of the GP in relation to drinking is to:

- be aware of problems related to drink;
- offer advice when requested;
- seek an opportunity to offer advice in any consultation;
- advise on how to cut down or stop drinking;
- supplement advice with appropriate literature;
- follow-up attempts to reduce drinking.

The advice offered should include:

- reference to the presenting problem when possible;
- information about the safe level of drinking;
- information about the health and personal hazards of excessive drinking;
- information about the nature and meaning of dependence;
- emphasis on the benefit of reducing drinking;
- a plan for a short-term goal;
- ways to cope with the difficulties after reducing drinking;
- warning of the dangers of relapse;
- explanation of the need for follow-up.

Further reading for patients and doctors

ABC of alcohol. BMJ November 1981; February 1982. A straightforward guide to helping patients with alcohol problems.
Alcohol—reducing the harm. London: Office of Health Economics, 1981. An excellent booklet covering the problems of alcohol in Britain.
Alcohol and alcoholism. BMJ September 1981; January 1982. An excellent series by Richard Smith covering the problems of alcohol in Britain.
Alcohol and alcoholism. London: Royal College of Psychiatrists, 1979. A comprehensive account of alcohol problems.
Alcohol and disease. British Medical Bulletin 1982; 38:1. A detailed account of many of the medical and social problems of alcohol.
Alcoholism. Max Glan. Teach Yourself Books: Care and Welfare Series, 1982. A very complete but readable account of alcohol problems and their treatment.
Alcoholism. Neil Kessel and Henry Whitton. Penguin Books, 1969. An easily read account of alcohol problems and their treatment.
Drinking sensibly. London: HMSO, 1981. A discussion document in the prevention and health series.
Women and alcohol. Camberwell Council on Alcoholism, 1980. A comprehensive coverage of the problems of alcohol as experienced by women.

ONE HUNDRED YEARS AGO In a recently published report by the Board of Trade on the overcrowding of emigrants in lodging-houses, Dr Brodie states that he has been personally informed that, when the beds in the lodging-houses are full, an itinerant musician is introduced to attract the attention of the sleepers, who are encouraged to rise and join the dance in which those who have been unable to obtain sleeping accommodation are engaged. When this rule has succeeded in getting them in bed to reach their places, the beds are then occupied by those who first commenced the dancing, so that the same bed or floor-place may be occupied by different persons during the night. It is not difficult to add, to comprehend the, the fearful state of overcrowding that must necessarily exist, and the consequent poisonous state of the air. (*British Medical Journal*, 1882.)