resulting in a well designed area of shelving placed on two opposing walls of our reception room. This cost us £60 in 1979 and is certainly much cheaper than some of the preformed systems that are available.¹

We then divided all our record envelopes according to sex and then according to year of birth. Then we started the time-consuming task of pruning our records and converting them to A4 folders. This took us about three months for a practice list of 3800 patients and required much help by us and by the receptionists. The receptionists made up the A4 folders and we then inserted the relevant records. Being new to the practice we had considerable interest in making the record system workable, and we also found it rewarding because we got to know many patients at an early stage through pruning and indexing their records according to disease.

The layout in the A4 record is best suited to our own purposes. It differs slightly from other recognised methods of A4 layout.¹ We decided at first to use as few inserts as possible in the record for simplicity. On the first spine we have a summary sheet of major past history and recent important findings or investigations, and on the reverse of this sheet immunisations are recorded. The second insert page holds laboratory reports and reports of x-ray examinations. On the second spine we mount hospital letters with the most recent uppermost on the left-hand side and facing this the clinical notes sheet on the right-hand side. We use separate cards for contraceptive and obstetric care and these can be filed in the front pocket of the A4 record. Drug sensitivities and other important information, such as treatment with monoamine oxidase inhibitors, are recorded on the inside front cover of the brown folder and also at the top of the clinical notes page.

For many practices it is very difficult to convert records so quickly and the initial effort required to start converting an already workable system of medical record envelopes would be considerable. Many practices convert a small number of those envelopes to A4 folders regularly until all records are converted, as in Ladywell Medical Centre in Edinburgh, where in a practice of 11 500 patients conversion of records was carried out over almost four years.² Since we were radically altering not only the individual records but the whole filing system, this had to be completed relatively quickly.

We have found that new staff quickly learn the colour-coding system. The patients quickly become accustomed to giving their name and year of birth at surgery attendances, and when phoning for home visits, repeat prescriptions, or for any other reason. Telephone queries can be dealt with speedily as the record is to hand in a few seconds, which helps both our patients and us; it also helps when local consultants phone with queries or requests in reducing the amount of time spent on the telephone and in giving accurate information.

Our visual age-sex register makes it easy for us to check and pull the case sheets of, for example, those 35-year-old women requiring cervical smears or 10 or 11-year-old girls needing rubella immunisations. We can also quickly check payments from the family practitioner committee—for example, for patients over 65 years and over 75 years. With a conventional age-sex register, firstly the relevant group of patients is defined, usually on a card index system, then the case notes are retrieved and the information gathered. With our system it is so easy to identify and check to see whether investigations or treatments

have been done that screening processes become very much simpler. We also keep a reserve card index age-sex register in our own home in case of fire or vandalism in the surgery.

Instead of storing our empty medical record envelopes inside the pocket of the A4 we file them alphabetically in cabinets for medical record envelopes. This takes up little space and provides a most useful back up for our main filing system. Any queries from the health board, hospital, social workers, patients' relatives, etc., when they do not know the year of birth, are dealt with by this method.

Benefits and possibilities

Having now rearranged our records and put them into a filing system that is reliable, easy to use, and has been tested over two and a half years, we find that our thoughts on computerisation have changed slightly. Our data for screening, research, and day-to-day medical practice is now so easy to obtain that the benefits of computerisation appear to be few. On the other hand, if computerisation becomes readily available we would agree with others¹⁻³ that one needs a data base like ours that is in a form that would make conversion to a computer feasible and comprehensible and gives us the ability to feed good data into the system.

Summary

We have established an A4 record system in which records are filed according to colour codes based on age and sex. This gives us an almost foolproof filing system with quick retrieval and has the advantage of a visible age-sex register, which is valuable for screening and research.

We are grateful to all those doctors in west and central Scotland who were so generous with their time and with their ideas, which we have adopted to our needs.

References

1. Marsh GN, Thornham JR. Changing to A4 folders and updating records in a "busy" general practice. *Br Med J* 1980;281:215-7.
2. Cormack JJC. Medical records II. A4 folders—a step into the present. *Br Med J* 1981;282:953-5.
3. Elliott A. Medical records V. A4 record system and all that. *Br Med J* 1981;282:1363-5.
4. Tulloch AJ. Medical records IV. Medical records in a large practice. *Br Med J* 1981;282:1203-6.
5. Maycock C. GP records—a simple and inexpensive system. *Br Med J* 1978;ii:1510-1.
6. Walker R. Medical records I. A personal view. *Br Med J* 1981;282:869-72.
7. Campbell IK. Medical records III. Improving records in a group practice. *Br Med J* 1981;282:1126-8.
8. Tomson P. Medical records VI. Middle-sized group practice. *Br Med J* 1981;282:1438-41.
9. Tait JG. The clinical record in British general practice. *Br Med J* 1977;ii:983-8.
10. Tait J. History of our records. *Br Med J* 1981;282:702-4.

(Accepted 18 May 1982)

conditions are referred by their doctors or health visitors (though many are self-referred) and attend a six-week course of classes held in a general practitioner's surgery. The course covers subjects such as the ill-effects of chronic drug use, alcohol, and tobacco, and other maladaptive coping mechanisms, but it emphasises positive ways of dealing with stress, including acquiring skills in muscular relaxation.

In a short article it is possible to refer only briefly to some areas of the preventive work that health visitors may do. So I conclude by welcoming the recent recognition of the importance of prevention by the other key member of the primary health care team, as demonstrated by the publication of the reports of the Royal College of General Practitioners' Working Party on prevention. At last there is a glimmer of hope among health visitors that their medical colleagues will begin to appreciate what they have been up to all these years!

The GP and the Specialist

Geriatric medicine

C T CURRIE

Although most elderly people live at home and enjoy good health and independence the care of elderly patients is the major task of the National Health Service, with most of this care being provided by the general practitioner, who is likely to see 10 or more patients over the age of 65 each day. Except in the few areas where geriatricians offer comprehensive age-related hospital services, the hospital care of most elderly patients is crisis of mobility, continence, or family support, and those with multiple and possibly interrelated physical, psychological, and social problems will be appropriate. Referrals are commonly non-specific, and laced with such phrases as "gone off her feet," "relatives at the end of their tether," and "neighbours demanding something must be done." Although the undergraduate curriculum has rarely equipped the doctor to cope with such presentations diagnosis remains his first duty. If the diagnosis or diagnoses are not apparent to the general practitioner the geriatrician may be able to help. Many geriatricians encourage early referral, because an old lady who takes to her bed on Monday can be in serious trouble (dehydration, confusion, immobility, constipation, pressure sores, hypostatic pneumonia) by Friday and early intervention carries obvious benefits, not least for anxious and exhausted relatives.

Like other specialties, geriatric medicine commands its own skills and resources. Geriatricians, accustomed to the confused or incontinent patient, familiar with the problems of families supporting old people, practised in the multiple and commonly atypical presentation of disease in the elderly, have perhaps one supreme advantage over other hospital specialists when the referral of an elderly patient is contemplated: they never object to the patient being old. The geriatrician has both an eye for the potentially remediable and an ability to think in terms of compensation for irretrievably lost function. By treating the treatable, managing the manageable, and improving around the rest, he can, in collaboration with the general practitioner, ensure the most satisfactory last few months or years for the patient. Many patients die fairly soon after referral and, when the time comes to settle for comfort, the approach to terminal care is generally both sympathetic and practical.

References

1. *The function of the health visitor*. London: Council for the Education and Training of Health Visitors, 1967.
2. Caplan G. *An approach to community mental health*. London: Tavistock Publications, 1961.
3. Bryant GM. Use of the Denver Developmental Screening Test by health visitors. *Health Visitor* 1980;82:2-5.
4. Bryant GM, Davies KJ, Newcombe RG. Standardisation of the Denver development screening test for Cardiff children. *Dev Med Child Neurol* 1979;21:353-64.
5. Williamson J, Lowther CP, Gray S. Use of health visitors in preventive geriatrics. *Geriatrics* 1966;11:362-9.
6. Laker K. Health visiting and the elderly. *Occasional Paper No 35*. Nursing Times 1981;77:137-40.
7. DHSS. *Assessing heart attack*. London: HMSO, 1981.
8. O'Connor P. Prevention of coronary heart disease: is there a role for the health visitor? *Health Visitor* 1981;84:28-30.
9. Goodson S. Curbins the cavenem in us. *Nursing Mirror* 1980;150:No 20:22-4.

Practising Prevention

The health visitor and prevention

SHIRLEY GOODWIN

Prevention has been the raison d'être of health visiting since the profession sprang into existence during the middle of the last century in response to the high infant morbidity and mortality rates in some northern cities. The official functions of the health visitor, as laid down by the statutory training body, include the prevention of mental, physical, and emotional ill-health, the early detection of ill-health, recognising and identifying need, health teaching, and providing care.¹ Preventive activity at each of Caplan's three levels of prevention² is clearly represented in these areas.

Mothers and children

The traditional focus for health visitors has been maternal and child health and, inevitably, preventive strategies with these groups have become more highly developed than with those groups who have come more lately to the attention of health visitors. Foremost of such strategies must surely be the routine home visiting of all families when a baby is expected or when there are children under 5 years of age. The opportunities for educating and giving advice to promote healthy living and for the early detection of illness or abnormality are obvious. But it is difficult to show the beneficial effect of such intervention since health visiting is a statutory service, parts of which may not be withdrawn in order to provide a control group against which to measure any effect. The introduction, however, of a regular home visiting programme (fortnightly from birth to 3 months, and then monthly until 6 months of age) to families with babies who are identified as having a statistically increased risk of "cot death" seems to have been associated with a reduction of six deaths per 10 000. R G Carpenter, in a paper on the evaluation of home visits for high-risk children in Sheffield, said that the data suggest that this reduction in mortality (comparable to the number of lives saved by the treatment of paediatric cancers) could be attributed to the visits by health visitors.

Although health visitors themselves are convinced of the value of routine visiting as a means of primary prevention, they sometimes have difficulty in persuading general practitioners that visiting healthy families, and thus a waste of time when there are sick patients to be followed up and supported and others with health-related social problems with whom the health visitor (in the opinion of the doctor) could more usefully spend time. There is probably less disagreement about the value of developmental assessment as a preventive measure. In the practice where I worked until recently a weekly child health clinic is held and children are seen by the health visitor and doctor at 6 weeks, 8 months, 21 years, and 4 years of age. A medical examination is conducted at each check-up, and the health visitor assesses the child's developmental progress. A

screening test for hearing is carried out at 8 months at a separate session. The Royal College of General Practitioners' child developmental record card is used to record observations and is kept in the medical notes. This arrangement is popular with parents, children, and health workers, not least for the continuity of care that such a system provides, and in my experience there are fewer failures to attend than when check-ups are available only at the health authority clinic.

The need for routine medical examination to be an invariable part of developmental assessment has been challenged recently, and a new approach to screening is being adopted in which the health visitors conduct the assessments, using a structured screening tool. One example is the Denver Developmental Screening Test.³ This test was developed in the United States but has been standardised⁴ for use with British children, and has been used by health visitors in South Glamorgan since 1974. One important aspect of the test is that it contains a clear-cut indication of when it is necessary to refer a child for more detailed medical assessment. I have been concerned in supervising student health visitors who are taught to use this test and have been particularly impressed with the thoroughness and practicality of the training programme, and how fast students gain confidence in the knowledge and skills relating to normal child development.

Others

At the other end of the life cycle are the elderly people who are becoming a greater proportion of the general practitioner's list and the health visitor's case load. Health visitors are constantly aware that they are in contact with only a minority of the individuals over retirement age to whom they could be of some service in terms of prevention. Case-finding with the help of the age-sex register is possible to do, but most health visitors have too great a case load to permit them to take on routine screening and visiting of all the elderly people that they, or their medical and nursing colleagues, would wish. This is unfortunate since the effectiveness of health visitors in this area has been long recognised,⁵ and a recent study⁶ has shown the positive effect of monthly visits by a health visitor on the health of elderly women, an effect that persists for at least six months after visits have ceased.

"The most important single cause of death in the United Kingdom is coronary heart disease," according to the Department of Health and Social Security,⁷ and health visitors are very much aware of the part they have to play in its prevention.⁸ I begin with the advice that they can give on infant feeding, and continue with the information about the harmful effects of smoking, and the benefits of exercise and relaxation, which may be incorporated into health education to individuals and groups, particularly of schoolchildren.

For some years I have been teaching groups of adults how to understand and cope with stress more effectively.⁹ Patients who are being weaned off tranquillisers, or who have hypertension, migraine, phobic symptoms, and other stress-related

an equally urgent crisis of support. To the ill and newly dependent patient the geriatrician offers a period of simultaneous medical and rehabilitation management, with the restoration of independence as its goal.

DAY HOSPITAL

A day hospital should not be confused with the day centre, which is staffed by social services departments. It extends the services of a geriatric unit very considerably. Patients who are even only marginally fit to remain at home may be brought up for assessment, rehabilitation, and nursing care without resort to the expensive and perhaps dependency-inducing facilities of an inpatient unit. Inpatients may be discharged earlier, families may be relieved, and continuing care may be avoided or at least delayed.

RESPIRE ADMISSIONS

The goodwill of relatives supporting patients who would otherwise require institutional care may be preserved by respite admissions—either planned, as for annual holidays, or responding to an unforeseen family crisis such as the illness of a supporter.

LONG-STATE CARE

Responsive and effective acute geriatric services backed up by good community care will minimise pressure on the scarce resource of the geriatric service—the long-state bed. Though a last resort in terms of both cost and quality of life, it will be necessary for the highly dependent patient for whom all other possibilities have proved inadequate. In practice the management of such beds is only a minor part of the geriatrician's duties, though for a general practitioner faced with a desperate family the availability of a long-state bed may be the sole standard by which a service is judged. Hence the plea for early referrals. Early involvement, successful therapeutic interventions where possible, and arrangements for support as outlined above not only improve the quality of life for the patient and the supporters but also permits all other possibilities besides long-state care to be considered before the point of desperation is reached.

Common problems

DRUGS

Drugs are a major cause of ill health in elderly people, who are more susceptible to side effects and more at risk from polypharmacy than younger patients. Many of the most effective interventions by a geriatrician will be achieved simply by stopping or radically simplifying medication. Repeat prescriptions should be used critically and sparingly, and any unwell old person should be assessed with the possibility of drug effects in mind.

FALLS

These are a common and non-specific presentation in elderly people. They may be accidental ("the kind of thing that could happen to anyone"), prodromal—as in an old person developing a silent bronchopneumonia or myocardial infarction—or the outcome of a variety of perhaps age-related dysfunctions, such as postural hypotension, impaired postural control, or stiff and painful joints. Searching for cardiac causes is rarely rewarding. As always, drugs should be suspected. Labyrinthine disorders

are rare, and the various labyrinthine sedatives commonly prescribed are more likely to cause drug-induced Parkinsonism than to prevent falls.

ACUTE CONFUSION

The sudden or rapid onset of confusion in an old person is likely to be a non-specific presentation of a specific and possibly treatable pathology. Infection in the urinary tract or respiratory tract is a likely cause. Drugs are a major but easily reversible cause of confusion in the elderly: sedatives, antidepressants, hypoglycaemics, and digoxin are frequent offenders. Hospitalisation increases confusion, which is therefore best diagnosed and managed at home if possible.

INCONTINENCE

Tactful but fearless questioning may be required to elicit details from a patient on this notoriously underreported symptom. Relatives may be more forthcoming, and for them incontinence may be a last straw. Easily treatable causes of urinary incontinence include urinary tract infection, faecal impaction (readily missed by omitting the rectal examination), inappropriate diuretic treatment, and senile vulvitis. Less easily treated is the incontinence associated with an unstable bladder, which is common in demented patients. Pads and pants, with incontinence pads at night, may make life more bearable for such patients. Faecal incontinence is most commonly caused by faecal impaction and overflow, which is grossly underdiagnosed.

Commonly missed diagnoses

DEMENZA

Appreciable mental impairment occurs in 10% of people over 65 and in 22% of those over 80, but only about one case in five is known to the general practitioner. Most cases of insidious mental loss are caused by either primary neuronal degeneration or cerebrovascular disease and are therefore at present untreatable, but diagnosis still matters because supportive management will defer a crisis and counselling will help relatives and other supporters. Dementia impairs or abolishes drug compliance and increases susceptibility to adverse reactions.

DEPRESSION

Depression is common and commonly missed. Pleomorphic and insidious in onset, depression is perhaps more easily diagnosed by someone who has never seen the patient before. Symptoms disproportionate to signs, sleep disturbance, withdrawal from previously enjoyed activities, and late-onset alcohol abuse should suggest the diagnosis. Suspect unreliable patients, though some will simply be unlikely. Small, one-day doses of the standard drugs, given at night, are tolerated and effective.

PARKINSON'S DISEASE

Again, the insidious onset of the disease may conceal the diagnosis from the patient's usual medical attendant. Tremor is commonly absent. Watching the patient rise from a chair is worthwhile, the agonisingly slow progress up and forward being characteristic. Many commonly prescribed drugs cause Parkinsonism and can be withdrawn, to gratifying effect. True Parkinson's disease usually responds well to L-dopa combination treatment, and in cases of doubt a trial of treatment is justified.

London W5 1SP
SHIRLEY GOODWIN, SRN, MNCERT, health visitor