Contemporary Themes

Attendance at a breast screening clinic: a problem of administration or attitudes

K FRENCH, A M D PORTER, S E ROBINSON, F M McCALLUM, J G R HOWIE, M M ROBERTS

Abstract

In a study of why a sample of women, aged 45-64 and registered with a group practice in Edinburgh, attended or did not attend the Edinburgh Breast Screening Clinic demographic, aetiological, social, and perceptual characteristics of attenders and non-attenders were compared. Similar proportions of attenders and non-attenders knew the chance of a breast lump being cancer and were aware of the benefits of early diagnosis and treatment. The study, however, suggests that non-attenders saw the screening clinic as a place of risk while the attenders saw screening in a positive light: 79% of non-attenders as compared with 36% of attenders said that they were afraid of cancer being found, and most women attended either to reassure themselves that they had not got breast cancer or to receive early treatment if they had. Furthermore, 72% of non-attenders as compared with 13% of attenders were anxious that their lives would be disrupted if cancer were found at the screening clinic. There may well be an important irreducible element to non-atten-

dance due to attitudinal factors; the ethical implications of attempting to eliminate this require careful consideration.

Introduction

Although there is evidence to suggest that the early diagnosis of breast cancer leads to an improvement in prognosis, the benefits of screening for the disease in asymptomatic women are still being evaluated.¹ Reports of the Health Insurance Plan study in New York indicate an improvement in mortality over nine years in the group offered screening.²

In Britain, after several feasibility studies,³⁻⁵ a national programme involving women aged 45-64 years is currently in progress.⁶ In Edinburgh this takes the form of a randomised trial. Since 1979, all women in this age group in practices randomly allocated for screening have been invited to attend a screening clinic. The letter of invitation, computer produced, is from the director of the clinic, but the interest of the individual general practitioner is emphasised.

Clearly any benefit achievable by screening will be reduced if women do not respond to their invitation, or if the procedure is not acceptable to them. The feasibility study in Edinburgh⁶ had shown that 80% of women in selected practices attended at first visit, whereas in New York it was 65%⁷ and in south Manchester 54%.⁸ Since 1979, however, roughly two-thirds of the women invited in Edinburgh have attended for screening. It is important in terms of both the cost effectiveness and the ethics of this project to determine the reason why women do not take up the invitation for screening. In association with the university department of general practice, which had been randomly assigned for screening, it was decided to set up the study reported here.

The aim of the study was to investigate the reasons why women did not attend for breast screening and to compare their

Department of Clinical Surgery, University of Edinburgh Medical School, Royal Infirmary, Edinburgh

K FRENCH, BA, research co-ordinator

S E ROBINSON, BA, SRN, research nurse

M M ROBERTS, BSC, MD, senior lecturer and director, Edinburgh Breast Screening Clinic

Department of General Practice, University of Edinburgh Medical School, Edinburgh

A M D PORTER, BA, MPHIL, lecturer

F M McCALLUM, RGN, SCM, practice nurse

J G R HOWIE, MD, FRCGP, professor

characteristics and their reasons for non-attendance with the characteristics and reasons for attendance of attenders in the same practice.

Method

Structured interview schedules were developed and piloted at the Edinburgh Breast Screening Clinic with attenders from other practices. It was not feasible to pilot the interviews with the non-attenders, but the wording of the two schedules was identical for demographic characteristics, knowledge and experience of breast disease, aspects of health behaviour, and home and work commitments. Information was also sought on factors important in the aetiology of the disease, reasons for attendance or non-attendance, and attitudes to and anxiety about screening. (Copies of the schedules used are available from the authors.) The schedules and the techniques of their administration were extensively discussed by the interviewers (KF, SR, FMCC) to minimise the possibility of bias.

The sample of attenders was drawn by randomly selecting a woman with an early appointment at the clinic and then approaching those women whose appointments fell at half-hourly intervals thereafter. Attenders were approached on arrival at the clinic and interviewed after screening but before the full results of their examinations were available.

There were three groups of non-attenders—women who wrote declining the invitation, women who confirmed that they would attend but did not, and those who made no response to the letter. The sample of each group drawn by random numbers was sent a letter explaining the purpose of the study and asking if the practice nurse (FMCC) could either visit them at home or see them at the practice.

Results

RESPONSE

Of the total practice population of 5000, 499 women aged 45-64 formed the basis of the study. Three months after the original appointment offered, 273 women had attended for screening (55%) and 205 had not (41%). Fifteen women (3%) had died or moved away and six had postponed their appointments.

Of the 90 attenders sampled, 75% agreed to be interviewed but seven were lost to the study as there was no interviewer waiting when they emerged from screening. Thus 61 women (22% of all attenders) were interviewed. Of the non-attenders sample, 54 women (49%) agreed to be interviewed. This sample comprised 26% of all non-attenders. The age distribution was similar for both samples and closely mirrored the age distributions for all attenders and non-attenders. All percentages are calculated on the 61 attenders and 54 non-attenders, and the statistical tests were all χ^2 (with Yates's correction factor) on the real numbers.

DEMOGRAPHIC AND AETIOLOGICAL CHARACTERISTICS

As shown in table I, the two samples were similar in age but slightly more of the non-attenders were single, widowed, divorced, or separated. There was a social class difference as 67% of attenders were manual working class as against 87% of the non-attenders ($p < 0.05$).

More women who attended for screening had a family history of breast cancer compared with the non-attenders, and slightly more were premenopausal or had undergone hysterectomy. On the other hand, although the numbers were small, it should be noted that twice as many non-attenders had had a previous breast complaint.

KNOWLEDGE, BELIEF, AND EXPERIENCE OF BREAST DISEASE

The two samples had several broadly similar beliefs: that early treatment always or usually made a difference to chances of survival (attenders 85%, non-attenders 77%), that breast lumps had a likelihood of 50/50 or greater of being benign (90%, 91% respectively), and that breast lumps were a symptom of breast cancer (91%, 81%). The samples had had similar exposure to programmes about breast cancer on television and to articles about it in newspapers or magazines. More attenders (90% as against 61%), however, knew someone who

TABLE I—Demographic and aetiological factors in attenders and non-attenders

	Attenders (n = 61)	Non-attenders (n = 54)
Mean age (years)	54.2	54.8
Marital status:		
Married	47 (77%)	36 (66%)
Other	14 (23%)	18 (33%)
Social class:		
I and II	11 (18%)	5 (9%)
III non-manual	9 (15%)	2 (4%)
III manual	22 (36%)	15 (28%)
IV and V	19 (31%)	32 (59%)
Menstrual status:		
Premenopausal	12 (20%)	6 (11%)
Previous hysterectomy	13 (21%)	8 (15%)
No of pregnancies:		
None	5 (8%)	6 (11%)
1-4	46 (75%)	37 (67%)
≥ 5	10 (16%)	11 (20%)
Age at first pregnancy:*		
≤ 20	15 (25%)	15 (28%)
≥ 21	0	1 (2%)
Family history of breast cancer:		
Mother/sister	2 (3%)	3 (5%)
Other	5 (8%)	0
Previous breast complaint:		
Hospital referral†	3 (5%)	6 (11%)
Mammogram‡	1	5
Breast operation‡	1	3

*Not known in every case.

†Excluding abscesses during lactation.

‡Included within hospital referral.

had had a breast lump, more attenders (87% as against 44%) knew someone who had had breast cancer, and fewer attenders thought that pain was a symptom of breast cancer (11% against 48%). These differences were significant at the 0.001 level.

ASPECTS OF HEALTH BEHAVIOUR

More attenders than non-attenders had had cervical smears (84% as against 65%) and more had had smears at their own request (20% as against 9%). Attenders were more likely to have regular dental check-ups (33% as against 15%) and were significantly more likely to have all or some of their own front teeth (52% as against 20%).

REASONS FOR ATTENDING

The 61 attenders gave 176 reasons for going to the screening clinic and these, together with the single most important reason given by each attender, are set out in table II. This table shows that, in order of importance, the main reasons for attendance were the desire to know or make sure that nothing was wrong (41%) followed by "passive" reasons (25%), the belief that screening would prevent disease (15%), and the importance of early treatment (10%). "Passive" reasons included statements such as:

"I didn't decide—I just got the letter."

"I didn't have any particular reason—the letter arrived and I thought why not?"

TABLE II—Reasons for attending screening clinic

Reasons	All reasons (n = 176)	Main reasons (n = 61)
Wishing to know:		
Make sure nothing wrong, to find out, put mind at rest, better to face things	24%	41%
Passive reasons:		
Sent by GP, got letter, didn't think, why not—no cost, don't know, screening is "official"	22%	25%
Prevention:		
Be on the safe side, preventive measure, a good opportunity	16%	15%
Importance of early treatment:		
Early treatment cures, gives better chances, catch it "in time," postponing treatment means poor outlook	15%	10%
Awareness and vulnerability:		
Right age group, symptoms, breast awareness, family history, possibility of cancer, aware of cancer	15%	2%
Miscellaneous:		
Only fair on family, screening better than self examination, etc	7%	8%

"When my doctor sent for me—when he wants anything done I just go."

"I am a great believer in trying to help people that are trying to help you."

When asked if a decision to attend for screening had been influenced by a specific person, 20 of the attenders (33%) agreed that they had been influenced by their general practitioner's "interest" in screening.

REASONS FOR NON-ATTENDANCE

The 54 non-attenders gave 95 reasons for not going to the screening clinic and these, together with the main reason given by each woman, are set out in table III. This table shows that the main reasons given in order of importance were practical difficulties (33%), a lack of interest or the belief that screening was irrelevant (26%), and fear (22%) or "not feeling like it" (9%). Only one woman mentioned the hazard of radiation dosage as a reason for non-attendance.

TABLE III—Reasons for not attending screening clinic

Reasons	All reasons (n = 95)	Main reasons (n = 54)
Practical reasons:		
Away on holiday, work commitments, family and health commitments, travelling difficulties, poor physical health	41 %	33 %
Not interested/irrelevance of screening:		
Had enough of check-ups and hospitals, been tested before, too old, can't be bothered, feel quite well	24 %	26 %
Fear:		
Can't face it, "feardy," distrust of medicine, negative influence of others	17 %	22 %
Didn't feel like it	6 %	9 %
Other:		
Forgot about it, not enough explanation in letter, screening dangerous	12 %	9 %

HOME AND WORK COMMITMENTS

Although practical difficulties were more often presented by non-attenders, more of the attenders were working (75% as against 57%) and more were working full time (36% as against 22%). Furthermore, for 43% of attenders their working hours and clinic hours were mutually exclusive; this was the case for only 28% of non-attenders. Comparable proportions of the two samples had dependants living with them, but more of the attenders had other calls on their time, such as elderly relatives or grandchildren to feed. Nevertheless, when asked how easy it was to get away from either home or work commitments for a couple of hours during the day, more of the non-attenders, in both cases, said that they thought it was difficult (overall 44% as against 21%). It should be noted that 41% of attenders had to change the time of their appointment in order to attend the clinic.

FEAR AND ANXIETY

Both attenders and non-attenders were asked if any factors on a check list about fear and anxiety relating to screening applied to them or not. The results of these questions are set out in table IV. The two

TABLE IV—Fear and anxiety related to screening

Reasons	Attenders (n = 61)	Non-attenders (n = 53)†
Feelings that one should not go looking for trouble*	7 (11 %)	30 (58 %)
Fear of cancer being found*	22 (36 %)	42 (79 %)
Anxiety about the trouble and bother entailed if anything found*	8 (13 %)	38 (72 %)
Modesty and dislike of being examined	8 (13 %)	7 (13 %)
Anxiety about screening examination	18 (30 %)	24 (45 %)

*Significant at 0.001 level.

†One non-response.

samples were similarly anxious about the screening examination and had an equal modesty and dislike of being examined. The non-attenders, however, were significantly more likely to feel "one shouldn't go looking for trouble," were more afraid of cancer being found, and were more anxious about what would happen if cancer were found ($p < 0.001$). The attenders were asked why they thought some women did not attend for screening. Of the reasons thought of by them, 40% were fear, followed by avoidance of the problem (17%) and ignorance, silliness, or feeling invulnerable (14%). Only 11% of all comments acknowledged practical obstacles to attendance.

Discussion

The findings presented in this paper have to be interpreted cautiously. At first sight it may appear that the sample of non-attending women approached for interview was small and the number of refusals high. The size of the sample was determined by the belief of the research team that it would be wrong to expect accurate recall of reasons for and feelings about refusal later than Christmas when the original invitation to attend the clinic had been in September. This left eight interviewing weeks for one part-time research worker, and a maximum of only about 100 interviews was thought feasible. The non-attenders who did not agree to be interviewed were not pressed further for ethical reasons. It should be recalled that they had already declined one invitation to attend for screening and the approach we made in relation to this study was, in effect, already a second approach on a matter that (as our researchers had shown) creates real anxiety. Nevertheless, the still substantial refusal rate by non-attenders may have concealed important factors.

There is also a danger that the researchers made women think of reasons for non-attendance when in fact they have had no clear or specific reason. (For many practices, an important factor for women's non-attendance is the accuracy of health board lists of people registered with the general practices. In this study only 3% of letters of invitation were returned as "not known at this address." This rate of return is considerably lower than the average rate for practices.)

More women who attended for screening said that they had more work or home commitments, or both, but more non-attenders said that they had found it difficult to get away from their commitments. Two particular responses that non-attenders raised—the feeling that "you shouldn't go looking for trouble" and the anxiety about the trouble and bother there would be if anything was found—seemed to match their perceived difficulties with their commitments.

For the non-attenders, the screening clinic seemed to be seen as a place of risk; over three-quarters of non-attenders said that they had been afraid that cancer would be found. The attenders also largely attributed non-attendance to fear.

Even though over half the attenders had been anxious about screening and one-third admitted—after the event—that they had been afraid of cancer being found, the attenders seemed, on the whole, to see screening in a positive light. Two of the main reasons for going to the clinic cited by attenders had been for the reassurance that they had not developed the disease, and that they would have gained the benefits of early treatment if they had.

Apart from the non-attenders' belief that pain was a symptom of breast cancer, the two samples were equally informed about the disease. The attenders may have felt more vulnerable or aware of breast cancer as more of them knew women with the disease. Nevertheless, more of the non-attenders had experienced breast problems in the past. Possibly their experiences "put them off" further voluntary investigation. An alternative explanation is that they had thought nothing could be wrong with their breasts because they had been previously reassured, and any further investigation would be unnecessary.

It seems unlikely, however, that there is a single motivating factor that differentiates between the samples; the attenders accorded awareness and vulnerability as of little overall importance. More importantly, attendance and non-attendance for

breast screening was consistent with the women's previous health behaviour.

Implications

It is difficult to be confident when commenting on an area as subjective as motivation, but whether due to personal, social, or cultural characteristics, the differences between attenders and non-attenders with regard to commitments, previous health behaviour, anxiety, and fear are so substantial as to suggest that they did not occur by chance.

The study reported here suggests that there are women who are unlikely to attend for breast (or any) screening. To do so would be out of character with their previous health behaviour, and they are further disinclined both by fear and by their perception of their home and work commitments. It seems from this research that the overall uptake rate reflects women's attitudes rather than their practical commitments. We do not know whether improved factual knowledge of breast disease would lead to a greater acceptance of screening, but we do recognise that attitudes and patterns of behaviour are difficult to change. The non-attendance rates may not be reducible without putting pressures on patients that would be widely regarded as unacceptable, given the present uncertainty over the long-term benefits of a screening programme.

The department of general practice is grateful for financial support from the Scottish Health Education Group.

This study was made possible by a grant from the Cancer Research Campaign to one of us (MMR).

References

- 1 Anonymous. Early diagnosis and survival in breast cancer. *Lancet* 1981; ii:785-6.
- 2 Shapiro S. Evidence on screening for breast cancer from a randomised trial. *Cancer* 1977;39:277-82.
- 3 Chamberlain J, Roger P, Price JL, Ginks S, Nathan BE, Burn I. Validity of clinical examination and mammography as screening tests for breast cancer. *Lancet* 1975;ii:1026-30.
- 4 George WD, Gleave EN, England PC, et al. Screening for breast cancer. *Br Med J* 1976;iii:858-60.
- 5 Edinburgh Breast Screening Clinic. Screening for breast cancer. *Br Med J* 1978;iii:175-8.
- 6 Department of Health and Social Security Working Group. Trial of early detection of breast cancer: description of method. *Br J Cancer* 1982; 44:616-27.
- 7 Fink R, Shapiro S, Lewison J. The reluctant participant in a breast cancer screening programme. *Public Health Rep* 1968;83:479-90.
- 8 Hobbs P, Smith A, George WD, Sellwood RA. Acceptors and rejectors of an invitation to undergo breast screening compared with those who referred themselves. *J Epidemiol Community Health* 1980;34:19-22.

(Accepted 23 June 1982)

Three brothers, aged 12, 10, and 8, all have juvenile diabetes. There is no family history of the disease. Is it common for siblings to be affected?

There is certainly a strong genetic component in the aetiology of type I insulin-dependent diabetes, although studies on identical twins have shown that for type I diabetes, in contrast to type II, they may be discordant in respect of diabetes for very many years.¹ This emphasises the importance of other agents, such as virus illnesses and diabetogenic dietary factors.² Where more than one sibling has type I diabetes they tend to be of similar HLA tissue types. The occurrence in three siblings of one family with neither parent being affected would suggest that one relevant haplotype was inherited from each parent, thus accounting for the greater susceptibility to diabetes of these children.³ In the Birmingham Diabetes Survey 4.8% of siblings of diabetics aged up to 29 also had diabetes, whereas the figure was only 0.4% in the controls.⁴—J M STOWERS, professor of diabetes and endocrinology, Aberdeen.

¹ Pyke DA, Nelson PG. Diabetes mellitus in identical twins. In: Creutzfeldt W, Köberling J, Neel JV, eds. *The genetics of diabetes mellitus*. Berlin: Springer-Verlag, 1976.

² Helgason T, Jonasson MR. Evidence for a food additive as a cause of ketosis-prone diabetes. *Lancet* 1981;ii:716-20.

³ Walker A, Cudworth AG. Type I (insulin-dependent) multiplex families; mode of genetic transmission. *Diabetes* 1980;29:1036-9.

⁴ Working Party appointed by the College of General Practitioners. The family history of diabetes. *Br Med J* 1965;ii:960-2.

Is there any risk of an amateur engaged in glass engraving developing silicosis? He uses various types of glass with a modified dentist's drill as an engraving tool. What reasonable precautions should he take?

Glass contains silicates with no free silica. There is no risk of developing silicosis, and glass dust is classified as a nuisance dust.¹ Some glasses used for engraving may contain up to 33% lead. There is thus a potential hazard, although whether it exists for the amateur is difficult to say. A study a few years ago of some professional glass cutters showed some raised blood lead concentrations.² Noise is another hazard from glass engraving, and this might be quite pronounced with the face held close to a dentist's drill. Ear muffs are sometimes used by amateur glass engravers. It is not clear from the question whether the engraving is being done wet or dry. If the former then the dust would probably be removed as a slurry. If the engraving is done dry, however, then the glass dust might be a general

irritant, particularly to the eyes. It is unlikely that the subject would want to wear safety goggles, for he would want to watch the work closely without any optical distortion. Furthermore, the hazard would be from a nuisance dust and not from sharp glass. Therefore, his ordinary spectacles or even good safety spectacles with a plain lens should be sufficient.—W R LEE, professor of occupational health, Manchester.

¹ Patty FA, ed. *Industrial hygiene and toxicology*. Vol 2. 2nd revised ed. New York: Interscience Publishers, 1967:2270.

² Finch WM. *Lighting for the cutting of crystal glass: a comparison and analysis of conditions in two factories, and an investigation into possible improvements*. Manchester: University of Manchester, 1981. (MSc thesis.)

Are depression or behavioural changes recognised side effects of long-term administration of oral contraceptives containing ethinylloestradiol and levonorgestrel?

Any combined oral contraceptive preparation may cause depression, anxiety, or loss of libido. It has been suggested^{1,2} that the cause may be altered tryptophan metabolism, and also that there is a relationship between depression and endometrial monoamine oxidase activity. These suggestions remain unconfirmed, and there may be other factors.¹ With older types of oral contraceptive the incidence of mild depression may be as high as 10-20% in the first months of use, but this quickly decreases with time. The Royal College of General Practitioners Survey in 1974 found no increased incidence of severe depression. Among patients taking newer oral contraceptives containing ethinylloestradiol and levonorgestrel behavioural effects do occur: one study³ reported "nervousness" and loss of libido in 4.5% of patients, but it is not clear whether these investigators asked specifically about depression. Circulating androgen concentrations (which may be related to libido in women) seem to be lower⁴ among women taking this type of contraceptive, but the relation between androgens and sexuality is not a simple one. The difficulty of assessing psychological side effects is illustrated⁵ by the high incidence of loss of libido and nervousness reported by women taking placebo in the guise of oral contraceptives.—JAMES OWEN DRIFE, lecturer in obstetrics and gynaecology, Bristol.

¹ Hawkins DF, Elder MG. *Human fertility control*. London: Butterworth, 1979.

² Andrews WC. Oral contraception. *Clin Obstet Gynaecol* 1979;6:3-26.

³ Briggs MH, Briggs M. Clinical and biochemical investigations of a variable-dose combined type oral contraceptive. *Curr Med Res Opin* 1977;5:213-6.

⁴ Bancroft J, Davidson DW, Warner P, Tyrer G. Androgens and sexual behaviour in women using oral contraceptives. *Clin Endocrinol* 1980;12:327-40.