receptive to new ideas, will be interested in the formation of the society and any guidance it can give them.

JUNE THOMPSON
Public Relations Officer

Health Visitors' Association,
London SW1V 1PF

SIR,—It is somewhat unfair to the health visiting profession to assume, as Dr R P Snaith is tempted to do (14 August, p 512), that the absence of specific mention of postnatal depression in Shirley Goodwin's article (17 July, p 182) indicates a lack of concern with psychiatric disorder on the part of health visitors in general. As part of a programme of research undertaken here by the General Practice Research Unit, and funded by the Department of Health and Social Security, we are currently studying health visitors in one London borough, with particular reference to their role in the identification and management of psychosocial problems in the families they visit. Our preliminary results indicate that health visitors do not concern themselves merely with the detection of abnormalities in child development and with physical health but are keenly aware of the significance of emotional problems and of psychiatric ill health in families.¹ Indeed, the support and encouragement for this research provided by the Health Visitors' Association as well as the local health visitors themselves is, we feel, further evidence that the health visiting profession shares Dr Snaith's concern with the need to improve training and practice in this area.

MONICA E BRISCOE
ANTHONY W CLARE

General Practice Research Unit,
Institute of Psychiatry,
London SE5 8AF

¹ Briscoe ME, Lindley. *Health Visitor* 1982;55:165-8.

Childhood drownings in private swimming pools: an avoidable cause of death

SIR,—We would endorse the suggestion by Dr W Barry and his colleagues that private swimming pools be required by law to be safely enclosed (21 August, p 542). This would

also increase awareness of parents and the public about the hazards of drowning presented to toddlers in many other situations.

Between April and October 1981 four children under 3 years old were admitted to the Royal Berkshire Hospital, Reading, after drowning or near drowning. One had found his way through shrubs into an incompletely fenced private pool and died. One had fallen into a shallow garden pond and survived with severe brain damage. Two survived without brain damage: one was recovered from a private pool and the other from a bath, having been left unattended briefly.

One of us (R S M-K) has constructed a raised private pool which is cheaper and safer than the conventional sunken concrete pool (fig). Its walls are four feet high and access is by a three-step ladder from which the middle rung is removed so that, even should the ladder inadvertently be left down, a toddler would find it impossible to climb in.

WILLIAM TARNOW-MORDI

John Radcliffe Hospital,
Oxford OX3 9DU

R S MONEY-KYRLE

Medical Centre,
Wallingford, Oxon OX10 9DU

ABC of Diabetes

SIR,—In his admirable series of articles on the ABC of Diabetes, Dr Peter J Watkins rightly stresses the increased need of insulin "even when appetite declines and vomiting begins" (31 July, p 360). I hope he will not mind me pointing out that this need is even more pressing when the vomiting is accompanied by diarrhoea, as in attacks of gastroenteritis, which are so liable to precipitate ketoacidosis in diabetics.

A M NUSSEY

Birmingham B38 8DB

SIR,—We read with interest the article by Dr Peter J Watkins on unstable insulin-dependent diabetes (17 July, p 192) and wonder whether the following points might be contributory.

Firstly, the definition of brittle diabetes needs to be standardised. If this is taken to be wide,

unpredictable swings in blood glucose and ketones leading to severe disruption of life then (as Dr Watkins points out) there are many possible causes, including inappropriate treatment by the physician, deliberate or accidental errors in diet and insulin dosage on the part of the patient, intercurrent illness, menstruation, etc. We would like to summarise the evidence that, in addition to the above causes, there exists a "special type" of unstable diabetes (whose existence is doubted by Dr Watkins and others), which we believe may be characterised by the patients' pathophysiological responses to subcutaneous insulin administration.

In the last few years we have investigated 18 "difficult" diabetic patients who were distinguished in that they remained uncontrolled on the ordinarily optimal treatment of continuous subcutaneous insulin infusion, which argued against treatment errors as the cause of their instability. They were aged 12-37 years (mean 19 years) and all but one were women. Seventeen had been referred to us by other hospitals. On switching to intramuscular¹ or intravenous insulin infusion, metabolic control was immediately and greatly improved, making cheating unlikely and pointing to a specific defect in subcutaneous insulin absorption. This hindrance to subcutaneous absorption is further suggested by the higher daily subcutaneous insulin dose in these patients who were unresponsive to continuous subcutaneous insulin infusion compared with that in a group of stable diabetic patients who were matched for age, sex and duration, of diabetes.²

In similar but perhaps not identical subcutaneously resistant diabetics this defect has been attributed to increased subcutaneous insulin degradation, largely on the indirect evidence that the protease inhibitor aprotinin lowers insulin requirements and improves metabolic control.³ We have not, however, been able to show accelerated in-vitro insulinolysis in subcutaneous tissue biopsy specimens taken from some of our brittle patients compared with that in stable patients and normal subjects.⁴ Abnormal vascular responses do, however, seem to be a feature of the brittle group. Using the method of photoelectric plethysmography to examine changes in blood flow close to superficial subcutaneous injection sites, we found that insulin injection causes local hyperaemia in stable diabetics and normal subjects, but not in brittle diabetics.⁵ Since blood flow is thought to be a major determinant of insulin absorption,⁶ failure to increase local blood flow in these patients could therefore impair absorption and contribute to their instability. Unabsorbed injected insulin, sequestered subcutaneously, could be subject to partial degradation or be later irregularly absorbed, perhaps causing the unpredictable hypoglycaemic episodes characteristic of some brittle diabetics. Moreover, the beneficial effect of aprotinin in some of these patients may be due to the powerful local hyperaemia at its subcutaneous injection site,⁷ rather than to inhibition of insulin breakdown.

Other mechanisms, such as metabolic hypersensitivity to emotion, stress, relative insulin deficiency, etc, may, of course, operate as well, and could modulate and even cause impairment of subcutaneous insulin absorption, by, for example, influencing injection-site blood flow. Cheating may also coexist, as patients who are frightened by their unpredictable diabetes and its medical and social consequences may be tempted to engineer readmission to the relative security of hospital.

The natural history of this type of "difficult" diabetes is obscure. The finding of only a few patients in older age groups may be due to spontaneous resolution of the brittleness, as Watkins suggests, or to an early fatal outcome of multiple emergency hospital admissions. In our series, no patient has died, although most have required treatment in the intensive care unit, but only one has remitted and the duration of brittleness has been up to 22 years. There is therefore a need for a largescale longitudinal study of the course of brittle diabetes.

The management of these patients is difficult and demanding, and the various options are too complex to discuss in detail

Raised swimming pool showing removable ladder with middle rung removed.