PRACTICE OBSERVED

Practice Research

Study of angina in patients aged 30 to 59 in general practice

RESEARCH COMMITTEE, NORTHERN REGION FACULTY, ROYAL COLLEGE OF GENERAL PRACTITIONERS*

We studied patients aged 30 to 59 in general practice with angina pectoris to try to find out how many might eventually need coronary artery by-pass surgery in the regional cardiological unit. Professor Julian of the department of cardiology in Newcastle upon Tyne had asked the research committee for an estimate to plan for the future of the unit. We also found out how patients in the community are affected by their angina. The published reports of hospital studies of angina seemed distorted, presumably because the results were derived from referred patients, who were likely to have had severe disease.

Method

General practitioners recruited by the research committee for the study were on the whole active members of the Royal College of General Practitioners. To get total figures for the prevalence of angina from known practice populations it was essential that all patients in a practice take part. In the event, 51 general practitioners from 20 practices serving a population of 125 000 people participated.

DEFINITION OF CASES

A patient was defined as suffering from angina if he or she had either collected a prescription for angina or consulted about it in the previous two or three months. An aide memoire was provided for all participating doctors that described several ways in which cases could be "recalled": repeat prescriptions for relevant drugs, continuing care, sick notes, disease indexes, and memory were used. The doctor completed a form for each patient that gave the criteria on which the diagnosis of angina had been made. Since this was a clinical diagnosis the major criterion was either a definite history or a suggestive history of anginal pain on effort. The doctor also noted previous episodes of

coronary thrombosis, changes in electrocardiographic readings, and confirmation of angina by other doctors or by the therapeutic effect of glyceryl trinitrate or other drugs prescribed for angina. The doctor noted the year that the angina had been first diagnosed and the drugs that the patient was taking.

PATIENT QUESTIONNAIRE

All patients completed a questionnaire about their angina and the effect that it had on them day to day. The doctor checked through the card afterwards. Questions were asked about the frequency of attacks of angina, the severity with regard to stress and exercise, the drugs that the patient took, and the patient's occupation and whether the angina had had any effect on that. The effect that the angina had on participation in sport, housework, shopping, or sexual intercourse was recorded. Past and current smoking habits were noted. A final question attempted to discover how eager patients would be to have surgery if it could be offered.

Results

Over four months (February-May 1979) 51 GPs from practices covering a population of 125 000 patients found 336 patients who could be included in the survey. The reporting from practice to practice was varied, with some doctors only notifying two or three patients, but there was a nucleus of 18 GPs in six practices who reported 185 (55%) patients, an average of 10 each. The number of patients reported by each practice was related to the size of the practice. Reporting fewer patients may have selected the severe cases, but an analysis comparing the age of patients, age at onset, the effect on occupation, and the factors thought to indicate severity showed no difference among them.

For analysis of the results, patients were divided into two groups: "severe" and "not severe" angina. People with "severe" angina were defined as those who got attacks not only when walking as an ordinary pace on the level, but also got attacks once or more a day and their angina affected one or more of the following activities: sport, sex, housework, and shopping—22% of the patients (21 women, 52 men).

BRITISH MEDICAL JOURNAL VOLUME 285 6 NOVEMBER 1982

TABLE II—Patients' use of glyceryl trinitrate tablets: tablet put under tongue (A) when pain occurs, (B) to prevent pain occurring

Angina	No.	%	No.	%	No.	%	No.	%
"Severe"	21	(26)	4	(19)	29	(87)	4	(16)
"Not severe"	101	(30)	1	(1)	42	(16)	181	(64)
Total	125		5		71		185	

The percentages are for each severity group.

severe angina and those whose angina was not severe, not between those who had had a myocardial infarction and those who had not. Angina had no effect on the work of 54% (115) of patients. A further 31% (103) had time off occasionally or had changed to lighter work on the same job or had to change to less heavy work; 32% (106) had stopped work altogether. Twelve patients failed to reply to this question. Fifty-five per cent (40) of patients with severe angina had to either change to less heavy work or stop work altogether, compared to 35% (93) of those with angina that was not severe. This difference was statistically significant ($p < 0.01$).

The effect of angina on participating in sport, sex, housework, and shopping for these patients is shown in table III. Questions about housework and shopping did not apply to most of the men and, similarly, those regarding sport to most women. The morbidity was high overall—even with sexual intercourse, as half of the patients had less sex or had stopped. The total morbidity together with occupation changes must have had an important effect on the whole family, though no inquiry was made.

TABLE III—Percentage of patients whose angina affected sport, sex, housework, and shopping (245 men, 51 women)

	Sport		Sex		Housework		Shopping	
	Men	Women	Men	Women	Men	Women	Men	Women
Unchanged	11	1	31	21	20	15	25	21
Stopped	32	15	18	13	3	4	7	4
Reduced	10	14	45	40	19	76	19	69
Does not apply	42	80	18	59	58	59	5	52

Discussion

Angina can be a disabling condition. The results of this survey show the effect that it has on social activities and occupations. Many patients still suffer from daily attacks despite treatment. "Severe" angina was thought to be present in 22% of the angina patients. Fry¹ thought that in 27% of his patients with angina it was moderate and in 2% it was disabling, but the criteria were not stated.

The patient may be diagnosed as having "severe" angina because of the extent of the pathology or because treatment is inadequate. Although 82% of patients with "severe" angina were taking glyceryl trinitrate, only 44% of these were taking it prophylactically. Some of the others may have been so instructed but had stopped or forgotten to do so. Ninety-two per cent were taking beta-blockers, but only 7% were taking sildofinil or perhexiline. Past treatment that was ineffective and consequently discarded was not discovered, but it seemed that some patients with considerable cardiac disability were not receiving the best drug treatment available. Similarly, smoking does not seem to have been given the attention that its danger warrants: 41% were still smoking cigarettes or a pipe, compared with 57% in a similar age group in the population at large. This underlines the need to review the function, treatment, and smoking habits of angina patients from time to time, rather than rely on repeat prescriptions and self-reporting.

There was a prevalence rate of angina of 11 per 1000 patients aged 30 to 59, using the figures from the high reporting practices. In population studies Rose² found a prevalence rate in men of

38 per 1000 patients, whereas our recorded rate in men was 16 per 1000. The method used to record practice data has obviously not given total prevalence figures, and emphasises the need to study total populations.

With regard to the question of how many persons in the northern region of England would require investigations for coronary artery disease, obviously only those patients known to the doctor to be suffering from angina can be considered for referral. In our study 2.4 per 1000 patients had "severe" angina, suggesting that 2800 patients between the ages of 30 and 60 in the northern region of England have severe angina which, despite treatment, interfered substantially with their lives. These are the patients who would have to be considered for referral for cardiological opinion. Fry¹ suggested that only 5% of angina patients require further investigation but does not say how he arrived at this figure.

In the year before the study started 14 patients were newly diagnosed as suffering from "severe" angina. If this reflects the region as a whole, there will be 350 new patients with severe angina a year. Bearing in mind that angina may improve spontaneously and that patients are keen to have further investigation, the actual number of referrals would probably be much lower. Only three (0.9%) of our patients had had cardiac surgery.

The apparent decrease in the incidence of persistent angina after the age of 55 remains unexplained (fig 2). Fry¹ found that the incidence rose until the age of 70. We studied only survivors with angina and though it is possible that the mortality of people

who develop angina may be increased, thereby excluding them from the survey, this was not considered to be the cause.

Conclusions

The severity of angina pectoris, its effect on day-to-day living, and its treatment were studied in 336 patients aged 30 to 60 in general practice. Twenty-two per cent of the patients had severe angina, which appreciably affected their lives. The results strongly suggested that drug treatment was not always used to best advantage nor was advice regarding smoking heeded. It was estimated that in the northern region 2800 patients would initially have to be considered for referral for cardiological opinion.

We thank Miss V J Male of the Statistics Department, Northern Region Health Authority, and all those general practitioners who participated in this study.

References

- Fry J. Natural history of angina in a general practice. *J R Coll Gen Pract* 1976;26:64-6.
- Rose G. Variability of angina. Some implications for epidemiology. *Br J Prev Soc Med* 1968;22:12-15.

(Accepted 18 August 1982)

Practising Prevention

Dietary advice

GODFREY FOWLER

Leave gourmandising; know, the grave doth gape for thee three wider than for other men. Shakespeare: *Henry IV*, part II, act V, scene 5.

Historically, malnutrition implied deficiencies in the diet which led to the traditional deficiency diseases, and this is still the major nutritional problem worldwide. In contrast, however, in this century in some populations diseases of overnutrition have emerged.

"Western diet"—that typical of industrial societies—is thought to contribute to a wide range of "diseases of affluence." These include ischaemic heart disease, diabetes, gall stones, diverticular disease of the colon, irritable bowel syndrome, constipation, appendicitis, large bowel cancer, haemorrhoids, varicose veins, dental caries, and, of course, obesity.

The "Western diet" is characterised by a preponderance of fatty foods, of foods with a high sugar and low fibre content, and an excess of alcohol. Moreover, 75% of food eaten in the United Kingdom is "processed" at least once, which alters the

composition and makes identification of ingredients difficult. Such diets commonly include high proportions of sweets, cakes, biscuits, butter, margarine, cream, cooking fats, and fried foods. What bread is eaten is mostly made from white flour. Total energy intake is generally excessive and there is usually also a lot of salt.

During this century the proportion of protein in the diet has remained roughly the same, the proportion of fat has increased (see figure), and that of carbohydrate has fallen—because, although there has been an increase in dietary sugar, less bread and fewer potatoes are eaten. In the past 20 years the proportion of alcohol in the diet has doubled, and this now represents about 5% of the daily energy intake. Originally the prerogative of the richer sections of society, with improved living standards such diets have now become universal in the UK.

Relation between diet and disease

There is accumulating evidence, mostly from epidemiological sources, of the association between dietary excesses and disease.

Coronary heart disease is the major killer in industrial societies. In an early study of the relation between diet and deaths from ischaemic heart disease in seven countries, Keys¹ showed that there were strong positive correlations with the amount of

Department of Community Medicine and General Practice, University of Oxford, Oxford OX2 6HE

GODFREY FOWLER, MD, FRCP, clinical reader in general practice

The distribution of patients by age and sex is shown in fig 1. The male:female ratio was about 3 to 1 in all age groups. The age at diagnosis (fig 2) reflects the incidence and falls in both men and women aged 55 and over. This has no effect on the prevalence of angina in patients who had angina for more than five years and only 10% (33) for more than 10 years.

FIG 1—The age and sex distribution of 336 patients with angina.

FIG 2—The age of 336 patients when angina was diagnosed.

Nearly all patients were taking some medicine (table 1), though 28 patients, of whom two had "severe" angina, were taking no drugs. All the drugs were used alone or in combination. As expected, glyceryl trinitrate and a beta-blocker were most commonly used together. Half of the patients who were taking long-acting trinitrates were also taking glyceryl trinitrate. No patient was on both perhexiline and nifedipine. The use of glyceryl trinitrate is shown in detail in table II. No questions were asked of the 24 patients who used no tablets prophylactically or when chest pain occurred. Presumably they either swallowed the tablets or used them on some schedule of their own.

Forty-one per cent (137) of the patients still smoked cigarettes or a pipe, and 35% (117) had smoked. This may be a slight underestimate as cigar smoking was not mentioned in our questionnaire. There was no statistical difference in smoking patterns between patients with

TABLE 1—Drug treatment of patients with angina

Drugs	No.	%	No.	%
Glyceryl trinitrate	46	(14)	185	(71)
Beta-blockers	87	(26)	185	(71)
Long-acting trinitrates	56	(17)	121	(46)
Perhexiline or nifedipine	5	(1)	22	(8)

The percentages are for each severity group.

saturated fat and dietary sugar eaten and that blood cholesterol concentrations were related to the proportion of calories intake contributed by saturated fat. Other studies showed that migrating populations, such as Japanese moving to California, in adopting a higher fat diet also acquired a higher incidence of ischaemic heart disease. More recently, the results of intervention studies such as the Oslo Heart Study,² a controlled trial in which dietary and smoking advice were studied, have shown a reduction in coronary heart deaths in men who reduced their fat intake. There is also evidence of a positive correlation between coronary heart disease and alcohol intake.

Changes in this century in the proportion of dietary carbohydrate, fat, and protein as a percentage of food energy. Source: *Practising Prevention: eating for health*.

Hypertension is associated with obesity, and losing weight leads to a fall in blood pressure. There is also a correlation between salt intake and high blood pressure, evidence that a high intake of salt may cause hypertension in susceptible individuals, and further evidence that reducing salt intake can reverse the process.

Diabetes—the incidence correlates strongly with an excessive intake of energy and fibre may be protective. High fibre diets also improve control of diabetes. Diverticular disease is associated with a deficiency in dietary fibre and an intake of dietary fibre of 30 g or more a day has been associated with a lower prevalence of diverticular disease. Although no intervention studies have been done to show the protective effect of dietary fibre, the evidence is strong and it has been effective in managing the condition.

Constipation is a common complaint in Western society but rare in countries with high fibre diets and in vegetarians. Increasing dietary fibre increases stool bulk, reduces bowel transit time, and relieves constipation. It also benefits patients with the irritable bowel syndrome.

Dental caries is a major problem in the UK. In a recent survey 80% of 5 year olds required treatment for dental caries and in 10% more than half of their teeth were seriously decayed. On the other hand, caries is rare in countries where food is unrefined.

Gall stones are associated with obesity and there is also evidence that together with appendicitis gall stones occur more often in those on a diet that is deficient in fibre.

Obesity is known to have major consequences for health. Much of the evidence comes from life insurance statistics, and there is a clear association between being overweight and premature death from several conditions—ranging from diabetes to accidents. Although obesity arises only when energy intake is greater than expenditure, the problem is not as simple as this. Weight reduction improves life expectancy, lowering blood pressure, among other things.

Cancer is in many instances attributable to aspects of diet. In

recent report Doll and Peto³ estimated that about one-third of all deaths from cancer are attributable to diet—about the same proportion as those caused by smoking. Epidemiological evidence indicates that diet may be important in determining the occurrence of cancer of the stomach, large bowel, uterus, body, gall bladder, and liver. It may also contribute to cancer of the breast and pancreas. The mechanisms are unclear, but there is some evidence that dietary fat may be implicated and that beta-carotene, the vitamin A precursor, may be protective.

Giving advice

In the light of these facts, what should the GP do? Firstly, the important role of the GP and other members of the primary health care team, especially the health visitor, in giving such advice must be acknowledged. The Latin derivation of "doctor" is "teacher," and although doctors seem reluctant to see themselves as health educators a recent survey showed that "of all the many and varied sources of health information available to the adult population, it is the GP who is most trusted and whose advice has most impact."

Secondly, advice must be concerned with the broad composition of the total diet and must at the same time be specific. Advice should also be reinforced and supplemented with written literature.

The most important objective is to maintain weight within 10% of the "ideal" or "optimal." This weight (obtained from life insurance data) is that associated with the lowest level of mortality. A useful table is that in *Which? Way to Slim* which is derived from Metropolitan Life Insurance Tables. Alternatively, body mass index (BMI) may be calculated. This is the ratio of weight (kg)/square of height (metres) and should be 20-25 for men and 19-24 for women.

Energy requirements vary greatly, but for many this means lowering energy intake by reducing the consumption of sugar and fat. Regular weighing is a necessary corollary.

Advice on individual dietary constituents should include the following: —Sugar: the average individual annual intake of sucrose is almost 100 lb! About half of it is added to beverages, cereals, etc. The amount should be halved.

—Fat: many international and national professional and governmental committees have shown a strong consensus that lowering total fat in the diet to about 30% of total energy intake can reduce the risk of coronary heart disease. The debate continues about the relative importance of saturated and polyunsaturated fats, but there is general agreement that dietary saturated fats should be reduced. A reduction in plasma cholesterol concentrations follows such changes.

—Fibre: dietary fibre, particularly cereal fibre, is of special interest. To its virtue as a laxative other benefits are now being added, and an intake of 30 g a day from foodstuffs derived from whole grains is recommended. Wholemeal bread and bran-rich breakfast cereals are obvious examples, and the increased consumption of both of these is encouraging.

—Protein: protein intake in the UK has remained remarkably stable this century, despite other major dietary changes. There is no evidence to suggest that changes are necessary, though eating less meat as a source of protein and more cereal and vegetable sources of protein incidentally reduce intake of saturated fat.

—Salt: the present average salt intake of 12 g a day is excessive and could be halved without detriment and with possible benefit in preventing hypertension.

—Alcohol: during the past 20 years alcohol consumption in the UK has doubled. To preserve health, alcohol intake should not exceed 4% of total calories, which means about 20 g or two "units" a day—the equivalent of two pints of beer or two glasses of wine or measures of spirit.

Dieting is not complicated, and an obsession with calories is unnecessary. The basic rules include: (1) a prudent diet with reduced sugar and fat and increased fibre content and less salt

and alcohol; (ii) regular weighing; (iii) avoiding eating between meals—"nibbling" should be of fresh fruit or nuts; (iv) awareness of the relation between individual energy intake, expenditure, and storage (fat); (v) motivation.

Summary

- (1) There should be a simple, standard approach to diet.
- (2) The main task is to avoid obesity. Energy intake should be adjusted to maintain optimal body weight (as defined by life insurance statistics or calculation of body mass index).
- (3) Average sugar intake should be halved. This means cutting down on sweets, chocolates, puddings, soft drinks, and sweetened beverages.
- (4) Fat intake should be reduced to about 30% (from an average now of about 40%) of energy intake and the reduction should be of "saturated" fats. This means cutting down visible fats—for example, butter, margarine, cream, fat on meat, and fried foods.
- (5) Intake of fibre should be increased to about 30 g a day (from an average now of about 20 g), and this should be from whole grain cereals.
- (6) Salt intake should be halved. Adding salt should be discouraged.
- (7) Alcohol intake should be reduced to not more than two "units" (pints of beer/glasses of wine or spirits) a day.

Law and the General Practitioner

Health and Safety at Work Act 1974: GP's duties to staff

ANNE CULLINAN, NORMAN ELLIS

For some time general practitioners could assume that legislation on health and safety at work simply need not worry them. But some general practitioners have been surprised by a visit to their premises by a health and safety inspector. The obligations laid down on the employer by the Health and Safety at Work Act etc are not onerous. Most general practitioners undoubtedly occupy premises that meet with the requirements of the legislation, and apart from a few mostly technical matters there is little in this new legislation that should cause concern. Nevertheless it may be helpful to explain the responsibilities laid on the employer, if only to forewarn general practitioners of some matters that an inspector may raise.

The 1974 Health and Safety at Work Act is not just another Factory Act. It lays down a new approach to occupational health, safety, and welfare, and it applies to many premises that are not covered by previous legislation, including laboratories, hospitals, and general practice surgeries. Although the occupiers of these

British Medical Association, London WC1H 9JP
ANNE CULLINAN, LLB, industrial relations officer, Oxford region
NORMAN ELLIS, MB, PhD, under secretary

References

- 1 Keys A, Aravanis C, Blackburn HW, et al. Epidemiological studies related to coronary heart disease: characteristics of men aged 40-59 in seven countries. *Acta Med Scand* 1966; 180, suppl. 1.
 - 2 Hietanen I, Holm L, Vahri B, et al. Effect of diet and smoking intervention on the incidence of coronary heart disease. *Lancet* 1981; ii: 1305-10.
 - 3 Doll R, Peto R. *The causes of cancer*. Oxford: Oxford University Press, 1981.
- Further reading**
- Department of Health and Social Security. *Prevention and health: everybody's business*. London: HMSO, 1976.
- Department of Health and Social Security. *Prevention and health: eating for health*. London: HMSO, 1978.
- Department of Health and Social Security. *Prevention and health: drinking for health*. London: HMSO, 1981.
- Which? Way to Slim. London: Consumers' Association, 1978.
- Anderson P. *Practising prevention: alcohol*. *Br Med J* 1982; 284: 1758-60.
- Prevention of coronary heart disease: report of expert committee. WHO Technical Report Series 678. Geneva: World Health Organisation, 1982.

Free literature

Sample literature on diets. *Fat—what made it?* and *Fibre in your food* are available from the Health Education Council, 78 New Oxford Street, London WC1A 1AH, and from local health education units.

premises already owed their visitors the "common duty of care," as laid down in the 1957 Occupier's Liability Act, this duty only required reasonable precautions to be taken for their wellbeing and was far less onerous than the duties of many employers to their employees as specified in Factory Acts, the Offices, Shops, and Railway Premises Act, and the Mines and Quarries Act.

The new Health and Safety Executive, combining the Factory Inspectorate and other central Government inspectorate, is responsible to the Health and Safety Commission. Local authority environmental health officers inspect premises and enforce the legislation in the non-industrial sector, and they follow guidance from the Commission. Because general practitioners' premises are grouped with hospitals and laboratories in the category of "health services" it is the Health and Safety Executive that inspects their premises.

The Act lays down new duties on employers (including the self-employed) to provide and maintain a safe place of work. It establishes new powers and penalties to enforce safety laws, and its aim is to make all employers and employees aware of the need for safety at the place of work. Although much of this legislation is concerned with employers who already recognise trade unions—there is a requirement to appoint union safety representatives and establish a safety committee if requested to do so—even general practitioners who have only a few employees and do not recognise trade unions have duties to fulfil.

1324

General practitioner's duties to his staff

The most important duties laid down in the Act are those of an employer to an employee. (The general practitioner has an ancillary staff who are employed by a health authority have other lesser duties.) The Act states: "It shall be the duty of every employer to ensure, so far as is reasonably practicable, the health, safety, and welfare at work of all his employees." Thus the general practitioner is required to do all that is reasonably practicable to ensure the safety of his employees. Any proceedings that may be taken under the Act are criminal. An employee can report a breach of the employer's statutory duty under the Act to the Health and Safety Executive, who may bring criminal charges. Although the Act does not in itself confer a right of civil action on the employee, it does not prevent the employee, if injured, suing the employer for being in breach of his liability as an employer or for negligence under the common law.

What is reasonably practicable?

In assessing whether it was reasonably practicable for a general practitioner to avoid a specific hazard or risk of injury a court may well look at the cost of any preventive measures, particularly if the general practitioner's resources are limited, and weigh this against the risk of personal injury and its likely severity. Thus, a larger practice may have a heavier burden imposed than one with few resources.

The Act requires that all equipment and systems of work must be safe and without risk to health. This includes waste-bins, sterilisers, photocopying machines, electric fires, electric typewriters, furniture, fire extinguishers, plugs, and any other potentially hazardous equipment. Importance is attached to the maintenance and renewal of equipment, and inspectors are particularly concerned about the arrangements for regular servicing. Attention is also paid to the age, reliability, appropriateness, and position of equipment.

Safe systems of working must be understood and used by all persons. Obvious examples include ensuring that staff do not lift heavy weights without help if there is a risk of back injury; that arrangements are made for the safe disposal of clinical waste, and that care is taken in storing flammable liquids, materials (such as paper), and drugs.

Any offences under the Medicines Acts, the Poisons Acts, or the Misuse of Drugs Act 1971, lie outside the remit of the Health and Safety at Work Act. So any unauthorised access to drugs is more likely to be of interest to the Home Office than the Health and Safety Executive.

Misuse of drugs

Since there may be some overlap of the work of these Government bodies, brief reference should be made to the Misuse of Drugs Act 1971 and its regulations. Most general practitioners can expect to encounter drug abuse in one form or another. This Act aims to restrict the availability of certain specified drugs of dependence. The following safety precautions apply to such drugs. They must be kept in a "locked receptacle" which can be opened only by the practitioner or a person authorised by him. This does not mean that your duties are absolved if you simply hide the key of the receptacle in a drawer or some other "secret place," since this could be the first place where an unauthorised person might search. Also, the courts do not accept a "locked car" as a "locked receptacle" within the meaning of the regulations, so emergency bags, if they have to be left in cars, should also be locked.

In large dispensing practices, where substantial quantities of controlled drugs may be stored, the advice of the local crime prevention officer on appropriate security precautions should be obtained. Both the police and inspectors of the Home Office

Drugs Branch are concerned with the enforcement of the Misuse of Drugs Act, and any breach is a criminal offence. Further information is available in a booklet *Doctors and the Misuse of Drugs Act, 1971* from the Home Office, 50 Queen Anne's Gate, London SW1.

Written statement of safety policy is required

The general practitioner should provide information, training, and supervision to ensure the health and safety of his ancillary staff. Specifically, unless he employs fewer than five ancillary staff, he must prepare a written statement of his general policy "with respect to the health and safety" of his employees and the organisation and arrangements for carrying out this policy. For example, any safety rules that apply to potentially hazardous operations undertaken by medical and nursing staff—the dangers of homologous serum jaundice and "High Risk Specimen" warning stickers required under the Howie Code of Practice for the Prevention of Infection in Clinical Laboratories and Post-Mortem Rooms by all Pathological Laboratories and the immediate action required by staff in respect of accidents and mishaps that might occur. He should consult his employees about its form and provide a copy of it to each employee. This small practice to give everybody a copy of this written statement may be advisable to do so. A group practice may easily overlook a change in their legal obligation when the number of employees changes. The statement could be pinned on the notice board, but if the general practitioner wished to show that it had been drawn to everyone's attention it is advisable to give everyone a copy. Once the requirement to provide a written statement is established—that is, you have five or more employees—it should cover all persons working on the premises.

The statement should be simple and avoid excessive wording that may detract from its impact. But the safety rules should be as comprehensive as necessary so that both the general practitioner and the safety inspector can ensure that the procedures are adequate. For example, all staff should be reminded to report immediately to the safety officer any incident that might cause a risk to health or safety. The written statement could be included in each employee's contract of employment. The statement might be as follows:

Health and Safety at Work

The practice's policy on health and safety at work is to provide a safe and healthy working conditions as possible and to enlist the support of its employees towards achieving these aims.

While the overall responsibility rests with your employer, all staff have a legal duty to take reasonable care to avoid injury to themselves or to others by their work activities and not interfere with or misuse any clothing or equipment provided to protect health and safety.

The main hazards that staff should be aware of are: (i) medical instruments, etc, in the context of non-injury, e.g., needles, etc; (ii) any accident to a member of the staff or a member of the public should be reported to the doctor in charge as soon as possible. You must record in writing a factual statement covering to the fullest possible extent all the circumstances of the accident at the time so that any necessary action may be taken to prevent a recurrence.

Maintaining the building

The general practitioner, "so far as is reasonably practicable as regards any place of work under (his) control," must ensure that the building is maintained in a safe condition, without risks to health. He must provide safe access to the building and exit for his staff. But if the health centre is under the direct control of a health authority the general practitioner can only do what is reasonable on his part, and any blame may ultimately lie with the authority. For example, if there is loose guttering which the health authority is under a duty to maintain according to the licence between it and the practice, so long as the general practitioner

titiner has notified the authority of the danger, preferably in writing, he should be exonerated from liability if an accident does occur. He should, however, also follow up his letter to check that action was being taken by the authority. (Liability would fall on the authority under the Defective Premises Act 1972 and the Health and Safety at Work Act, since it is both the owner of the premises and those in control of maintenance who are responsible.) Similar provisions may arise where premises are rented from a private landlord. It may be advisable for a general practitioner to examine the licence or lease under which he holds the premises to clarify whose burden it is to maintain specific parts of the premises. The staff also have a responsibility to report any risks to health and safety.

A practice may not be exonerated of all responsibility for the safety of their premises, even if they are not the owners; they may be held responsible for hazards such as highly polished and slippery floors and unsafe electrical fix. But if the doctor is responsible for maintaining the premises he must concern himself with other parts, including, for example, the condition of steps and stairways, floors, and walls.

The Act also states that, as far as is reasonably practicable, the employer must provide and maintain a working environment for his employees that is without risks to health, with adequate provision for their general welfare. The wording appears to apply to more than just the physical environment of the workplace. Here a practice would be well advised to consider whether there is an adequate rest room, refreshment facilities, and toilet and washing arrangements, etc. Again, the extent to which this can be done must depend on the practice's resources.

Seaside Practice

Bexhill: retirement haven

J H EMMANUEL

Bexhill-on-Sea is a seaside town on the south coast, nestling quietly between the more celebrated resorts of Eastbourne and Hastings. In its Victorian heyday it was a fashionable resort in its own right. Now, however, its clean air, benign climate, and flat, wide promenade make it famous for a newer activity—retirement. The imposing seafront hotels that once accommodated the rich and famous at home have now been converted to house permanently elderly residents, most of them widowed and far from any other members of their families. This is surely the capital of the "Costa Geriatrica." Like all good grafts, the oft-quoted slogan "Dover for the Continent, Bexhill for the Incontinent" contains a sharp but cruel ring of truth.

This predilection for retirement to Bexhill has led to appreciable age imbalance in the town, and therefore in the practice. Of my personal list of 2800 patients, 34% are over 65 years of age (more than twice the national average). Almost half of these are 75 years or older. Just these few statistics point up the potential problems for the general practitioner who has to organise primary medical care for his patients.

Bexhill-on-Sea, East Sussex TN39 3EE
J H EMMANUEL, MB, MRCP, general practitioner

Safety representatives and safety officers

Not all practices will have safety representatives and safety committees. These are normally appointed only when trade unions are recognised for negotiating. In premises where there is more than one employer, however—for example, health centres—they will normally be required. Employer safety representatives can inspect the workplace, investigate accidents, make representations to the general practitioner, and also insist on a joint safety committee being appointed. Safety representatives have the legal right to challenge an employer on all matters affecting health and safety standards in the workplace.

But the Health and Safety Commission have recommended that similar arrangements for safety committees and safety representatives should be made even if staff are not unionised. The Health and Safety Commission think that since both the employer and employee share a statutory duty to ensure the health, safety, and welfare at work of all an employer may find it helpful if employees contribute to the development and improvement of health and safety procedures. The Commission have therefore recommended that in non-unionised workplaces the employer should set up a safety committee drawn from both management and staff. This is hardly practicable in a small practice, so the general practitioner may prefer to nominate a member of his staff to serve as the "safety officer" and monitor standards. The practice manager could well undertake this duty. The "safety officer" should report to the general practitioner who is the "controller" of the premises.

This is the first of a three-part article on the Health and Safety at Work Act.

Some difficulties

Several features distinguish geriatric practice from practice among a predominantly younger population. Firstly, some medical conditions are almost unique to the elderly—polymyalgia rheumatica, Stokes-Adams attacks, and brain failure, to name a few. Secondly, the elderly patient frequently presents with disease or degenerative changes in several body systems. Thus, a chest infection may lead on to mental confusion, incontinence, and heart failure; the patient deteriorates from a functioning, independent citizen to a major management problem within 48 hours. The elderly patient with arthritis will soon start falling about as if her Parkinson's disease or cataracts are allowed to develop too far without timely medical intervention. Finally, social isolation is extremely prevalent in this sort of population and commonly has a devastating effect on the patient's ability to lead a fulfilling, independent existence. This factor is exacerbated by the popular dream of retiring to a bungalow by the sea. Sadly, so many couples find that having achieved this long awaited ambition they miss their life-long friends and family, and find it difficult to make a new social life. If one spouse soon becomes ill or even dies, one has the makings of a social catastrophe.

In summary, the elderly patient epitomises the need to consider what Marinker calls "the triple diagnosis," for in this

1326

age group above all, the physical, psychological, and social factors in the illness often assume equal importance, and successful management of the problem may hinge on getting all these factors right.

Surprisingly, and contrary to my initial expectations, the practice statistics show no evidence of the increased work load that a geriatric population is claimed to generate. My patients consult in surgery at the rate of 2.3 contacts per year, and home visits are requested at the rate of 0.9 a year. Both figures are close to the national average.¹ It is evident, however, that even though the quantity of the work appears average, the quality of the work is certainly extraordinary in both its scope and the demands it makes upon the caring practitioner and his team. The truth of this assertion is soon grasped by any trainee, incoming partner, or visitor to the practice—sometimes even as he enters the waiting room. Commonly a surgery of some 25 consultations does not include anyone under 65 years of age. The consultation itself requires important skills that can only be learned by practical experience and not from textbooks. The patient with arthritis may need assistance to reach the consulting room safely. When inside the hearing aid may need to be adjusted to allow communication to proceed. A shopping list of symptoms or problems, written in a tremulous hand, is often produced and has to be patiently read. Should cardiovascular or abdominal examination be required there may be layers of tightly applied corsets to be removed first, and reapplied afterwards. As mentioned before, several systems may need checking. The plan of management may require contact with other members of the team, or other agencies. Finally, treatment must be explained to the patient, and it is important to ensure that he has understood. So much for "six minutes for the patient." This type of medicine cannot be hurried, and the implications for ensuring enough time within the appointment system are obvious. Home visits are often more time-consuming because the patient is obviously more handicapped, and relatives need explanations and help.

Anticipatory care

What should the general practitioner's priorities be in caring for a predominantly aged population? Certainly most of one's time is spent diagnosing, reassuring, and treating, or mobilising specialised resources so that patients with difficult problems can receive adequate support to remain in the community. It is also important to remember the enormous potential for preventive medicine in the elderly. In this age group true anticipatory care is primarily intended not to prolong the life-span, but to prevent conditions that cause disability and dependence on others. For example, arranging attendance at a day hospital for assessment and rehabilitation of Parkinsonism or disabled patients before they go completely off their legs and require hospital admission; effective management of risk factors such as hypertension, transient ischaemic attacks, arrhythmias to prevent heart disease; careful supervision of the diabetic patient with stasis ulcers, cataracts, or vascular disease—any of which could rob him of his independence if this possibility is not anticipated.

Such is the challenge of practising in this town. What resources are required to meet the challenge effectively? They are threefold—the GP himself; his primary health care team; and the geriatric hospital services. Firstly, the practitioner himself must be an enthusiast who can establish and maintain a caring relationship with a patient in the face of chronic disability, degenerative disorders, and multiple pathology. In my experience this is not as difficult as many of my colleagues from other areas appear to believe. Most of my patients are less demanding than their younger counterparts, and usually present their complaints in a humble and apologetic manner, often with the preface, "I know it's my age doctor, but..." There is always an element of self-blame for these patients, and the doctor who shelters his indifference behind such phrases as "What can he expect at his age?" or "You

were born too soon" is therapeutically bankrupt and would probably be happier in another type of practice.

Secondly, the members of the primary health care team must be equally resilient. In the surgery the receptionists need to be particularly patient and understanding. On the district one needs not only excellent nurses, but also other specialists in the community care of the elderly. The diabetic patient with a leg ulcer may require daily attendance by a district nurse to administer insulin, monitor blood glucose concentrations, dress the ulcer, and assess the need for aids to daily living. In addition, the health visitor may be required to support the spouse, liaise with social services to obtain financial entitlements, arrange home help or meal on wheels, and, where necessary, arrange short-term admission to a nursing home to relieve the strain on the partner. Another common problem is when a patient develops dementia, possibly the cruellest of long-term geriatric problems. The spouse has to be supported heavily while the life-long partner gradually changes into a confused, aggressive, unskinted individual who cannot be left alone. In this common situation a community psychiatric nurse for the elderly can be invaluable to monitor the progression of the disorder, give appropriate treatment, arrange short-term social admissions so that the spouse can have a holiday, and eventually help the spouse through the guilt that supervenes when the patient's condition requires admission to the long-term psychogeriatric ward.

Thirdly, the overall community management of this population is made easier by enthusiastic and forward-looking consultants in geriatric medicine and allied services, such as day hospitals, short-term assessment wards, occupational therapists, and medical social workers. (Incidentally I should reveal at this point that here the department of geriatric medicine caters only for patients aged 76 and over. Younger fry remain the responsibility of the general medical firms!) In this town we are blessed with such a set-up, and life would become very difficult without them. Furthermore, the management of patients with brain failure was revolutionised by the appointment of a consultant in psychogeriatric medicine. The icing on the cake for us is the availability of GP beds in our community hospital. Patients appreciate its informality and its accessibility compared with that of the district general hospital six miles away. We general practitioners appreciate the superb quality of nursing care provided there and the stimulation of managing our own cases. Unfortunately, the predominance of degenerative joint disease stretches the resources of the orthopaedic unit to the detriment of elderly patients with painful arthritis and the younger population with more routine surgical problems.

I have perhaps belaboured the problems of the elderly because this is the main feature of medical care in this particular seaside town. We do have some holidaymakers in summer, with the usual problems of medication left at home, sunburn, and other summer ailments. These cause considerable influx here, however, consists of foreign students attending language schools and boarding with local families. This can tax one's linguistic skills, non-verbal communication abilities, and knowledge of foreign languages. Further along the coast at Rye, where there are caravan parks and holiday camps, my friend and his colleagues have to cope with a doubling of the population during the summer months—a startling increase in work load, but profitable in temporary rent fees.

Apart from the clinical challenge there are personal compensations for practising here—a pleasant environment for one's children and family, a quiet pace of life compared to that in "the big city," and lack of traffic jams and parking problems. After 10 years here I can thoroughly recommend it.

References

- 1 Cornick JF, Marinker M, Morrell DC, eds. *Teaching general practice*. Brentford, Middlesex: Kluwer Medical, 1981.
- 2 *Practical Geriatrics*. London: 1977. London: Royal College of General Practitioners, 1977.