Practice Research

Team care of elderly patients in general practice

JEFFRIE R STRANG, NOREEN CAINE, ROY M ACHESON

The development of the primary care team is one of the important changes in general practice in the past 20 years. Associated with this was extensive financial investment in health centres and privately owned group premises.¹ Nevertheless, the concept of teamwork has tended to receive only lip service, and many primary care workers are still sceptical about its usefulness. Impediments to effective communication and teamwork have arisen from problems in building design² and from each of the health professions wanting to establish its own role (Brunt G, Seminar on primary care and community medicine, Cambridge, March 1979). The division of labour that is necessary in a team is based on recognising and understanding different yet complementary roles, but this is not sufficient. We believe that the "team" must consist of a whole that is greater than the sum of its parts.³ Interaction and interdependence among members is likely to increase the effectiveness of the team when working towards a common goal.

Most research has considered the contributions of each professional group separately, although Marsh and McNay described the interaction between one doctor and his team.⁴ Our criteria of interaction were the frequency with which team members discussed the care of individual patients, and how often the care of a patient was shared or cross referred between team members. Elderly patients are the most frequent users of primary care services and often require the help of several members of the primary care team. Thus they are an ideal group to use to study teamwork in general practice.

Our aims were therefore to (i) develop a simple encounter form to record work load, which would be acceptable to all members of the primary care team for use during a busy day; (ii) quantify the interaction between members of the primary care team by studying patterns and numbers of cross referrals.

Method

Forty six doctors and their nursing colleagues in 17 practices collected data for 14 days between April 1979 and February 1980. The practices were in urban, rural, and semirural areas in East Anglia. Four doctors were single handed, five practices had two partners, and eight were training practices with groups of four or five partners.

A team consisted of doctors, nurses, health visitors, and attached social workers. By our definition the work of a "practice nurse" was treating patients in the surgery, most nurses were employed by the doctors. A "community nurse" treated patients in their homes and was employed by the area health authority. When one nurse did both

Department of Community Medicine, University of Cambridge
JEFFRIE R STRANG, MR, MPH, research associate (presently a general practitioner, Redcar Health Centre, Redcar, Cleveland)
NOREEN CAINE, BA, research associate (presently a general practitioner, Redcar Health Centre, Redcar, Cleveland)
ROY M ACHESON, DM, FRCP, professor of community medicine
Correspondence to: Professor Acheson, Department of Community Medicine, Addenbrooke's Hospital, Hills Road, Cambridge CB2 2QQ

jobs she was asked to record information about each separately. Community nurses and health visitors were members of the team in all the practices. Each of the group practices and three practices with two partners had practice nurses; two group practices had a social worker attached. Each time medical care or advice was given in person or by telephone to a patient aged 65 or over by a member of the primary care team during two consecutive weeks an encounter form was completed (fig 1). Furthermore, each team member recorded on an encounter card a tally of all encounters with patients who were under 65. Thus at the end of the two weeks, by summing the number of encounter forms and the tallies on the encounter cards, we had information about the number of patient encounters made by every member of the team. Administrative personnel tried to ensure that encounter forms were completed.

Results

The total number of patients served by the 17 practices was 113 066, of whom 16.7% were aged 65 or over. During the two weeks of the study 6744 (26.8%) of the 25 134 encounters were with elderly patients.

The form is titled 'Encounter Form'. It has a 'Staff code' field. Below it are 'Date of encounter' (with checkboxes for 65+, 75-84, 85+) and 'Age of patient' (with checkboxes for 65+, 75-84, 85+). There are 'Sex' (Male/Female) and 'Place/method of encounter' (Home/Practice/Other) fields. A 'Patient's residence' field is also present. There are checkboxes for 'Is this a new episode of care?' and 'Action' (No further action, See me again, Refer to team member, etc.). At the bottom, there are checkboxes for 'Doctor', 'Practice nurse', 'Community nurse', 'Health visitor', and 'Social worker'.

FIG 1—Encounter form for use by all members of the primary care team.

WORK LOAD

Figure 2 shows the distribution of work load among team members for all 17 practices. The doctor and community nurse were responsible for nine out of every 10 encounters with elderly patients, whereas the health visitor was concerned in only one of every 100 encounters.

FIG 2—Distribution of work load with elderly patients of members of primary care teams.

Table 1 shows how the distribution of work load varied among the group practices. The doctors' work load ranged from 33% in practice 6 to 86% in practice 5, such a difference cannot readily be explained by patterns of morbidity. Annual consultation rates, standardised for age, with elderly patients were 3.4% in practice 6 and 6.2% in practice 5. Neither of these differences was due to doctors creating their own work load by initiating visits and consultations. We did, however, find in the 17 practices that the larger a partner's list the lower his work load with elderly patients was (Spearman's rank correlation, $r = -0.422$, $p < 0.05$). The practice nurse overall had 10% of the encounters in the group practices but the nurse's contribution ranged from 1% in practice 3 to 26% in practice 8 (table 1).

TABLE 1—Percentage of encounters with elderly patients by each member of the primary care team in eight group practices

Group practices	1	2	3	4	5	6	7	8
Doctors	52.6	61.4	55.3	48.5	60.5	32.6	47.5	59.8
Practice nurses	13.1	1.5	1.1	10.3	2.2	22.2	11.9	26.0
Health visitors	1.3	2.4	1.3	2.4	1.3	2.2	2.2	0.3
Social workers	13.4	28.6	41.5	37.1	24.0	52.9	47.0	18.0

INTERACTION

To fulfil our second aim to study interactions among team members we recorded whether their encounters had been initiated by them or by other members of the team. On about 2% of occasions doctors saw patients referred by non-medical colleagues, whereas doctors referred about 30% of encounters for nurses, health visitors, and social workers. Regardless of practice size, however, the fewest referrals were made to the doctor, and when the practice nurse was present most referrals were made by her (table 1). Apart from the doctor, the health visitor is the most independent member of the team. Her contribution to the care of elderly patients in all the practices was only 1% (see fig 2), but that of her had initiated between 84% and 90% of her encounters. In contrast, community nurses carried more than one third of the work load but about one in every three of their encounters was initiated by another team member. In the group practices on average half of the work of the practice nurses and of the two social workers was initiated by other team members, largely as referrals from the doctor. In one practice this meant that 13% of the total work load with elderly patients was done by the practice nurse but initiated by the doctor (table 1).

TABLE 11—Percentage of encounters initiated by other team members according to size of practice

	Group practices		Two partners		Single handed	
	No	%	No	%	No	%
Doctors	48	1.9	15	1.9	8	2.1
Practice nurses	241	48.9	0	0.0	0	0.0
Community nurses	464	26.1	97	11.6	95	28.7
Health visitors	9	16.4	2	10.7	2	10.5
Social workers	12	54.5				

Referral—The larger the practice the more often doctors referred patients to other team members (table 11). In the group practices, of the 156 referrals by doctors to other team members, 87 (55.1%) were to the practice nurse compared with 61 (38.4%) to the community nurse. Overall there were few referrals.

TABLE 111—Percentage of encounters referred to or discussed with doctors and other team members according to size of practice

	Group practices		Two partners		Single handed	
	No	%	No	%	No	%
			<i>Referred</i>			
Doctors	156	6.2	20	2.6	2	0.5
Others	112	4.8	3	0.9	9	2.6
			<i>Discussed</i>			
Doctors	48	1.9	11	1.4	0	0.0
Others	110	4.7	14	4.2	16	4.6

Discussion of cases—For every 100 elderly patients seen, about five were discussed with a colleague in the team (table 111). The larger the practice the more often the doctors discussed a case, but even so only 2% of cases. Among nurses, 5% of cases were discussed regardless of practice size, even though group practices often had common rooms.

OTHER ASPECTS

The percentage of encounters concerning continuing care of a known illness ranged from 52.9% to 77.5% for doctors and from 73.2% to 95.1% for nurses. The size of the practice did not affect the proportion of such cases.

The mean percentage of doctor encounters that took place in the patient's usual residence was 42%, and it did not vary with practice size. This compares with 45% in the *General Household Survey of 1972*. Variability between practices was considerable, however, ranging from 35% to 59.4%. The rate was not associated with how far patients lived from the practice. Of the two group practices with the lowest home visiting rate (15%), one was in an area where only 1% of the patients on the list lived three or more miles from the surgery, whereas in the other 43% of patients lived three or more miles away (table IV).

Discussion

This study made no attempt to assess whether patients served by a primary care team whose members are interdependent and communicate freely receive better care than may be provided when health professionals work independently. It accepts

TABLE IV—Percentage of encounters in the home between doctors in eight group practices and elderly patients and distance patients lived from surgery

Group practices	Home visits (%)	Distance patients lived from surgery (miles)	% of patients
1	35.8	<3	58.9
2	44.6	3-4	68.9
3	41.5	4-5	68.9
4	40.6	5-6	69.7
5	42.5	6-7	69.3
6	35.3	7-8	69.3
7	59.4	8-9	69.3
8	43.3	9-10	69.3

Norell's proposition that inefficient management, contradictory advice, and duplication of effort, which may be avoided by effective interaction between health professionals, is not good for the patient.⁵ The doctor who delegates some of his traditional work should have more time to practise his unique skills. Nurses may benefit by widening the range of their work—for instance, doing minor clinical procedures in the surgery. The community stands to gain economically from more rational deployment of scarce professional resources.

Knowing that details of process may only be inferred from our cross sectional data, we therefore attempted to determine how extensively primary care teams worked with elderly patients in some East Anglian practices. The burden of providing primary care in Britain was originally shouldered by the general practitioner working single handed. The contribution of all other team members is a modification of the original. Thus we shall discuss the work of the other members of the team.

HEALTH VISITOR

A striking finding was the small proportion of time the health visitors spent with elderly patients. Her particular contributions are education and the maintenance of health, both of which are needed by the elderly. Unlike the community nurse, the health visitor has not worked closely with the general practitioner in the past. Traditionally she was responsible to the medical officer of health and directed all her energies to preventive care of mothers and young children who were brought to her attention by the notification of births. Since the reorganisation of the National Health Service in 1974 in some parts of Britain she has continued this work independently of the general practitioner. Elsewhere she has contributed to caring for elderly patients by using an age-specific register to identify those over 65 and visiting unknown patients to assess their needs and directing them to the appropriate service—which may be provided by a doctor, a clinical nurse, a chiropodist, or a social worker. The results of some studies have shown that as many as 20% of elderly people are out of touch with the health service.⁶ Apart from assessing medical or social needs or both, health visitors could monitor compliance and advise patients to drugs among old people, especially those who do not respond to the services of the community nurse. The potential for preventive health is unknown, but many admissions to geriatric wards are due to the misuse of drugs.

Only 1.3% of encounters with elderly patients were with health visitors; the results of an unpublished study of several practices in Cambridgeshire, conducted from our department, showed that only 6% of all home visits made by health visitors

were with elderly people, and in a recent national survey 9%¹² (despite the fact that for every 100 babies born about 80 people reach their 65th birthday and about 55 reach their 75th birthday). If, however, health visitors were to fulfil their potential contribution to the care of the elderly their numbers would need to be increased.

PRACTICE NURSE

Clearly the practice nurse affects considerably the distribution of work load within the team. In one practice of four partners the practice nurse dealt with 26% of all encounters with the elderly, and about half of her work was initiated by the doctors. Without her many of these patients would have been seen by the doctors. The results of randomised controlled trials in Ontario, Canada, have shown that doctors, patients, and nurses could not differentiate the quality and outcome of care that nurses gave from that of doctors doing similar tasks.¹³⁻¹⁵

Only three of the smaller practices had employed nurses in the surgery (in one practice nursing tasks were combined with administrative work). About half of all family doctors in Britain work single handed or in two. Thus if our data are representative there could be a substantial gain in efficiency in British surgery, perhaps community nurses, spent time in such surgeries, provided that facilities were available. It is hoped that the problems arising from terms and conditions of employment may be solved to general satisfaction.

COMMUNITY NURSE

Our data indicate that the community nurse makes a large and important contribution to the care of sick elderly patients. Cartwright and colleagues found that in addition to clinical duties the community nurse may also advise.¹⁶ Nevertheless, her training and skills are mainly clinical, and she could be encouraged to provide the extra support to the doctor in the surgery, which he evidently needs, and leave to the health visitor health promotion and advice for patients who do not require clinical care.

An alternative might be to have one nursing professional concerned with the elderly, trained to offer the combined skills of a clinical community nurse and a health visitor, to care for old people. Steps have been taken in this direction with the expanded, statutory training programmes for community nurses that were introduced in 1981.

SOCIAL WORKER

Only two of 17 practices considered social workers to be members of the primary care team, even though there were social workers in all the local authority districts in which the practices were located. This reflects the problems of communication that still exist between the health and social services.

Conclusions

If primary care is to be provided by a multidisciplinary team working interdependently then planning for the distribution of the resources of one of the health professions cannot be done in isolation of planning for the others. It is possible to define the "ideal" list size for the general practitioner without considering the impact of his nursing colleagues on his work load. Planning for the care of elderly patients must be based on agreement about which tasks members of the team can and should do, and this must be backed up by effective interaction.

This work was supported by the Department of Health and Social Security, the Cambridgeshire Area Health Authority (Teaching), and the East Anglian Regional Health Authority. We are grateful to all members of the practices that we studied.

References

- Department of Health and Social Security. *Provision for health and personal social services in England. A consultation document*. HMSO, 1976.
- Department of Health and Social Security. *Health building: health centres No 36*. London: HMSO, May 1978. (Drafting).
- Bates JG. *Sick health centres and how to make them better*. Tunbridge Wells: Pitman Medical, 1978.
- Lamberts H, Riphagen FE. Working together in a team for primary health care—a guide to dangerous company. *J R Coll Gen Pract* 1975; 25: 135.
- Marsh GN, McNay RA. Team work load in an English general practice. *Br Med J* 1974; 3: 31.
- Department of Health and Social Security. *Provision for health and personal social services in England. A consultation document*. HMSO, 1976.
- Spitzer WO, Sackett DL, Sibley JG, et al. The Burlington randomised trial of the nurse practitioner. *N Engl J Med* 1974; 290: 257-6.
- Sweeney GP, May WJ. The Burlington experience: a study of nurse practitioners. *Can Fam Physician* 1973; 19: 101-10.
- Cartwright A. *Health care in the home: life before death*. London: Routledge and Kegan Paul, 1973.
- Dunnell K, Dobbs J. *Nurses working in the community*. OPCS Survey, London: HMSO, 1982.
- Spitzer WO, Sackett DL, Sibley JG, et al. The Burlington randomised trial of the nurse practitioner. *N Engl J Med* 1974; 290: 257-6.
- Sweeney GP, May WJ. The Burlington experience: a study of nurse practitioners. *Can Fam Physician* 1973; 19: 101-10.
- Cartwright A. *Health care in the home: life before death*. London: Routledge and Kegan Paul, 1973.

(Accepted 20 December 1982)

Diary of Urban Marks: 1880-1949

After passing the second conjoint, I went to Dr Calley, the dean of the medical school and asked him to put me down as a dresser for Mr Page. This meant acting as an assistant in the surgical wards and in the out-patient department during which time I could read for the intermediate MB and sit for it in the following July. Dr Calley did his best to dissuade me from going into the "House" until I had completed the school work, and said that once I went on to the final work I would never come back to the school. I did not do so then I would never obtain the MB, and this I would rue to my dying day. With the cohesiveness of youth I would not listen to his advice.

How many times in the course of my life have I regretted not taking Calley's advice. Yet, had I remained in the school and worked there solely for the MB it might have meant the ruin of my life as you who read will discover as my narrative unfolds. Often I wonder as I look back whether everything is not ultimately for one's good in this world. Is there a Guiding Power which in its wisdom shapes our ends? Life is made up of "ifs," but would one be better or worse off if any of the "ifs" had developed? If, for instance, I had never become a medical man but had entered my father's business I would have been better off than I am at present? If again I had gone to a university would I have been in the position I am in today? Who can tell?

I went into the casualty department for six weeks and on 1 January 1902 commenced to dress for Mr Page. Just as I was completing the fourth month, I was taken ill with cystitis, which was very persistent. I had to cease work and went to Jersey. This did not cure me and I went to Skegness. There I found a cure but on returning to London and finding all my contemporaries far advanced of me I threw up the idea of taking the London degree altogether and commenced to read in earnest for my finals.

In connection with St Mary's there was a swimming club and for this I played back at polo while Lees played forward. We used to visit various baths in London and play another team perhaps belonging to some reason was always known as "Strut," got drunk at the supper which followed. This was "Strut's" habit and in this state he suffered from kleptomania. He would purloin spoons, forks, knives, vases, and in fact anything which came handy. We used to keep a strict watch on him and always returned the articles on the next day after he had been told his name and made him return them. On one occasion leaving the supper room someone noticed that "Strut" had a big vase under his coat. We all got on a bus, horse driven in those days, but we would not let "Strut" get on. We let the conductor into the secret and it was a funny sight to see "Strut" running after the bus clutching at the vase while knives, forks, and spoons jingled in his pockets. He made his way to the bus and on getting into it he found that the vase was missing. He was found, not in a hospital, as at first sight might be imagined, but in a factory, where he became famous in the Indian Medical Service as an eye specialist. Still later, having retired from this and become consulting surgeon to a large eye hospital, he fell foul of the General Medical Council for

advertising and at the time of writing has been "crossed off the Roll." It is a sad ending to a brilliant life. Apart from such incidents as this, the main part of the diary is a record of the many necessary lectures, dissections, and the various examinations. It is a record of the life of an allotted member in the various wards, spent a month on the district attending births under the supervision of a superior officer and so found ourselves in the hands of the profession and the larger and more useful sphere which was to follow this.

FACTORY LABOUR A stranger, whose eye had not become familiarised with the sight, would at once be struck with the inferiority and with the general aspect of debility of these operatives. Old age is unknown to them but by the anticipation of its attributes; for their services are valued beyond forty or fifty. Instead of the usual vigor of youth, there is a sickly sallowness, languor in the place of vigorous elasticity, and weak and flabby muscles supply that tonic of vigour produced by employment under favourable circumstances. In complexion, in size, in stature, they stand forth blighted, not so much from original defects in the germ, as from the inappropriateness of the soil in which it has been reared. The diseases which prevail among them are, for the most part, diseases of debility. Acute affections are of but rare occurrence among those individuals who have for some time, been subjected to the influence of these agents. Scrofula in all its gradations, from its mild to its worst form attacking the joints, the glands, the eyes, and the viscera, is so widely spread that the disease seems here to have found its peculiar habitat. Among those who entered on the employment at an early period, and have survived the perils of youth, there are numerous cases of deformities. Many exhibit a fungus of foot which impedes progression, others, curvature of the long bones in different directions of their length, or lateral distortion at the ankle and knee joints. It has further been remarked, by those engaged in that department of practice, that women who have passed their childhood in these places very frequently require instrumental assistance at parturition. In the domain of the physician, rheumatism, chronic affections of the liver, the innumerable forms of dyspepsia, local and general effusions, bronchial disease, and phthisis are abundantly found. In an examination instituted some years since, for the information of parliament, of the persons employed in a number of mills, the physician and surgeon reported that out of the number then actually at work not one fourth of them was free from disease. Out of 266 who were working in one mill they found that sixty-three laboured under a cough, and that seventy-five were delicate. There were fourteen cases of distortion, eight were stunted, and there were eight cases of scrofula. Besides these there were others with enlarged joints, difficulty of breathing, headache, rheumatism and pain in the legs. And these were found, not in a hospital, as at first sight might be imagined, but in a factory, where he became famous in the Indian Medical Service as an eye specialist. Still later, having retired from this and become consulting surgeon to a large eye hospital, he fell foul of the General Medical Council for