PRACTICE OBSERVED

General Practice in the Year 2000

Point of view of an overseas doctor

ABBAS VIRJI

The conventional use of the term "overseas doctor" was coined to embrace all doctors working in the United Kingdom who had obtained their basic medical qualifications in medical schools abroad. Numerically, the largest group of overseas doctors is from the Indian subcontinent—17% of all doctors in the National Health Service (30% of hospital doctors and 20% of general practitioners). Those from white anglophone countries and Arab countries account for only 6% of all NHS doctors. Graduates from Eire are traditionally considered "home grown," as are doctors born overseas who qualified in Britain. In this article the term "overseas doctor" is used in a limited sense to define those doctors with particular ethnic, cultural, and linguistic characteristics that influence their future.

The history of the overseas doctor is one of the most incongruous and anomalous quirks of the twentieth century. How did large numbers of doctors trained in one of the poorest parts of the world end up working for one of the richest? Most doctors came from ex-colonies of Britain, especially India where medical schools are run on the British model and students are taught in English. Substantial immigration to the United Kingdom took place in the 1950s and 1960s for several reasons, one of the most important being the lack of British graduates to fill vacancies in the NHS. Furthermore, British training and qualifications were highly regarded, and immigration of overseas doctors was facilitated by the lack of the usual entry restrictions and the recognition of their degrees by the General Medical Council.

Thus having been allowed freely into Britain, these doctors discovered that the streets of Vilayat (United Kingdom) were not paved with gold as they had been led to believe. Handicapped by cultural and linguistic differences, the unfortunate immigrant doctor was regarded as having lower standards of

practice, and often his basic qualifications were questioned. Many found that the only jobs available were the unpopular junior hospital posts. In 1975 the report of the Merrison committee on the NHS voiced official reservations felt towards overseas doctors by the British medical profession. This resulted in the introduction of the Temporary Registration Assessment Board (TRAB) examination, which doctors who qualified overseas were now required to pass to prove their knowledge of medicine and skill with the English language. Fuelled by this development, the Overseas Doctors Association was formed officially to express the difficulties faced by its members.

Overseas Doctors Association

The Overseas Doctors Association levelled much criticism at the NHS for failing to improve conditions for overseas doctors. The main areas of concern were the difficulties in obtaining desired jobs and the subsequent uptake of the less popular specialties such as psychiatry and geriatrics. Overseas doctors were working long hours with little or no time for training or studies, and they were caught in a "catch 22" situation—that is, the lack of training facilities prevented them from being appointed to desired jobs that required previous training and experience. This often resulted in failure to obtain the further qualifications that had attracted the overseas doctor to the United Kingdom in the first instance. Many returned home without their FRCS or MRCP, while others sought refuge in the NHS, filling the jobs that no one else wanted.

There are 20 000 overseas doctors working in the NHS. Every year 2000 enter the United Kingdom, and of these, about three quarters return home with or without further qualifications. Among the 500 who stay some achieve their desired goal of consultancy and others choose to become principals in general practice. Unfortunately, there are also many whose only option is either to provide casual medical labour or to form a body of

Liter Health Centre, Camden Square, Peckham, London SE15 2LW
ABBAS VIRJI, MD, MRCP, general practitioner

2026

BRITISH MEDICAL JOURNAL VOLUME 286 25 JUNE 1983

junior hospital doctors "waiting in the ranks" for opportunities. Whether overseas doctors pick up an ailing NHS for you to decide, but the image of overseas doctors is a function of their role in the NHS, their peculiarities of religion, culture, language, and to a certain extent their "way of life." It is easy to see the problems they face and to appreciate why they often feel victims of prejudice and exploitation.

Future

What of the future? It certainly looks bleak if you are trying to settle in a foreign land and receiving endless rejections of your "work anywhere do anything" application forms. Only an extremely optimistic forecaster would predict total and satisfactory employment for all overseas doctors by the year 2000. Some will have to come to terms with the limitations, relative to absolute, imposed by the fabric of the NHS and dictated by a shortfall in their own training and experience.

I foresee that in 20 years most doctors working full time in providing casual medical labour will be overseas doctors, their ranks being swelled by women doctors with family commitments working part time and by local graduates who are temporarily unemployed. The establishment of a national local organisation will revolutionise the way night and weekend cover is handled and should end the monopoly currently enjoyed by some private organisations. The local organisation will come under the umbrella of the NHS, and all locums and casual part timers will be employed and paid by the organisation. In effect this should bring order and respectability into what has been a somewhat shambolic part of general practice, particularly in inner cities. It will be feasible for locums to become permanently attached to health centres and groups in an area. The practices wishing to use this service will be able to interview and select suitable locums to provide regular weekend and night cover and the occasional surgery. Contracts will be drawn up and terms of service agreed. Thus locums will feel more a part of a team than before and it is hoped will assume a great deal more responsibility for their work.

Furthermore, more practices will have computerised patient records and the night doctor will be able to summon relevant information from the data base. He will thus have a resume of therapies and past medical history and plans of continuing management in chronic cases. The locum's actions and decisions will be fed into the main computer at the health centre and be instantly available for the rest of the team. This will require a computer terminal in the "locumobile," which will also be equipped with the latest electronic wizardry, such as portable x-ray camera, ultrasonicography, cardiotachometer, and a complex transport facilities for the patient. Regular meetings will take place at the health centre between locums and principals to discuss cases, policies, plans, and equipment.

OOHAAH

One can visualise yet another bureaucratic complex of the national local organisation with divisions and subdivisions. One of these could easily have a jolly acronym OOHAAH—out of hours area health authority. The locum organisation, however, will be an answer to many an overseas doctor's prayers. It will provide job prospects, pension schemes, and a "future" for thousands of doctors who would otherwise either be unemployed or suffer exploitation in the hands of private organisations. With a guaranteed minimum income, a permanent place of work, and a feeling of being as much part of a community as the rest of the health centre team, the "out of hours" doctor will be at the pinnacle of our predictions for the year 2000.

In tandem with this development the general practitioner will retain his independent status but will have a much more predictable work schedule due, firstly, to list sizes shrinking to that magic number 1700, and, secondly, to greater acceptance of

computerisation. Whether the "dragon" will be replaced by a "nanosized android" in a brain the size of the earth by the year 2001 is not sure, but storage, retrieval, follow up, diagnosis, therapeutics, referral, and, indeed, interdepartmental co-ordination will be smoother and surer. Many doctors will cultivate outside appointments to widen their horizons.

Overseas doctors who have given up long hospital careers to enter general practice have a vast and enviable store of experience in various specialties. Some use this experience by working as clinical assistants in, for instance, rheumatology or chest medicine. In future there will be a great expansion of this interface between general practice and hospital, and hospital practitioner posts will become widely available. Many of these appointments will be three year rotations with options for renewal of contract by mutual agreement and will most likely be in geriatrics, psychiatry, rheumatology, ear, nose, and throat medicine, and dermatology, but others will follow suit. One would expect the emphasis to be on training rather than service to overworked departments.

Better life for the elderly

By the year 2000 the average life expectancy will have approached 90, and consequently there will be enormous increase in the geriatric age group. Although most hospital specialists will accept a greater share of responsibility for the aged, geriatrics will be numerically the largest hospital department. The general practitioner's role in the care of the elderly patients will expand as more and more geriatric wards will contain general practitioner beds. Thus the general practitioner will care for geriatric patients with both short term and long term problems and care for the dying where appropriate.

The overseas doctor, particularly from Asia, will have an important contribution to make towards society's attitude as regards the aging population. He will be ideally placed to share his cultural attitudes about the care of the elderly. In his home his cultural attitudes about the elderly, and the reason is that his country age equals wisdom, and strong family bonds are rarely severed because relatives become old and physically infirm. Continuing social stimulation should keep the octogenarians (and even nonagenarians) active and viable citizens like their counterparts in the East. Thus the family unit will adopt an axiomatic role in the life of the citizen of the twenty first century, with care of the elderly becoming part of normal family life. As for ethnic minorities in the United Kingdom in the year 2000, the results of recent population studies have shown lower birth rates among certain groups than was forecast. This second and third generation Asians will still be a minority group, but they will have none of the language problems or cultural entrenchments of their forefathers. One can foresee communication gaps narrowing and previous prejudice dwindling into insignificance. In areas of high ethnic populations the local general practitioner will no longer need to look Urdu or Punjabi, but there will still be a need for the Hakim. His herbal potions will still be prescribed in the next millennium, which may seem inexplicable in the face of high technology offering major advances in medicine. But the reason is that although most major diseases will bite the dust, the virus will be vanquished and the rheumatoid toxin, the "symptoms" will always remain. Aches and pains, giddy spells, intractable headaches, and the down and gloom of the polysymptomatic under-achiever will always ensure a place for the Hakim, the ayurvedic herbalist, or the homeopathist. Moreover, the simplicity, efficacy, and economy of acupuncture will win this form of alternative medicine a wide acceptance among the public and the profession, particularly among the overseas doctors for whom it has a special appeal.

What of the future of the Overseas Doctors Association? Hitherto its role has been to safeguard the interests of overseas doctors by ensuring equal job opportunities and providing a platform to voice grievances and complaints. It is not too optimistic to foresee better management of the overseas doctors

by the NHS while control of influx of new doctors and integration of those who are here now will eliminate the association's raison d'être. Perhaps it will be renamed the Overseas Doctors Association or exist only to serve the needs of doctors from overseas taking part in training programmes.

British medical practice will continue to be held in high regard, and the great demand for training graduates from overseas will continue. Training programmes with a planned succession of jobs both in hospitals and in general practice will be available, tailored to the needs of the overseas graduate to ensure sufficient

training and experience. Entry to the United Kingdom will be impossible unless a place on one of these programmes is obtained beforehand. This would not only guarantee a higher calibre of candidate but also greatly increase his chances of success in the specified period of time.

Financial security, better working and living conditions, excellent training facilities, freedom from exploitation, better integration, and equal job opportunities will herald the dawn of the new century for the overseas doctors. By then the term used to describe these doctors will seem anachronistic.

Practice Research

Improving the care of asthmatic patients in general practice

MICHAEL MODELL, JENNY M HARDING, ELIZABETH J HORDER, PETER R WILLIAMS

The management of asthma still poses many problems. Despite the advances in treatment made in the past 20 years, control is often inadequate and asthma may cause much disability.¹⁻³ Patients with asthma do not always make full use of health services,⁴ and poor communication between patients and doctors may result in poor care.^{5,6} These important reasons for the failure of care are of special concern to us as general practitioners,⁷ and influenced us to design a study (a) to discover whether a team of general practitioners could agree on a plan of management for asthma; (b) to examine their adherence to the plan; and (c) to devise ways of measuring the severity of asthma and use them to assess the effect of a management plan on a group of patients with asthma during one year.

The results of the study showed that though there was an appreciable improvement among younger patients, many of those who remained most severely affected at the end of the study had been inappropriately assessed and inadequately treated. We suggest ways of overcoming this.

Methods

The study took place in a group practice of eight doctors with a list of 13 000 patients. Patients with asthma aged between 5 and 55 were identified from the practice disease register and after the records were scrutinised were included in the study if they had either more than one recorded episode of wheezing in the past 12 months or a diagnosis of asthma recorded by a general practitioner in the previous two years.

Because it was too difficult to identify a control group of asthma patients, defined according to our criteria, we assessed the study group before and after a 12 month period, so that the patients acted

as their own controls. To achieve objectivity a research worker identified and interviewed the patients and examined the records.

The general practitioners discussed the care of asthma patients and without much difficulty agreed on a plan of management and treatment. This emphasised the importance of (a) giving drugs appropriate to the severity of the asthma; (b) discussion and education in the consultation; and (c) instruction in self management by the patient.

A new record card that was printed with details of the management plan was used by the doctors as an aide-memoire and a record of action taken and by the research worker to assess how well the doctors adhered to the management plan. The plan was implemented when patients came spontaneously to consult about their asthma.

Consultations for asthma during the year included the following: taking a detailed history, including past and present drug treatment and family history; measuring height and weight; examining the chest, taking peak expiratory flow readings; discussing the natural history of the disease, provoking factors, the appropriate use of medicines between and during attacks; and guidelines about when to seek medical help; discussing the plan of treatment proposed for the patient; estimating the length of the consultation; assessing severity on a four point scale.

The patients were interviewed by the research worker before the management plan was introduced and 12 months after. On both occasions she assessed the severity of asthma using the following indices:

Peak flow—The best of three peak expiratory flow readings was compared with that predicted for a non-asthmatic person of equivalent age, sex, and height and the difference expressed as standard deviations from the expected mean.⁸

Disability—This was expressed as "days of disability" in the four months before interview. Days spent in bed or in a chair score two units; entire days lost from usual activity scored one unit; and days of reduced activity scored a half unit.

Symptoms—Shortness of breath, wheezing, tightness in the chest, coughing, and sputum production were graded according to whether they occurred less than (grade 1) or more than (grade 11) once a week during the four months before interview.

The research worker also used a prepared questionnaire to assess the patients' attitudes to asthma and their knowledge of it and its treatment. This information was not given to the doctors. Of the 150 asthma patients identified, 111 were interviewed at the start of the study (25 had already left the practice list, and 30 preferred not to take part), and 92 were interviewed again 12 months later (19 had left the area and four others wished to participate). Both interviews were conducted in winter.

James Wigg Practice, Kenilworth Health Centre, London NW5
MICHAEL MODELL, MD, MRCP, general practitioner and senior clinical lecturer in general practice, University College, London
JENNY M HARDING, BA, MSc, research assistant
ELIZABETH J HORDER, BA, MSc, general practitioner
PETER R WILLIAMS, MD, MRCP, general practitioner

Correspondence to: Dr Modell, General Practice Unit, School of Medicine, University College London, University Street, London WC1E 6BJ.

2029

BRITISH MEDICAL JOURNAL VOLUME 286 25 JUNE 1983

Patients aged under 15 and patients over 15 were analysed separately, because although the children were encouraged to speak for themselves, half were helped by a parent and this influenced their responses. The distribution by age and sex of the group is shown in table 1.

TABLE 1—Distribution by age and sex of the final study group

Age (years)	No. of males	No. of females	Total No.
Under 15	26	16	42
15 and over	19	21	40
Total No.	45	37	82

Forty seven patients had atopic conditions. During the year three children and one adult were admitted to hospital and 17 attended the outpatient department. There were no deaths during the year.

Results

ADHERENCE TO THE MANAGEMENT PLAN

From the record cards prepared for the study and from the research worker's interviews with patients the doctor's adherence to the plan of management was assessed. Seventy one per cent of the cards of patients who were interviewed were completed in the first six months and 87% by the end of the study. The doctors were asked to see those patients with incomplete record cards eight months into the study; 42 patients were then recalled. The cards showed that the doctors had made considerable efforts to follow the agreed plan, although this meant longer consultations with additional examinations and discussion of a wide range of topics. The assessment consultation took an average of 14 minutes.

TABLE 2—Age, type of drug treatment, and severity of asthma graded by doctor at recorded on management cards

Type of treatment	Patients aged under 15 (n=42)	Patients aged 15 and over (n=40)	All patients (n=82)
None	10	3	13
Bronchodilators only	20	12	32
Bronchodilators and asthma control drugs	12	14	26
Bronchodilators, asthma control drugs, and inhaled steroids	7	13	20
Inhaled steroids only	3	1	4
Change in prescription recorded during study year	11	9	20
Severity graded by doctor			
Grade 1	20	21	41
Grade 2	10	10	20
Grade 3	5	6	11
Grade 4	7	3	10
Not recorded	2	5	7
Unclassified 1			
Unclassified 4			

TABLE 3—Severity of asthma at recorded at first and final interviews by research worker

Indicators of asthma severity	Percentage of patients aged under 15 (n=42)	Percentage of patients aged 15 and over (n=40)	Total (n=82)
(1) Disability: No. of disability days			
11 "Peak flow" days	45	50	48
11 "Peak flow" days	17	25	20
(2) Peak expiratory flow readings			
11 "Standard deviation from mean predicted"	15	35	40
11 "Standard deviation from mean predicted"	21	28	30
11 "Standard deviation from mean predicted"	10	24	34
(3) Symptoms: Frequency			
11 "Less than once a week"	64	76	70
11 "Once a week or more"	36	24	40

*Two patients were unable to obtain readings.

TABLE 4—Final interview: number of patients at all ages with peak expiratory flow reading related to disability and symptom grades

Disability grades	Symptoms grades	Total
1 11 111 1111	1 11	11

Peak expiratory flow reading grade 111 10 5 6 8 8 11

GRADING OF SEVERITY AND TREATMENT MANAGEMENT

When filling out the cards the doctors considered that most of their patients had "mild" or "moderate" asthma (grades 1 & 2, see table 1). They tended to recommend the less potent treatment regimens; 13 patients had no drugs, and 56 had bronchodilators or bronchodilators with cromoglycate only. Fewer children than adults were on treatment and fewer received steroids, although the severity grades in the paediatric and adult groups were closely similar.

Changes in the severity of asthma during the year as measured by the research worker (table 11) show a general tendency towards improvement in all three measures. The most significant improvement was in peak flow readings, which improved from 62% of whom improved peak flow by changing from category 111 to 11 or from 11 to 1 compared with only 28% of adults. Forty one per cent of children and 24% of adults reported fewer disability days at the end of the study.

At both interviews disability was noted between the subjective indicators—disability and symptoms—and peak flow readings (table 11). The subjective indicators gave a retrospective estimate of the severity of asthma, whereas the peak flow reading gives an accurate measurement at the time of testing. Peak flow readings are especially useful if repeated often over a long period.

ATTITUDES OF PATIENTS

Table V shows that at the end of the study nearly half of the patients still disassociated with their own understanding of asthma and also with the explanations given by their doctor. In table VI patients' views and behaviour are related to their disability, and this shows a clear relation between severity of asthma and lack of knowledge, understanding, and control. Disability may be aggravated by these factors but may also be rationalised in terms of them.

Adults were more likely than children to say that they disliked taking medicine and less likely to comply with the doctor's advice—perhaps because they do not have to make the decision. At the final interview 52 out of the 92 patients reported seeing their doctor about asthma from sources other than their doctor, mainly from books, magazines, other people with asthma, and television programmes. Seven children and 18 adults had had non-medical treatment; in addition to or instead of their prescribed medicines.

Twenty three patients remarked that during the study the doctor spent more time with them and showed more interest in their asthma, and they felt in less of a rush. Twelve remarked that they had been asked to visit their doctor specifically for a review of their asthma. Twelve patients did not attend for asthma during the study. Nine of these were assessed by the research worker as mildly affected according to all indicators. These patients seemed to be more co-ordinating