CLINICAL RESEARCH

Improved tumour localisation using indium-111 labelled antibodies

D S FAIRWEATHER, A R BRADWELL, P W DYKES, A T VAUGHAN, S F WATSON-JAMES, S CHANDLER

Abstract

Immunoglobulin G (IgG) antibodies to carcinoembryonic antigen (CEA) were labelled with radioactive indium (^{111}In) or iodine (^{131}I) and a comparison made of their value in locating CEA producing tumours. Eleven patients given ^{111}In -anti-CEA had 31 tumours as judged by a combination of all techniques. Of these, 28 were detected by ^{111}In -anti-CEA and 26 by conventional clinical techniques. Five of the patients also received ^{131}I -anti-CEA. These patients had 15 tumour areas. Thirteen were detected by ^{111}In and eight by ^{131}I . ^{111}In also produced a better signal to noise ratio in the scans and thereby showed lesions with greater certainty. In addition, the ^{111}In isotope continued to accumulate in the tumour areas for considerably longer than ^{131}I .

Absorbed doses (whole body) were similar for both isotopes. The results show that antibody scanning is greatly improved by using ^{111}In as the radiolabel in place of ^{131}I and should allow the detection of smaller or deeper lesions.

Introduction

There is considerable interest in the localisation of tumours using radiolabelled antibodies. Tumours of the gastrointestinal

tract,¹ thyroid,² germ cell tissue,^{3,4} and insulinomas⁵ have been detected using various different antibodies. In most reports iodine-131 (^{131}I) has been used as the radiolabel, but it is not ideal because the high energy gamma ray is poorly detected and the emission of β^- contributes to a high radiation dose. ^{123}I has been tried⁶ but has a short half life (13 hours) and cannot be used in a dual isotope procedure with technetium-99m ($^{99\text{m}}\text{Tc}$) for background subtraction. Two further problems with iodine isotopes are that they are not retained in cells and iodination of proteins is too time consuming for a nuclear medicine department.

Indium-111 (^{111}In) has been suggested⁷ as an alternative radiolabel for the following reasons: (a) the gamma energies (171 and 247 keV) are better detected by conventional gamma-cameras; (b) the half life (2.83 days) is more suitable for antibody imaging; (c) it is known to be chemically stable within cells⁸; and (d) protein labelling can be conveniently and simply carried out using a bifunctional chelating agent.⁹ Disadvantages of using ^{111}In are, firstly, that the available isotope has a lower effective specific activity than iodine and, secondly, that it is complex to prepare chelate linked proteins with enough groups available to bind ^{111}In in large amounts. A chelate that has been used for ^{111}In binding is diethylenetriamine-penta-acetic acid (DTPA), which has been successfully linked to albumin¹⁰ for use as a blood pool reagent and to anti-myosin¹¹ for imaging canine myocardium. Recently a monoclonal antibody to carcinoembryonic antigen (CEA) radiolabelled with ^{111}In via DTPA has been shown successfully to localise to human tumours transplanted into nude mice.¹²

We describe the preparation and labelling of an antiserum to CEA with ^{111}In to high specific activity and its use for locating tumours in 11 patients. Five of the patients were also scanned using the same antibody labelled with ^{131}I .

Methods and patients

ANTISERUM

Polyclonal antiserum to CEA was prepared in sheep by immunising with CEA extracted from hepatic metastases of a colonic carcinoma. Non-specific antibodies were adsorbed on glutaraldehyde polymers of

Department of Immunology, University of Birmingham, Birmingham B15 2TJ

D S FAIRWEATHER, BA, MRCP, honorary senior medical registrar
A R BRADWELL, MRCP, MRCPATH, senior lecturer
P W DYKES, MD, FRCP, consultant physician
A T VAUGHAN, PHD, lecturer
S F WATSON-JAMES, BSC, technician

Department of Nuclear Medicine, Queen Elizabeth Hospital, Birmingham B15 2TH

S CHANDLER, MSc, senior physicist

Correspondence to: Dr A R Bradwell, Department of Immunology, Medical School, University of Birmingham, Birmingham B15 2TJ.

and subtraction was necessary to enhance the image. In contrast, ^{111}In gave a clear, positive scan five days after injection (fig 1 (c)), and at nine days the tumour was still visible without subtraction. Table II gives the ratios of tumour to normal tissue for these scans, and even when these ratios were equal (day 1) the ^{111}In scan was visually superior, as assessed by the count rate per pixel in the tumour (table II). In the example in table II the subtraction scans with the ^{131}I antibody contained 100 counts at 24 hours and 189 counts at 48 hours over the whole tumour (2.2 and 5.2 standard deviations respectively above the background counting error). In the ^{111}In subtraction image the count rates were 226 at 24 hours and 825 at 48 hours—that is, 2.6 and 10.6 standard deviations respectively above the background counting error.

FIG 1—Unsubtracted lateral view of head and neck in patient with squamous carcinoma in region of right temporomandibular joint. (a) ^{131}I -anti-carcinoembryonic antigen (CEA) 24 hours after injection. (b) ^{111}In -anti-CEA at 24 hours after injection. (c) ^{111}In -anti-CEA five days after injection.

TABLE II—Count rates from scans of one patient given ^{111}In and ^{131}I labelled antibody

		Days after antibody injection			
		1	2	5	9
Ratio of tumour to normal tissue	^{111}In	2.1	2.7	3.1	1.8
	^{131}I	2.1	1.9	*	*
Counts/min/pixel	^{111}In	25.4 (25.4)†	24.1 (30.9)	11 (29.3)	7.4‡ (21.5)
	^{131}I	9.2	7.0	*	*

*Not scanned.

†Counts in parentheses corrected for physical decay.

‡Both ^{111}In photopeaks counted.

RADIOLABELLING OF ANTIBODY

Radiolabelling with ^{111}In had an overall efficiency of 50–75%, giving a maximum specific activity of 74 MBq (2.0 mCi)/mg protein. After separation on a G25 column, 100% of the counts were shown to be protein bound by filtration on Sephadex G75 and 90% precipitable by saturated ammonium sulphate at pH 3.0.

Overall titre of the antiserum, as assessed by reverse radial immuno-

diffusion, was unaffected by either ^{111}In or ^{131}I labelling, but the binding assay showed a reduction in high affinity binding sites in the DTPA linked antiserum (fig 2).

FIG 2—Binding of ^{125}I -carcinoembryonic antigen (CEA) to unmodified and radiolabelled anti-CEA. Each point is mean of three experiments for each antibody. Range of each value about 6% for unmodified and ^{113m}In labelled antibody and 9% for ^{131}I labelled antibody.

DISTRIBUTION AND DOSIMETRY OF LABELLED ANTIBODIES

In the first 24 hours after injection 33% (range 24% to 46%) of the injected ^{131}I activity was recovered in the urine and in the second 24 hours 20% (range 13% to 28%) was recovered. With ^{111}In labelled antibodies 10% (range 4.3% to 15%) of the injected activity was recovered in the urine in the first 24 hours and 3% daily thereafter (up to seven days).

In patients given oral iodide and perchlorate neither ^{131}I nor ^{99m}Tc accumulated in any organ except the bladder. In contrast, 20% of the injected dose of ^{111}In accumulated in the liver and remained there for the duration of the study (eight days). This was the only organ significantly to accumulate ^{111}In , and in particular there was little activity in the skeleton.

Whole body doses were the same for both isotopes, about 0.135 mSv (13.5 mrem)/MBq. Liver doses were 0.154 mSv/MBq for ^{131}I -anti-CEA administered, 0.783 mSv/MBq for ^{111}In -anti-CEA, and 0.089 mSv/MBq for ^{99m}Tc -colloid injected. Average bladder dose was 0.803 mSv/MBq from ^{131}I . (Conversion: 1 Sv = 100 rem.)

Discussion

These results show that ^{111}In -anti-CEA detected more tumour sites and with greater certainty than did ^{131}I -anti-CEA. Overall sensitivity of the ^{131}I scans (53%) was lower than in some reports,^{1,2} but this varies with the criteria used for staging. Furthermore, different groups of patients may not be comparable, and in our study comparisons were made within patients using the same dose of each isotope. The amount of isotope injected, however, was lower than used by some workers.¹ One patient received 66 MBq (1.8 mCi) but the others received between 18.5 and 27.7 MBq (0.5–0.75 mCi). Improved sensitivity was not shown simply by the excess of positive sites with ^{111}In , but when lesions were detected by both isotopes ^{111}In had more ++ scores, and when individual scans were compared the ^{111}In counts reached higher statistical significance.

This improvement may largely be related to improved count rates. There was some improvement in ratios of tumour to normal tissue, which may reflect different in vivo handling of the two isotopes. Nevertheless, a clear distinction must be made between increased label in the tumour and a reduction in blood pool activity. Our results suggest that most accumulation occurs in the first 48 hours; but the label remains there, and further

improvements in ratios are the result of blood clearance (see half life corrected figures in table II). This is in contrast with ^{131}I , which is less stable in tumours, presumably due to internalisation and deiodination.¹²

On cursory inspection ^{111}In might be thought to offer a great advantage over ^{131}I in terms of dosimetry because of the absence of energetic β^- particles. ^{111}In , however, produces several low energy x rays and electrons which, combined with its longer biological half life, offset this advantage. Critical organ dose from ^{111}In was to the liver, and from ^{131}I to the bladder. We emphasise that the dosimetry calculation was based on a small excretion of ^{111}In but a large excretion of ^{131}I . Since individual patients may vary widely in their rate of iodine excretion some may receive a higher dose from ^{131}I than that calculated. In addition, it is important to ensure adequate blockade of the thyroid and stomach with iodide and perchlorate.

The doses used in our study were cautious, and the British Institute of Radiology guidelines¹⁸ would allow 300 MBq (8.1 mCi) of either isotope to be administered to such patients for clinical indications (50 mSv (5 rem) whole body and 250 mSv single organ). Radiation dose with computed tomography is about 10 mSv (1 rem) per examination, and some other x ray techniques, especially fluorography, give comparable doses.

With regard to detector efficiency and spatial resolution, our high energy collimator has an efficiency of 63% compared with the medium energy collimator.¹⁹ When counting only the 247 keV ^{111}In peak there is a detector efficiency factor of 3.2 in favour of ^{111}In , reducing to 2.3 after 48 hours; if both ^{111}In peaks are included the factors are 7.8 and 5.6 respectively. Spatial resolution at 10 cm from the collimator face is 20 mm full width half maximum for ^{131}I and 17 mm full width half maximum for ^{111}In .¹⁹

With regard to the antiserum, the total binding sites were not affected by either labelling procedure, but incorporating up to five molecules of DTPA had a measurable effect on affinity profile. The substitution is chemically similar to that employed with the Boulton and Hunter reagent, and damage might be expected to be related to the number of groups substituted per molecule of IgG.⁷ Further refinements to the technique may enable reagents of equally high specific activity to be prepared without measurable effects on affinity, and, once synthesised, DTPA linked antibodies are easy to radiolabel. This has an advantage over iodination for a routine nuclear medicine department.

There have been several suggestions to improve antibody scans, of which increasing the number of counts emanating from the tumour is fundamental.²⁰ Theoretical considerations and our results show that with ^{111}In this is possible because of its physical characteristics and prolonged residence in tumours. This results in improved count rates and improved signal to noise ratios, which allow the detection of smaller or deeper lesions.

We thank Mr R Hawker for providing indium isotopes and useful discussion, Dr N Baggett for help with the chemical synthesis, and Dr R Flinn for help with computer analysis of the scans. S F Watson-James is supported by a grant from the Endowment Fund of the Central Birmingham Health District, and D S Fairweather by a project grant from the CRC.

References

- 1 Goldenberg DM, Deland F, Kim E, *et al.* Use of radiolabelled antibodies to carcinoembryonic antigen for the detection and localisation of diverse cancers by external scanning. *N Engl J Med* 1978;**298**:1384-8.
- 2 Fairweather DS, Bradwell AR, Watson-James SF, Dykes PW, Chandler S, Hoffenberg R. Detection of thyroid tumours using radio-labelled anti-thyroglobulin. *Clin Endocrinol (Oxf)* 1983;**18**:563-70.
- 3 Goldenberg DM, Kim EE, Deland FH. Human chorionic gonadotrophin radioantibodies in the radioimmunodetection of cancer and the disclosure of occult metastases. *Proc Natl Acad Sci USA* 1981;**78**:7754-8.
- 4 Halsall AK, Fairweather DS, Bradwell AR, *et al.* Localisation of malignant germ-cell tumours by external scanning after injection of radiolabelled anti-alpha-fetoprotein. *Br Med J* 1981;**283**:942-4.

- 5 Fairweather DS, Bradwell AR, Dykes PW. Monoclonal antibodies for in vivo localisation. *Lancet* 1982;iii:660.
- 6 Epentons AA, Britton KE, Mather S, *et al.* Targeting of iodine-123 labelled tumour-associated monoclonal antibodies to ovarian, breast, and gastrointestinal tumours. *Lancet* 1982;iii:999-1006.
- 7 Eckman WC, Paik CH, Reba RC. Radiolabelling of antibodies. *Cancer Res* 1980;**40**:3036-42.
- 8 Thakur ML, Segal AW, Louis L, Welch MJ, Hopkins J, Peters J. Indium-111-labelled cellular blood components: mechanism of labelling and intracellular location in human neutrophils. *J Nucl Med* 1977;**18**:1020-4.
- 9 Sunberg MW, Mears CF, Goodwin DA, Diamanti CI. Selective binding of metal ions to macromolecules using bifunctional analogs of EDTA. *J Med Chem* 1974;**17**:1304-7.
- 10 Wagner SJ, Welch MJ. Gallium-68 labelling of albumin and albumin microspheres. *J Nucl Med* 1979;**20**:428-33.
- 11 Khaw BA, Fallon JT, Strauss HW, Harber E. Myocardial infarct imaging of antibodies to canine cardiac myosin with indium-111-diethylene triamine pentaacetic acid. *Science* 1980;**209**:295-7.
- 12 Stern P, Hagan P, Halpern S, *et al.* The effect of the radiolabel on the kinetics of monoclonal anti-CEA in a nude mouse-human colon tumor model. In: Mitchel MS, Oettgen HF, eds. *Hybridomas in cancer diagnosis and treatment*. New York: Raven Press, 1982:245-53.
- 13 Egan ML, Lautenschlager JE, Coligan JE, Todd CW. Radioimmune assay of carcinoembryonic antigen. *Immunochemistry* 1972;**9**:289-99.
- 14 Krejcarek GE, Tucker KL. Covalent attachment of chelating groups to macromolecules. *Biochem Biophys Res Comm* 1977;**77**:581-5.
- 15 Watson-James SF, Fairweather DS, Bradwell AR. A shielded sterile apparatus for iodinating proteins. *Med Lab Sci* 1983;**40**:67-8.
- 16 Snyder WS, Ford MR, Warner GG. Estimates of specific absorbed fractions for photon sources uniformly distributed in various organs of a heterogeneous phantom. *MIRD Pamphlet*. No 5 (revised). New York: Society of Nuclear Medicine, 1978.
- 17 Flemming JS. A technique for the absolute measurement of activity using a gamma camera and computer. *Phys Med Biol* 1979;**24**:176-80.
- 18 British Institute of Radiology. Guidelines. Irradiation of human subjects for medical research. *Bulletin of the British Institute of Radiology* 1975; No 1.
- 19 Anger HO, Davis DH. Gamma-ray detection efficiency and image resolution in sodium iodide. *Review of Scientific Instruments* 1964;**35**:693-7.
- 20 Rockoff SD, Goodenough DJ, McIntire KR. Theoretical limitations in the immunodiagnostic image of cancer with computed tomography and nuclear scanning. *Cancer Res* 1980;**40**:3054-8.

(Accepted 25 April 1983)

ONE HUNDRED YEARS AGO The police-courts have lately made public several instances of revolting cruelty suffered by little children in the course of their training for performance as acrobats. It appears that, in the present stage of our civilisation, the supply of acrobats is a steady public demand, and that many British children of very tender years are regularly devoted to the acrobatic business, to be systematically trained, while their limbs and joints are yielding, by a long process of painful exercises enforced by punishment and privation, for the gratification of a debased public taste by their performances as "human serpents," tumblers, and contortionists. Mr Charles Mylne Barker, who acted as honorary solicitor in the proceedings taken some time ago by Mr Littler, QC, and his friends, regarding a number of young English children found living in slavery, or quasi-slavery, at Constantinople, writes to the *Times* to express his disappointment that a short Act of Parliament has not been passed during the recent session, making it illegal for any infant child of twelve years of age or under to be apprenticed to any trade or business without the consent of a magistrate. Mr Barker states that, with the assistance of the authorities of the Criminal Investigation Department, he had made several inquiries as to the manner in which infant children being trained as acrobats were treated when out of the jurisdiction of English courts; and it was found that, in many instances, the treatment was harsh in the extreme, especially when the children did not take kindly to their calling; and it was believed that a child was actually killed in Spain, its back having been broken by the cruel treatment to which it was subjected. Mr Barker thinks that some philanthropic member of Parliament, at a time when so much public sympathy is expended on the sufferings of pigeons, might have carried through a short Bill such as he has indicated. (*British Medical Journal* 1883;iii:440.)