PRACTICE OBSERVED

587

Overcoming Isolation

... by being there

ROGER JONES

The transition from teaching hospital nephrology to general practice was certain to be isolating. I would be excluded from the current of ideas that flows through our centres of excellence. I would no longer enjoy the badinage around the seminar room or the *New England Journal* review slip, and, direct of all, I would be cut off from "real" medicine. "Soon you will have forgotten the world and the world will have forgotten you."

I am delighted to report to the contrary; general practice has opened doors that I did not know existed and given the lie to the dogma that I had never doubted. The most vivid of these is the notion fostered by Lord Moran that hospital medicine requires the ablest and most dedicated of us and that general practice can make do with the less gifted and less certain. Two other misapprehensions go hand in hand with this one, in education, that information can only flow downwards from the hospital to the general practitioner, and the other that to be reputable general practice research must conform to the narrow archetypes of the respectable journals.

Of course, there are aspects of working in a close knit hospital community, where care and concern are shared, that are lost forever in general practice, and the opportunities for interchange of ideas may be restricted, even when working among sympathetic and concerned colleagues. None the less the new general practitioner principal may be heartened to learn that merely by pursuing the four main activities of primary care he will be able to participate and given the opportunity to make a contribution that would have been diluted in hospital. These four activities are the service commitment, performance review, research, and education. Every general practitioner's Venn diagram representing these four dimensions will be different, but for many of us there is much overlap among the four.

The Health Centre, Charlton Road, Andover, Hampshire SP10 3LD
ROGER JONES, MRCGP, MRCP, general practitioner

Service

The service commitment is by far the largest chunk of the general practitioner's work. "Service commitment"—how this bureaucratic phrase obscures the richness and infinite variety that we find in these seemingly endless, uncomfortable, brief encounters with our patients. There is a danger, in becoming oversophisticated, that we might fail to "capture and hold the excitement of the encounter and to explain to ourselves and others the value and meaning of what we are really being asked to do." By extending this fundamental commitment, however, we may break away from the one to one surgery confrontation; in my own practice special interests have resulted in establishing an open access endoscopy service for general practitioners in the town, training ambulance personnel in electrocardiographic diagnosis and cardiac defibrillation, establishing criteria for safe obstetrics in the community hospital, and developing family counselling and hypnotherapy for some patients. All of these endeavours make us look outside the surgery and force us to discuss our problems and learn from others with more experience and knowledge. Wherever there is a general practitioner hospital there will be a hospital staff committee, which is often a forum and a crucible for ideas and projects. In the wider setting it is possible to participate in the Association of General Practitioner Hospitals, which serves the interests of general practitioners throughout Britain. In the political arena participation in the local medical committee and district management team are ways in which we can influence local health stimulation, and it also provides a valuable input of information and representation for the local medical community.

Performance review

Performance review—a less threatening term than audit—may be reflex and informal for some, but for most of us it has to be a conscious act of will and also has to contain some objective elements. Performance review may operate on many

588

levels from personal through practice review to the participation of larger numbers of doctors. In the past year in the practice we have looked critically at antibiotic prescribing in otitis media, prescribing costs in hypertension and the rheumatic conditions, the use of the community hospital, and electrocardiographic diagnosis in patients with cardiovascular symptoms. All of these exercises have given us valuable insights and have often led to ghosts or corrected misconceptions about our behaviour and that of our colleagues. Perhaps the most interesting was the performance review which has concerned a whole medical community was a study performed with our practice trainee last year when 17 of the 18 general practitioners in the town took part in a prospective survey of outpatient referral. This engendered an enormous amount of interest and remarkable cooperation and candour about the way that the outpatient departments were used and the motives and expectations attached to such referrals.

Research

Research activities follow almost naturally from performance review, and analysis of the outpatient referral study is beginning to yield some interesting results about what determines referral, the characteristics of general practitioners that influence referral, and what use is made of secondary care generally. Other opportunities for research are constantly available; it is possible to undertake useful studies "in isolation," these may be financed, for example, through facilities of the Royal College of General Practitioners or by funds obtained from pharmaceutical companies. For example, I conducted a survey of the use of chaperones by doctors in the south of England by using funds obtained from several companies that manufacture contraceptive pills. This brought me into contact not only with the pharmaceutical industry but with academics in community medicine and statisticians in the medical school at Southampton. Another research activity was a clinical trial of single dose antibiotic chemotherapy in urinary tract infections, which meant participation with workers in antimicrobial chemotherapy. A further project that opened up wider perspectives was holding an Epithel Travelling Fellowship, which enabled me to study the resuscitation ambulance services in England, travelling to several distant centres to see medical and paramedical personnel.

Some general practitioners would undoubtedly like to express their "organised curiosity" more formally; what resources are available and what is needed to guide us in our projects? The Royal College of General Practitioners has three important roles here. Firstly, the facilities give grants and advice to members interested in research. Secondly, at regional level it has research units. Thirdly it has library facilities at its headquarters in London. The big college meetings also provide limited opportunities to present results. Locally, however, the postgraduate library of the district general hospital is a valuable resource; advice and encouragement should also be available in departments of general practice in the medical schools. An important role of departments that have been adequately established is to initiate and coordinate projects with non-academic general practitioners. The General Practitioner Research Club is a rather evanescent body that theoretically provides a forum to examine ideas for projects and preliminary results as well as polished data. The contribution of all these has for me been far exceeded by the support and warmth of a few young general practitioners, mostly from London, Oxford, and Hampshire, who meet in a small group started by one of the lecturers in primary medical care at Southampton.

Education

Finally, participating in education is guaranteed to produce a sense of belonging to a community—a tradition, and of passing on and continuously reviewing and reassessing the

values and knowledge of that tradition. General practitioners may participate at many levels. We have fifth year medical students from the University of Southampton and London medical schools, and one of the partners teaches in the third year seminars in the department of primary care at Southampton. We are also a training practice; the requirements for trainees ensure that they and anyone teaching in a university department of general practice are anything but isolated from thought in primary care. Having young people in the practice is invariably stimulating and often makes us question our preconceptions and look critically at our behaviour.

Just as important is our response to the need for our own postgraduate education. Although it is favoured by some more and more general practitioners find a diet of consultant stodge based in a postgraduate centre and predigested clinical review in glossy journals unpalatable and look to innovative educational strategies that are firmly based in general practice. Perhaps a lot of the activities of a single practice is unrepresentative of general practice in a wider setting, but it does show that any one of several activities is available and guarantees the doctor protection from isolation. We need not have unscrambled every patient's hidden agenda or developed a field theory of medicine; all we have to do is recognise the special nature of the work that we are doing, respond to its excitement, question the ways that we behave as general practitioners, and be willing to share our experiences, our doubts, and our certainties with our colleagues in medicine. Isolation may be overcome simply by being there.

References

- Higgs R. Three challenges for the future. *Br Med J* 1983;286:1113-4.
- Harrison L. L. Lushfield. Academic support practice. *Br Med J* 1982;284:872.
- Accepted 27 May 1983.

ONE HUNDRED YEARS AGO With alarming frequency we have lately heard of exhausted railway servants failing to keep awake at their posts. Last week, one of the chief and busiest railroads in the kingdom, a disaster was imminent, because an engine-driver and his fireman fell asleep on their engine, in front of the Irish bridge. We are informed that an engine had taken an excursion train to Bangor, and was returning with a train of empty carriages to Chester. Just at midnight the train left Llandudno Junction all right, and should have passed Colwyn some minutes afterwards, but, as half an hour elapsed and it was not signalled, the officials became alarmed. The Irish mail train from London to Holyhead was due to pass Llandudno Junction about one o'clock, but, happily, the driver saw the lights against him and stopped his train. An inspector went down the line, and found the empty excursion train at a standstill on the rails. The driver and fireman were fast asleep on their engine, and the fire in the fire-box was nearly out. Had not the black system been in perfect operation an appalling accident might have resulted, as the mail train runs at great speed from Holyhead to Chester, without stopping. It is reported that the driver and fireman who fell asleep misrepresented at Bangor the time they had been on duty, or they would not have been allowed to proceed on their return journey. This striking instance again makes clear the peril to which the travelling public is exposed from the risk of exhausted railway servants succumbing to fatigue at their posts. The legislature has long ago limited by stringent enactments the hours of toll in our factories, but it still leaves the work of the railway servant unrestricted. If railway companies were liable to legal process for overworking their servants, and not merely for proved damages arising in consequence of such overwork, and if efficient care were taken to render misrepresentation impossible on the part of railway servants, as to the time they have been on duty, it is not improbable that railway "accidents" might become less frequent. With increasing traffic on our railways, and with increasing speed in trains, the points to which we have referred urgently demand general and complete revision and safe adjustment. (*British Medical Journal*, 1883;ii:24.)

589

Practice Research

Screening in general practice: implications for cost and extra work

JOHN F GRACE

The report of the Royal College of General Practitioners on *Prevention of Arterial Disease in General Practice* suggested that a major but little performed task in general practice was the prevention of disease, especially coronary artery disease. We therefore decided to start a screening clinic in our practice to detect risk factors for the development of coronary artery disease that would run in parallel with the well woman cervical cytology screening clinic. We hoped to be able to evaluate the response of the population at risk, the rate of positive findings, and the increased workload and cost of screening.

Method

The practice has 11 000 patients in four geographically discrete villages in north Kent. My surgery has a list of 3525 patients. We decided to set up the two screening clinics at this surgery. All men between the ages of 35 and 40 were identified from the age-sex register and invited to attend the screening clinic. A practice nurse was recruited partly to run the clinic. A similar invitation to attend was sent to all women between the ages of 20 and 60 who apparently had not had a cervical smear in the previous five years.

Two questionnaires were drawn up for the nurse to use to identify the risks for both groups—family history, blood pressure, exercise, smoking history, height, weight, and urine analysis; for men—fasting blood cholesterol concentrations; and for women—cervical smear and breast examination. When all the men had been screened we compared the results from the two groups.

Results

Of 110 men invited, 72 attended the clinic (65%), whereas of 152 women invited only 52 attended (34%). Fig 1 shows the number of risk factors: 15 men had none, 21 had one, 16 had two, 14 had three, four had four, and one had five associated with the development of coronary artery disease. The equivalent rates for women are also shown.

RISK FACTORS (FIG 2)

Family history.—There was a positive family history of coronary artery disease in parents or siblings in 33 men and 25 women. Of the men with a positive family history, only four had high fasting blood cholesterol concentrations and in two of the men the positive family history was the only risk factor identified.

Exercise.—Twenty five men and 28 women admitted to taking no regular exercise, although "walking" was regarded as an acceptable exercise.

JOHN F. GRACE, MB, MRCP, general practitioner
Correspondence to: Ainstock, Tansard Hill, Shore, Kent DA12 3EN

Smoking.—Twenty four men and 19 women smoked more than the equivalent of five cigarettes daily.
Weight.—Thirty two men and 23 women were more than 10% above the medium weight for their height according to the charts in appendix 1 of the college report.¹

FIG 1—Number of risk factors in patients screened

FIG 2—Percentage of patients with each risk factor

Blood pressure.—The upper level of "normal" was set at 145/90 mm Hg, given the age of the group to be screened. Thus five men and two women needed follow up.

Urine analysis.—Only one woman had a trace of glycosuria needing investigation.

Fasting total blood cholesterol concentrations.—These were carried out on the men only, of whom three decided against the blood test.

590

Of the remaining 69, five (7%) had concentrations above the accepted upper limit of normal.

Cervical cytology breast examinations.—Neither produced positive results in the women, although the breast examination was more of an educational than diagnostic procedure.

OTHER FACTORS

Extra work for the doctor.—For most risk factors identified—weight, smoking, and exercise—advice and literature were provided by the nurse or the dietitian attached to the practice. The nurse will also follow up those identified in these groups. Those who had high blood pressure, glycosuria, or hypercholesterolaemia were followed up by the doctor. These totalled 13 patients, roughly one a month, and a negligible addition to the doctor's workload.

Cost.—The estimated cost is given in the table. After allowing for 70% reimbursement for the nurse's salary and the extra income generated by the extra cervical smears for which it was possible to claim, the whole screening exercise was almost self-supporting. We found that for our population of 3200 patients roughly five hours of the nurse's time was needed, firstly to overcome the backlog of patients who required screening when the clinic was set up, then to screen those entering the defined age groups at risk, and to follow up those found to be at risk before.

Estimated cost of the screening clinic

Practice nurse paid at 3.45 an hour	£20.00
Screening cost of 100 patients	£10.00
100 cervical smears at £0.10 each	£10.00
100 blood pressure readings	£5.00
100 cholesterol readings	£10.00
100 urine analyses	£10.00
100 breast examinations	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00
100 weight interviews	£10.00
100 blood pressure interviews	£10.00
100 cholesterol interviews	£10.00
100 urine analysis interviews	£10.00
100 breast examination interviews	£10.00
100 family history interviews	£10.00
100 exercise interviews	£10.00
100 smoking interviews	£10.00