- ⁵ Feinstein DI, Rappaport SI. Acquired inhibitors of blood coagulation. *Progress in Hemost Thromb* 1972;1:75-9.
- ⁶ Mueh JR, Herbst KD, Rappaport SI. Thrombosis in patients with the lupus anticoagulant. *Ann Intern Med* 1980;92:156-9.
- ⁷ Tan EM, Cohen AS, Fries JF, et al. The 1982 revised criteria for the classification of SLE. *Arthritis Rheum* 1982;25:1271-7.
- ⁸ Proctor RR, Rappaport SI. The partial thromboplastin time with kaolin. A simple screening test for first stage plasma clotting factor deficiencies. *Am J Clin Pathol* 1961;36:212-9.
- ⁹ Catterall RD. Biological false positive reactions and systemic disease. In: Walker G, ed. *Ninth symposium on advanced medicine*. London: Pitman Medical, 1973.
- ¹⁰ Moore JE, Lutz WB. The natural history of systemic lupus erythematosus: an approach to its study through biologic false reactors. *J Chronic Dis* 1955;1:297-316.
- ¹¹ Dubois EL. *Lupus erythematosus*. 2nd ed. Los Angeles: University of California Press, 1974.
- ¹² Byron MA. The clotting defect in SLE. *Clin Rheum Dis* 1982;8:137-51.
- ¹³ Schleider MA, Nachman RL, Jaffe EA, Coleman M. A clinical study of the lupus anticoagulant. *Blood* 1976;48:499-509.
- ¹⁴ Veltkamp JJ, Kerkhoren P, Loeliger EA. Circulating anticoagulant in disseminated lupus erythematosus. *Haemostasis* 1974;2:253-9.

- ¹⁵ Thiagarajan P, Shapiro SS, DeMarco L. Monoclonal immunoglobulin M λ coagulation inhibitor with phospholipid specificity—mechanism of a lupus anticoagulant. *J Clin Invest* 1980;66:397-405.
- ¹⁶ Johansson EA, Lassus A. The occurrence of circulating anticoagulants in patients with syphilitic and biologically false positive antilipoidal antibodies. *Ann Clin Res* 1974;6:105-8.
- ¹⁷ Laurell AB, Nilsson IM. Hypergammaglobulinemia, circulating anti-coagulant and biologic false positive Wassermann reaction. *J Lab Clin Med* 1957;49:694-707.
- ¹⁸ Green D. Circulating anticoagulants. *Med Clin North Am* 1972;56:145-51.
- ¹⁹ Carreras LO, Defreyn G, Machin SJ, et al. Arterial thrombosis, intra-uterine death and lupus anticoagulant: detection of immunoglobulin interfering with prostacyclin formation. *Lancet* 1981;i:244-6.
- ²⁰ McVerry BA, Machin SJ, Parry H, Goldstone AH. Reduced prostacyclin activity in SLE. *Ann Rheum Dis* 1980;39:524-5.
- ²¹ Firkin BG, Howard MA, Radford N. Possible relationship between lupus inhibitor and recurrent abortion in young women. *Lancet* 1980;ii:366.
- ²² Gonyea L, Herdman R, Bridges RA. The coagulation abnormalities in systemic lupus erythematosus. *Thrombosis et Diathesis Haemorrhagica* 1968;20:457-64.

(Accepted 28 June 1983)

SHORT REPORTS

Ascites in breast cancer

Metastatic breast cancer may manifest as serous effusions. Pleural effusions are a common complication of breast cancer, and recent studies have shown that they may be alleviated effectively by early chemical pleurodesis.¹ Patients treated in this way have a mean survival time of 16 months from presentation.² To determine whether similar relief might be obtained in patients with the less common complication of ascites, we reviewed patients under the care of this unit who developed ascites during the decade 1972-81; we established the incidence of this complication, the clinical features of the patients, and the duration of survival after presentation with ascites.

Patients, methods, and results

We reviewed the records of 56 patients with histologically confirmed breast cancer who developed clinical evidence of ascites. In 11 (20%) of them ascites had been the first manifestation of metastatic disease. The mean survival time was 5.7 months (range 0-48 months). An arbitrary division was made at three months: 38 (68%) of the patients with ascites died within this time (short term survivors), 25 (45%) dying within one month. The short term and longer term survivors were compared to establish clinical features at presentation that might identify potential longer term survivors and establish the most appropriate treatment.

There was no difference in age at diagnosis, clinical stage, disease free interval, and ascites free interval between the two groups. Similar proportions of both groups had undergone oophorectomy or received cytotoxic chemotherapy, or both.

The table shows the pattern of metastatic disease. Eleven (61%) of the longer term survivors compared with only eight (21%) of the short term survivors had previously had skin metastasis ($\chi^2 = 8$, $p < 0.005$). The proportions of patients with nodal and bone metastasis and pleural effusion were similar in the two groups. Hepatomegaly or jaundice was present in 60% of the short term survivors and in only 11% of the longer term survivors ($\chi^2 = 12.06$, $p < 0.001$). This suggests that palpable liver enlargement or jaundice indicates a bad prognosis.

Pattern of metastatic disease among 56 patients with ascites. Figures are numbers (%) of patients

Symptoms and previous metastases and treatment	Longer term survivors (n = 18)	Short term survivors (n = 38)
Ascites first symptom	2 (11)	9 (24)
Hepatomegaly or jaundice	2 (11)	23 (60)
Previous disease:		
Lymph node	2 (11)	4 (11)
Skin	11 (61)	8 (21)
Bone	6 (33)	22 (58)
Pleura	5 (28)	5 (13)
Previous treatment:		
Oophorectomy	5 (28)	7 (18)
Chemotherapy	4 (22)	16 (42)

Of the 25 patients with hepatomegaly or jaundice, 23 (92%) died within three months. Necropsy was performed in four patients who died early; in each the liver was almost completely replaced by tumour deposits. In contrast, laparotomy in three of the longer term survivors showed peritoneal seedlings without evidence of liver metastases. The usefulness of hepatomegaly or jaundice as a prognostic factor is not altered by additional information on previous skin metastasis.

Various treatments were used among both short and longer term survivors. This suggests that it is hepatic metastases rather than any particular treatment that determines survival.

Comment

These findings may have implications for the management of patients with ascites secondary to breast cancer. Of those patients who have associated hepatomegaly or jaundice, 90% will be dead within three months, probably as a result of liver failure, and will gain no benefit from active local treatment of their ascites. In contrast, the longer term survivors will emerge from those patients in whom hepatomegaly and jaundice are absent. These patients with supposed peritoneal metastases may benefit from local treatment provided that they have adequate hepatic function, as assessed by radio-isotopic scanning and blood biochemistry tests, and do not have overt disease elsewhere requiring systemic treatment.

Each of the longer term survivors was treated empirically, and ascites was controlled in one third of cases. A similar response rate would be predicted for patients with advanced breast cancer given endocrine treatment or cytotoxic chemotherapy. We do not know whether specific local treatment such as intracavitary adriamycin³ or peritoneovenous shunts⁴ gives any greater benefit than systemic treatment alone. A multicentre controlled trial would be needed to answer this question.

¹ Fentiman IS, Rubens RD, Hayward JL. Control of pleural effusions: a randomised trial. *Cancer* 1983;52:151-3.

² Fentiman IS, Millis RR, Hayward JL. Pleural effusion in breast cancer: a review of 105 cases. *Cancer* 1981;47:2087-92.

³ Kefford RF, Woods RL, Fox RM, Tattersall MHN. Intracavitary adriamycin nitrogen mustard and tetracycline in the control of malignant effusions. *Med J Aust* 1980;2:447-8.

⁴ Leveen HH, Christoudias G, Moon IP, Luff R, Falk G, Grosberg S. Peritoneo-venous shunting for ascites. *Ann Surg* 1974;180:580-91.

(Accepted 2 June 1983)

ICRF Breast Cancer Unit, Guy's Hospital, London SE1 9RT

I S FENTIMAN, MD, FRCS, consultant surgeon

R D RUBENS, MD, MRCP, consultant physician

J L HAYWARD, FRCS, director

Correspondence to: Mr I S Fentiman.