PRACTICE OBSERVED

Overlapping General Practice

Clinical psychologist

D W A JERROM, R J SIMPSON

Since 1945 clinical psychology has become a distinct discipline. Hitherto, general practitioners may have had contact with the educational psychologist through the school medical service. The clinical psychologist is one of many practical therapists who are becoming part of the network of community psychiatric care and will have an honours degree in psychology and postgraduate professional training of two or three years, including a probationer course. The specialised application of the knowledge and techniques of psychology to both mental and physical health problems adds a further dimension to treating and managing many conditions.

Ten or 20 years ago there was little overlap between clinical psychology and general practice. The psychologist before this time had a subordinate role and worked in an institution, providing psychological testing for all branches of psychiatry. During the past two decades clinical psychology has become an independent profession treating a greater variety of problems throughout the National Health Service, most recently in community primary care and away from the large institutions. A consequence of these changes in role and function, discussed in the Tretlow report,¹ which has been generally welcomed by psychologists, is that more and more the general practitioner will have a new resource person to refer his patients to, often in the health centre or surgery. The published surveys, investigations, and evaluation of the new role of clinical psychologists generally conclude that it is a useful and fruitful development in health care for the patient, the general practitioner, and the clinical psychologist.²⁻⁴

Forth Valley Health Board
D W A JERROM, MSc, principal clinical psychologist and clinical research fellow, University of Stirling
R J SIMPSON, MSc, MSc, general practitioner and psychiatrist

Correspondence to: Dr D W A Jerrom, Community Clinical Psychology Service, Department of Psychology, University of Stirling, Stirling FK9 4LA.

Our experience

In January 1979 the Forth Valley Health Board provided funding over three years for a community clinic to provide a clinical psychology service to the 78 general practitioners in one of the health board districts and to the sector psychiatrists and hospital specialists. The clinic was run from one health centre but patients were seen in other surgeries and health centres throughout the district. It was hoped that patients who either might not have had an opportunity for psychological treatment or who would have been referred to the psychiatric services when psychological treatment would have been more appropriate would be referred.

A steering committee, chaired by Professor H Barber, Norrie-Miller Professor of General Practice at Glasgow University, closely monitored and evaluated the project over the three years. An important feature was the management group, consisting of a psychiatrist and a general practitioner working with the two psychologists who provided the service, which was able to allay the fears of both general practitioners and psychiatrists that the independence of the clinical psychologist would create difficulties and helped doctors and psychologists to learn about the skills and limitations of the other. From the outset clinical responsibility was held according to the recognised professional guidelines⁵ and these have proved to be acceptable. The psychologist was responsible for his own methods of treatment while the general practitioner retained primary responsibility for the overall health care of the patient. By the end of the three years 698 patients had been referred—21000 a year for the district. Referrals had been received from 77% of the general practitioners. Those who had not referred were practising predominantly in rural areas where no "on the spot" service could be offered regularly.

The distribution of diagnoses in patients referred was as follows: 70% suffered from an anxiety state or phobia (32% generalised anxiety, 38% agoraphobia, social phobia, or specific phobias), 8% depression, 7% sexual deviation or dysfunction, and the remaining 15% a mixture of psychiatric syndromes, such as obsessional compulsive disorders and

anorexia, habit disorders, or relationship problems. Although the problems tended to be chronic, with a mean length of seven years' presentation, only a third of cases had received psychiatric or other specialist attention before. A questionnaire sent to the general practitioners who referred the first 30 cases showed that in the absence of such a service they would have continued treating two thirds of the referred cases themselves. We have received only three inappropriate referrals, and this low level was undoubtedly helped by a well attended introductory one day seminar. Patients were usually treated individually with either behaviour therapy or cognitive therapy. Treatment was usually given for seven or eight sessions averaging 20 to 30 minutes each at intervals of two to four weeks. A few severe cases were treated for much longer. The psychologist always wrote to the general practitioner about the patient after assessment and after treatment was concluded and sent interim progress reports if treatment was protracted.

We have evaluated the outcome of psychological treatment.⁶ In summary, we found a high level of satisfaction both among patients and general practitioners with the service, and we showed statistically significant reductions in drug usage, consultation rates with the patient's doctor, and severity of self-rated problems after treatment and at six months' follow up. Patients who fared best were young, were not taking medication at the outset, and had less intercurrent physical disease; but there was no relation between diagnosis and outcome. Furthermore, it was often difficult to predict outcome for individuals until four or five treatment sessions had elapsed, which is common with behaviour therapy and other psychological treatments. A major question that remains to be answered is whether in the long term early intervention with psychological treatment will have an appreciable effect on the high consultation rates noted among the few patients with chronic neurotic problems. If indeed the clinical psychologist can relieve the general practitioner of this particularly frustrating aspect of his work, which accumulates with each succeeding year in practice, then his contribution will be doubly valuable.

Further roles for the clinical psychologist in primary care

Some potentially fruitful roles for psychologists in primary care are other than direct concern in patient management: patient education, staff training, and stimulation of patient self help groups. By patient education we mean working with target patient groups in the practice and not general health education programmes, which are the responsibility of health education departments. For example, we are planning brief courses for adolescents with diabetes who have maintenance problems and on non-drug treatment for patients with mild and moderate essential hypertension. Such programmes will require careful design and evaluation; if proved effective, however, they could be provided in future by the general practitioner, community nurse, or paramedical staff.

Staff training covers teaching medical staff, community nursing staff, health visitors, and counsellors about psychological techniques for changing behaviour and attitudes. For example, teaching general relaxation therapy and techniques for

managing insomnia might give general practitioners further opportunities for curtailing the use of minor tranquilisers. In our survey of general practitioners in our health district 60% of them were interested in such training, but opinions on the best method of providing it were equally divided between seminars on general topics and the opportunity to discuss specific cases in the practice with the psychologist.

Self help groups may be used as a resource by general practitioners. We have helped to set up a group for people who suffer from both phobias and depression. Contributions might also be made in antismoking groups or weight watchers' groups working in collaboration with health visitors. It is important to evaluate the effectiveness of the psychologist's contribution to self help groups before either psychologists or other health care staff make a major commitment to them.

The clinical psychologist has a growing part to play as a member of the primary care network treating and managing psychiatric and psychological morbidity in the community. Linking in one practice the general practitioner, health visitor, nurse, and social worker with specialists such as the clinical psychologist, psychiatrist, community psychiatric nurse, specialist social worker, and trained counsellor should lead to more effective patient care. We are fortunate that the Forth Valley Health Board has supported our experimental project and has established the community psychologist service permanently. The most important aspect is that specialist treatment was available to patients who previously would not have received it. We have emphasised repeatedly the need for careful evaluation since we believe that such experiments too often become part of the service without effective evaluation. After five years we think that the value of the community psychologist has been proved in our district, although further work needs to be done to confirm these findings in the long term. There is already sufficient evidence to warrant the addition of a clinical psychologist to the other National Health Service specialists available in the community.

References

1. Simpson RJ. The team: an outmoded concept? *Scottish Medicine* 1981; 1:15-6.
2. Tretlow WH. The role of psychologists in the health service. London: HMSO, 1977.
3. McPherson I, Feldman MP. A preliminary investigation of the role of the clinical psychologist in the primary care setting. *Bulletin of the British Psychological Society* 1977; 30:342-6.
4. Eastman C, McPherson I. An outline of the general practitioners' perceptions of psychological problems and the relevance of clinical psychology. *Br J Clin Psychol* 1982; 51:226-31.
5. Jerrom DWA, Simpson RJ, Barber JH, Pemberton DA. General practitioners' satisfaction with primary care clinical psychology service. *J R Coll Gen Pract* 1983; 33:26-31.
6. Koch HGH. Evaluation of behaviour therapy intervention in general practice. *J R Coll Gen Pract* 1979; 29:337-41.
7. Ives G. Psychological treatment in general practice. *J R Coll Gen Pract* 1979; 29:341-51.
8. Clark DF. The clinical psychologist in primary care. *Soc Sci Med* 1979; 13:707-13.
9. Jerrom DWA, Garver D, Simpson RJ, Pemberton DA. Clinical psychology in primary care—issues in the evaluation of services. In: Mann CJ, ed. *Clinical psychology and medicine*. London: Pitman, 1982.
10. Bart L, Kinney J. Clinical psychology in general practice: a controlled trial evaluation. *J R Coll Gen Pract* 1982; 32:32-7.

Practice Research

Is the serological diagnosis of infectious mononucleosis always necessary?

D O HO-YEN

Abstract

Clinical and serological data from 100 patients who had a Downey lymphocytosis of $\geq 40\%$ of all white cells were assessed over three years. All had clinical evidence of infectious mononucleosis, suggesting that Downey lymphocytosis of $\geq 40\%$ is specific for the illness.

Introduction

Infectious mononucleosis is believed to be a result of Epstein-Barr virus infection, and the disease has distinctive features.¹ The diagnosis is usually not made unless a triad of clinical, haematological, and serological findings are present. The characteristic lymphocyte in infectious mononucleosis has been called "atypical"; however, as these cells may total up to 12% of all white cells in normal patients,² this description is imprecise. The historical term "Downey lymphocytes," in recognition of their detailed illustration by H Downey in 1923,³ is more appropriate. Recently it has been suggested that a haematological finding of Downey lymphocytes $\geq 40\%$ of all white cells is sufficient to diagnose infectious mononucleosis.⁴ This would allow clinicians in certain cases to make the diagnosis without serological evidence, and there would be a saving in each case of at least 57% in the cost of making the diagnosis.⁴

Patients, methods, results

Prospectively, over three years, the clinical and serological data were assessed in 100 consecutive patients in whom there was a Downey lymphocytosis of $\geq 40\%$ of all white cells. To reduce interobserver and intraobserver variation in counting Downey lymphocytes time was spent ensuring that there was concordance in identifying the types of Downey lymphocytes among the haematologists. Then, if at least 200 white cells are counted, the mean coefficient of variation is about 5%. Sera were examined for heterophile antibody by the Paul-Bunnell-Davidsohn test; any serum that did not contain heterophile antibody was tested for antibodies to Epstein-Barr virus capsid antigen, cytomegalovirus, and *Toxoplasma gondii*. Methods were performed as described.^{5,6}

Results of the serological tests were: 81 sera were positive for the Paul-Bunnell-Davidsohn test, 15 had IgM antibodies to Epstein-Barr virus capsid antigen, and four were undiagnosed. IgG antibodies to Epstein-Barr virus capsid antigen were found in all patients with IgM antibodies and in none of the undiagnosed patients. The undiagnosed

Haematology Section, University Department of Pathology, Ninewells Hospital and Medical School, Dundee
D O HO-YEN, MSc, MSc, MD, senior registrar

Correspondence to: Regional Virus Laboratory, Ruchill Hospital, Glasgow G20 9NB.

patients were tested three weeks after the start of the illness; two became positive for the Paul-Bunnell-Davidsohn test and two remained undiagnosed. The table gives the results of haematological investigations. All patients had clinical evidence of infectious mononucleosis.⁷ Tests showed no differences among the three groups of patients in cervical lymphadenopathy, sore throat, fever, malaise, and fatigue.

Mean (SD) of total white cell counts, lymphocyte counts (and percentage), and Downey lymphocyte counts (and percentage) in the three groups of patients

Patients	Total white cell count ($\times 10^9/l$)	Lymphocytes ($\times 10^9/l$)	Downey lymphocytes ($\times 10^9/l$)
(1) Positive for the Paul-Bunnell-Davidsohn test (n=81)	12.6 (3.5)	69.5 (8.5)	8.8 (2.8)
(2) IgM antibodies to Epstein-Barr virus capsid antigen (n=15)	11.3 (3.7)	72.7 (9.3)	8.3 (3.1)
(3) Undiagnosed (n=4)	6.6 (2.4)	62.5 (10.3)	4.3 (2.0)
Remained undiagnosed (n=2)	12.1 (2.6)	70.0 (7.1)	6.6 (2.6)

Discussion

Patients with clinical evidence of infectious mononucleosis do not always have haematological and serological features of the disease. This is particularly true in general practice where patients are often seen early in the course of the disease. Diagnosis may be further complicated by the haematological criteria required. Low concentrations of Downey lymphocytes are not significant, and a concentration $\geq 20\%$ of all white cells is required.⁴ In patients with such concentrations, and undiagnosed evidence of infectious mononucleosis serological investigations have shown that 69% have the illness, 5% have toxoplasmosis, 2% have cytomegalovirus, but 24% remain undiagnosed.⁴ This high level of undiagnosed cases probably reflects the fact that most patients were from general practice.

The degree of Downey lymphocytosis can be useful in the investigation of patients with clinical infectious mononucleosis. Patients with toxoplasmosis, cytomegalovirus, and undiagnosed infections have lower concentrations of Downey lymphocytes, usually in the range 20-25%.⁴ Adults and children with these illnesses have similar concentrations of Downey lymphocytes, though children with infectious mononucleosis may have exceptionally high concentrations.⁸ This study was designed to verify the suggestion that all patients with $\geq 40\%$ Downey lymphocytes have serological evidence of infectious mononucleosis.⁴ The only requirement for entry into the study was $\geq 40\%$ Downey lymphocytes, yet I found that all the patients had clinical evidence of mononucleosis. Although serological confirmation was obtained in only 98 patients, the remaining two

patients had identical haematological and clinical features. As some patients may form antibodies to Epstein-Barr nuclear antigen without forming other antibodies⁹ the two undiagnosed patients may have had antibodies to Epstein-Barr nuclear antigen for which they were not tested. Thus massive Downey lymphocytosis seems to be specific for infectious mononucleosis. The results of one survey of patients suspected of having this illness showed that in more than a third of cases the referring

doctor (mostly general practitioners) would have had to base the diagnosis on the amount of Downey lymphocytes alone.¹⁰ The results from my study show that this is a reasonable approach (figure). Apart from the concentration of Downey lymphocytes, the haematologist may also consider other factors that vary during the course of the illness such as leucopenia, neutropenia, and thrombocytopenia.¹¹ In patients with clinical and serological evidence of infectious mononucleosis $\geq 40\%$ have $\geq 40\%$ Downey lymphocytes.⁴ But in patients with only clinical features of infectious mononucleosis 15% have $\geq 40\%$ Downey lymphocytes⁴; this probably reflects the large number of diseases that can produce the clinical picture of infectious mononucleosis. Thus knowledge that $\geq 40\%$ Downey lymphocytes is specific for the disease obviates further investigation and saves much time and money.

I am grateful for the help and cooperation of Dr A J Munro, Dr N Saleem, Dr R A Bennet, Dr R A Sharp, Dr A S Todd, and Dr H B Goodall.

References

1. Ho-Yen DO. The significance of Downey lymphocytes and a serum inhibitor in the clinical picture of infectious mononucleosis. Dundee: Dundee University, 1981. (MD thesis).
2. Wood TA, Franklin EP. The atypical lymphocyte. *Am J Med* 1967; 43: 923-36.
3. Downey H, McKinlay CA. Acute lymphadenitis compared with acute lymphatic leukaemia. *Arch Intern Med* 1923; 32:12-12.
4. Ho-Yen DO. Diagnosis of infectious mononucleosis. *Br Med J* 1980; 280: 1338-9.
5. Ho-Yen DO, Martin K. The relationship between atypical lymphocytes and serological tests in infectious mononucleosis. *Journal of Infection* 1981; 3:324-31.
6. English EC, Geyman JP. The efficiency and cost effectiveness of diagnostic tests for infectious mononucleosis. *J Fam Pract* 1978; 8:977-81.
7. Gervais F, Jonck JH. An unusual antibody response to EBV during infancy. *J Infect Dis* 1979; 140:273-4.
8. Ho-Yen DO, Pinkerton LW, Reid D. Infectious mononucleosis. *Update* (in press).
9. (Accepted 24 August 1983)

Laboratory and radiological investigations in general practice

III: Abnormal results

K A MILLS, P M REILLY

Results

Throughout the study practice C had a higher rate of requests for investigations. We wanted to see if this led to the detection of appreciably more patients with abnormalities compared with the other practices. To allow a fair comparison the numbers of patients in whom abnormalities were detected by each practice during

the six months 1 September 1982 to 28 February 1983 are expressed as a percentage of the number of surgery encounters in that practice over the same period (table 1). We confirmed by means of the χ^2 test that the rate at which patients were investigated was significantly different among the four practices over the six months (see column 3). If practices A and B are considered alone, however, there was no significant difference between them, but there was a pronounced difference among practices A, B, and D. Of the patients investigated by each practice, a significantly different proportion of them had abnormalities (column 5). This difference is lost if practice C is omitted and the calculations repeated; thus practice C seems to have had a higher proportion of patients with abnormalities in its population of investigated patients than the other three practices. The four practices also differed significantly in the proportion of surgery encounters that yielded one or more abnormalities as a result of laboratory or radiological investigations. Again, practice C had the highest proportion, followed by D, and then A and B (column 6).

Department of General Practice, The Queen's University of Belfast
K A MILLS, MSc, research assistant
P M REILLY, MSc, MSc, senior lecturer

Correspondence to: Mrs K A Mills, Department of General Practice, Dundee Health Centre, 1 Dundee Avenue, Belfast BT9 7TH, Northern Ireland.